

**PHARMACOGENOMICS OF CHEMOTHERAPEUTIC AGENTS:
CARBOPLATIN AND PACLITAXEL**

A Dissertation
SUBMITTED TO THE FACULTY OF
UNIVERSITY OF MINNESOTA
BY

Taraswi Mitra Ghosh

IN PARTIAL FULFILLMENT OF THE REQUIREMENTS
FOR THE DEGREE OF
DOCTOR OF PHILOSOPHY

Dr. Jatinder K. Lamba

May 2017

ACKNOWLEDGEMENTS

At this special moment when I have reached the concluding stages of my Doctoral research and am ready to embark upon another voyage in my scientific career, I would like to take the opportunity to acknowledge the contributions of the people who have shaped up my life and career.

First and foremost, I would like to thank Almighty God for giving me the strength to overcome all the challenges, for giving me the motivation to learn and to acquire knowledge.

I would like to thank Dr. Jatinder Lamba, my adviser. I have learned a lot from her and am extremely thankful to her. She has helped me develop my scientific temper and inquisitive thinking.

I would like to express my sincere gratitude towards my committee members. Dr. Mark Kirstein, my committee chair, has been a constant support. He has helped me develop my concepts in Pharmacokinetics. His insightful questions during our discussions have encouraged me to think deep and broaden my knowledge base. Dr. Angela Birnbaum, my committee member and DGS, has always been a strong support. Working with her was a very nice and cherishable experience for me. She has provided me the confidence to undertake independent projects- an experience which I am sure will be helpful in my future as a Scientist. I would like to thank Dr. Natalia Tretyakova for her invaluable suggestions and helpful guidance in Drug metabolism.

I am grateful to my collaborators at the University of Kansas (Dr. Brooke Fridley and Dr. Rama Raghavan), Minneapolis VA Healthcare (Dr. Pankaj Gupta) and the Mayo Clinic (Dr. Ellen Goode) for their support during my research.

My heartiest thanks to Prof. Esam El-Fakahany, Prof. Brian Van Ness and Dr. Robert Straka for standing by me when things were not on brighter side. It was an honor for me to be associated with them throughout my scientific quest.

I am extremely grateful to all my professors and teachers who have enriched my knowledge and understanding of various fields. Especially, I would like to mention Dr. Demerath, Dr. Foster, Dr. Zimmerman and Dr. Susan Telke in University of Minnesota and Prof. Ashish Kumar DuttaGupta, Prof. Trilochan Midya, Prof. Arun Chatterjee, Dr. Nihar Ranjana Chatterjee, Dr. Srikanta Kumar Rath, Atanu Banerjee, Nandita Chatterjee, and Cindy Brown for their contributions.

I would like to acknowledge the help of all the members of ECP administrative staff Carol Ann, Dede, Karla, Mary, Erin for their help whenever I needed.

I am very grateful to Jan Morse for providing the support and confidence I needed during difficult times in my PhD career.

My special thanks to Tanya Feldberg for being such a wonderful person through her continued support and encouragement. I would like to thank Dr. Neha Bhise for being an excellent lab member and above all a great friend and confidante. I am also thankful to Dr. Lata Chauhan for her help and for being a cordial host in Florida.

I would like to thank my friends at University of Minnesota Manoj, Kinjal, Sam, Youssef, Mariam, Chay, Ritu, Shuang, Ashwin, Vaishnavi and University of Florida Miyoung and Bradley who have shared my journey through my graduate student life.

I am extremely thankful to all my friends Arpita, Shahida di, Nlanjana di, Bubai da, Subhrajit, Ujjal da, Sohan, Debarchana, Smita, Moumita, Joyeeta, Sraboni di, Rita di and Kathak dance buddies. Without their support my journey would have been impossible to complete.

I consider myself lucky to have very supportive parents. My parents have taught me that education is the door to freedom. I will always be grateful to them for instilling in me the urge to acquire knowledge. I realized during my PhD that they are willing to sacrifice anything for my education when they decided to leave their country and live with me during my final graduation year.

I have heartfelt love for my loving sister, Tanushri, who has always unconditionally believed in me. I always remember and admire all her support right from our childhood. I would also like to thank Abhijit, my brother-in-law, who takes care of my parents in my absence.

I am extremely grateful to my in-laws and my extended family. As I am writing these acknowledgements I feel sad that my father-in-law, who passed away last month, will not be here to share the joy of my PhD degree completion with me.

I would like to thank my friend and fortunately my husband, Amit, who has been the pillar of my strength through all the trials and tribulations over the past 17 years. He is the biggest critique as well as the biggest admirer of all my work. He has forever handled my gloomy phases with immense patience. This journey was possible because of him.

Last but not the least, I would like to express my deepest love for our son, Adhrit (Tintin), the emperor of our hearts.

DEDICATION

To my beloved companion, Amit

TABLE OF CONTENTS

LIST OF TABLES	IX
LIST OF FIGURES	XI
CHAPTER 1 Literature review.....	1
1.1 Ovarian cancer	2
1.1.1 Introduction	2
1.1.2 Risk factors.....	4
1.1.3 Prognosis of Ovarian cancer	6
1.1.4 Treatment of Ovarian epithelial cancer	6
1.1.5 Ovarian cancer clinical trials using Platinum/Paclitaxel chemotherapy	8
1.2 Lung cancer: Non-small cell lung cancer (NSCLC)	11
1.2.1 Introduction	11
1.2.2 Risk Factors	14
1.2.3 Symptoms and Diagnosis	14
1.2.4 Prognostic factors	14
1.2.5 Treatment of Non-Small cell lung cancer (NSCLC)	15
1.2.6 NSCLC clinical trials using Platinum/Paclitaxel combination therapy.....	15
1.3 Platinum drugs	19
1.3.1 Introduction	19
1.3.2 Mechanism of action of Platinum drugs	20
1.3.3 Metabolic pathway of platinum drugs.....	22
1.3.4 Pharmacogenomics of platinum drug pathway genes	35
1.4 Paclitaxel chemotherapy.....	83
1.4.1 Introduction	83

1.4.2	Mechanism of action of paclitaxel	84
1.4.3	Metabolic (PK/PD) pathway of paclitaxel	88
1.4.4	Pharmacogenomics of paclitaxel (PK/PD) pathway genes	93
1.5	Research Study Objectives.....	105
1.5.1	Gaps in literature	105
1.5.2	Objectives and Specific Aims	106
CHAPTER 2 Pharmacogenomic markers influence the variation of in vitro chemosensitivity to carboplatin and paclitaxel as single agents and as combination: a pathway based approach ...		108
2.1	Introduction	109
2.2	Materials and Methods.....	111
2.2.1	Creation of EBV-transformed Ovarian cancer LCLs.....	111
2.2.2	Cell Culture.....	111
2.2.3	Drugs	111
2.2.4	In vitro cytotoxicity assays	111
2.2.5	Caspase-Glo 3/7 activity Assay	112
2.2.6	Cell cycle analysis	112
2.2.7	SNP Genotyping	112
2.2.8	RNA Isolation and Gene expression assays.....	112
2.2.9	Data analysis	117
2.3	Results.....	119
2.3.1	<i>In vitro</i> profiling of carboplatin and paclitaxel chemo-sensitivity	119
2.3.2	K-Means clustering identified resistant subgroups	123
2.3.3	Analysis of carboplatin-paclitaxel combination treatment using CI theorem	126
2.3.4	SNPs in pathway genes associated with chemo-sensitivity.....	130

2.3.5	Differential GEP signatures of drug response	136
2.4	Discussion and Conclusions	140
CHAPTER 3 Response to platinum, taxane, and combination therapy in ovarian cancer: In vitro phenotypes, inherited variation, and disease recurrence		
		144
3.1.	Introduction	145
3.2.	Materials and Methods.....	147
3.2.1.	Patients, Lymphoblastoid Cell Lines and Cytotoxicity Assays.....	147
3.2.2.	Genotyping and Statistical Methods.....	149
3.3.	Results.....	151
3.4.	Discussion and Conclusions	158
CHAPTER 4 Genetic variation in platinating agent and taxane pathway genes as predictors of outcome and toxicity in advanced non-small-cell lung cancer.....		
		163
4.1	Introduction	164
4.2	Materials and methods.....	169
4.2.1	Patients cohort.....	169
4.2.2	Treatment schedules, dose & toxicity assessments	170
4.2.3	SNP analysis:	170
4.2.4	DNA isolation	170
4.2.5	Genotyping.....	170
4.2.6	Statistical methods.....	173
4.3	Results.....	175
4.3.1	Demographic features and clinical characteristics	175
4.3.2	Treatment, response and outcomes.....	176
4.3.3	Pharmacogenetics.....	178

4.4	Discussion.....	182
4.5	Conclusion.....	185
4.6	Summary.....	186
CHAPTER 5	Summary	187
References	193
Chapter 1.....		194
Chapter 2.....		225
Chapter 3.....		227
Chapter 4.....		230

LIST OF TABLES

TABLE 1.1 DETAILS OF CLINICAL TRIALS EVALUATING PLATINUM DRUGS/PACLITAXEL COMBINATION THERAPY IN OVARIAN CANCER.	9
TABLE 1.2 SUMMARY OF CLINICAL TRIALS OF CARBOPLATIN/PACLITAXEL COMBINATION THERAPY IN NSCLC.	17
TABLE 1.3 SUMMARY OF SNPs IN DRUG TRANSPORTER GENES ASSOCIATED WITH RESPONSE TO PLATINATING AGENTS.	40
TABLE 1.4 SUMMARY OF SNPs IN DRUG METABOLISM GENES ASSOCIATED WITH PLATINUM-BASED CHEMOTHERAPY.	46
TABLE 1.5 SUMMARY OF SNPs IN NER PATHWAY GENES ASSOCIATED WITH PLATINUM CHEMOTHERAPY.	65
TABLE 1.6 SUMMARY OF SNPs IN MMR, NHEJ AND SOS PATHWAY GENES ASSOCIATED WITH PLATINUM DRUG CHEMOTHERAPY OUTCOME.	75
TABLE 1.7 SNPs FOUND SIGNIFICANTLY ASSOCIATED WITH PLATINUM DRUG CHEMOTHERAPY OUTCOME IN GWAS STUDIES	82
TABLE 1.8 SUMMARY OF SNPs IN EFFLUX-INFLUX TRANSPORTER AND DRUG METABOLIZING (PK) GENES ASSOCIATED WITH PACLITAXEL CHEMOTHERAPY.	97
TABLE 1.9 SUMMARY OF SNPs IN MICROTUBULE BINDING, APOPTOSIS AND IMMUNE RESPONSE (PD) GENES ASSOCIATED WITH PACLITAXEL CHEMOTHERAPY.	102
TABLE 1.10 SNPs FOUND SIGNIFICANTLY ASSOCIATED WITH PACLITAXEL DRUG CHEMOTHERAPY OUTCOME IN GWAS STUDIES.	104
TABLE 2.1 GENES INCLUDED IN TAQMAN LOW-DENSITY ARRAY (TLDA) CARD.	115
TABLE 2.2 SUMMARY OF <i>IN VITRO</i> PHENOTYPE MEASURES IN LCLS FROM EOC PATIENTS.	120
TABLE 2.3 SUMMARY OF CORRELATION BETWEEN CELL VIABILITY & CELL CYCLE PHASES AND DRUG IC ₅₀	123
TABLE 2.4 SUMMARY OF DRUG RESPONSE PHENOTYPES IN SUBGROUPS IDENTIFIED BY KMEANS CLUSTERING. ...	126

TABLE 2.5 CI AND DRI AS MEASURES OF SYNERGISM AND ANTAGONISM OF DRUG COMBINATION TREATMENT IN LCLs FROM EOC PATIENTS.....	129
TABLE 2.6 TOP PATHWAY SNPs FOUND MOST SIGNIFICANTLY ASSOCIATED WITH <i>IN VITRO</i> CYTOTOXICITY PARAMETERS.	132
TABLE 2.7 TOP PATHWAY SNPs FOUND SIGNIFICANTLY ASSOCIATED WITH <i>IN VITRO</i> COMBINATION TREATMENT CASPASE ACTIVITY.	134
TABLE 2.8 SUMMARY OF RESULTS FROM SPEARMAN CORRELATION ANALYSIS BETWEEN TLDA GENE EXPRESSION AND DRUG CHEMO-SENSITIVITY OUTCOMES ($P < 0.1$).....	137
TABLE 3.1 MEAN (SD) <i>IN VITRO</i> PHENOTYPE MEASURES COLLECTED ON LCLs FROM 74 EOC PATIENTS AND COMPARISON ACROSS EXPERIMENTAL BATCH.....	148
TABLE 3.2 HAZARD RATIOS (P-VALUE) ASSOCIATING <i>IN VITRO</i> DRUG PHENOTYPES WITH TIME TO DISEASE RECURRENCE.....	152
TABLE 3.3 SNPs WITH $P < 10^{-6}$ ASSOCIATION WITH A DRUG RESPONSE PHENOTYPE.	153
TABLE 3.4 GENE REGIONS WITH SNPs ASSOCIATED WITH BOTH PHENOTYPES FOR A GIVEN DRUG.	156
TABLE 3.5 GENE REGIONS WITH SNPs ASSOCIATED WITH MULTIPLE DRUGS FOR ANY PHENOTYPE ($P < 0.0001$).	157
TABLE 4.1 SUMMARY OF SNPs GENOTYPED IN THE NSCLC PATIENTS.	172
TABLE 4.2 PATIENT DEMOGRAPHICS AND CLINICAL CHARACTERISTICS (N = 90).	175
TABLE 4.3 CHEMOTHERAPY REGIMENS, DOSE INTENSITY, RESPONSE AND SURVIVAL.	176
TABLE 4.4 HEMATOLOGICAL AND NON-HEMATOLOGICAL TOXICITIES.	177
TABLE 4.5 ASSOCIATION OF SNPs WITH PROGRESSION-FREE SURVIVAL AFTER ADJUSTING FOR COVARIATES (PERFORMANCE STATUS AND STAGE).	180
TABLE 4.6 ASSOCIATION OF SNPs WITH TOXICITY AFTER ADJUSTING FOR COVARIATES (AGE AS CATEGORICAL VARIABLE AND TOTAL NUMBER OF CYCLES).....	181

LIST OF FIGURES

FIGURE 1.1 A) INCIDENCE OF OVARIAN CANCER IN THE US, 1992-2012. B) NEW CASES, 2008-2012, ALL RACES, FEMALES (NCI SEER CANCER STATISTICS).....	3
FIGURE 1.2 OVERALL TREATMENT STRATEGIES FOR EPITHELIAL OVARIAN CANCER (EOC)	10
FIGURE 1.3 INCIDENCE AND MORTALITY OF LUNG CANCER; NON-SMALL CELL LUNG CANCER (NSCLC) (NCI SEER CANCER STATISTICS).	13
FIGURE 1.4 NSCLC TREATMENT SUMMARY BY STAGE (NCCN GUIDELINES; VERSION 1.2016).	18
FIGURE 1.5 CHEMICAL STRUCTURE OF PLATINUM DRUGS.....	19
FIGURE 1.6 MECHANISM OF ACTION OF PLATINUM DRUGS.....	21
FIGURE 1.7 METABOLIC PATHWAY OF PLATINUM DRUGS	27
FIGURE 1.8 MISMATCH REPAIR (MMR) PATHWAY.	30
FIGURE 1.9 NUCLEOTIDE EXCISION REPAIR (NER) PATHWAY.....	33
FIGURE 1.10 GENETIC VARIANTS IN GSTP1 ASSOCIATED WITH SURVIVAL/OUTCOME AND/OR TOXICITY IN PATIENTS TREATED WITH PLATINUM DRUG-BASED CHEMOTHERAPY (GREEN BOXES DENOTE CODING SNPs).	48
FIGURE 1.11 GENETIC VARIANTS IN ERCC1 ASSOCIATED WITH SURVIVAL/OUTCOME AND/OR TOXICITY IN PATIENTS TREATED WITH PLATINUM DRUG-BASED CHEMOTHERAPY (GREEN BOX DENOTES CODING SNP; YELLOW BOXES REPRESENT NON-CODING SNPs).....	64
FIGURE 1.12 A) BASIC STRUCTURE OF TAXANES SHOWING DIFFERENCES IN PACLITAXEL AND DOCETAXEL (R^1 AND R^2) ⁵³¹ B) MAJOR METABOLITES OF PACLITAXEL WITH THE ENZYMES INVOLVED ⁵⁵³	86
FIGURE 1.13 MECHANISM OF ACTION OF PACLITAXEL.....	87
FIGURE 1.14 METABOLIC PATHWAY OF TAXANES.....	92
FIGURE 2.1 WORKFLOW REPRESENTING DATA QC (QUALITY CHECK) OF CHEMO-SENSITIVITY ASSAY RESULTS...	119

FIGURE 2.2 BOXPLOTS REPRESENTING DISTRIBUTION OF A) DRUG CYTOTOXICITY AND B) CASPASE ACTIVITY FOLLOWING TREATMENT.....	121
FIGURE 2.3 SCATTERPLOT MATRIX REPRESENTING PAIRWISE CORRELATIONS BETWEEN IC ₅₀ AND AUC ESTIMATES OF PACLITAXEL AND CARBOPLATIN SINGLE-AGENT AND COMBINATION TREATMENT.....	122
FIGURE 2.4 RESULTS FROM KMEANS CLUSTERING OF <i>IN VITRO</i> CYTOTOXICITY DATA	124
FIGURE 2.5 PLOTS SHOWING DIFFERENCES IN RESPONSE TO CARBOPLATIN, PACLITAXEL AND COMBINATION TREATMENTS BETWEEN THE CLUSTERS GENERATED USING <i>IN VITRO</i> CYTOTOXICITY ASSAY DATA ON THE LCL PANEL.....	125
FIGURE 2.6 REPRESENTATION OF VARIATION OF THE RELATIVE IC ₅₀ S OF TREATMENT WITH SINGLE AGENTS AND DRUG COMBINATION THAT INDICATES VARIATION ACROSS LCL PANEL. A) CARBOPLATIN; B) PACLITAXEL. .	128
FIGURE 2.7 SCATTERPLOTS SHOWING CORRELATION BETWEEN COMBINATION INDEX (CI) AND DOSE REDUCTION INDEX (DRI) VALUES AT IC ₅₀	129
FIGURE 2.8 HEATMAP SHOWING HIERARCHICAL CLUSTERING OF THE LCL PANEL BASED ON DIFFERENTIALLY EXPRESSED CARBOPLATIN PATHWAY GENES. A) ALL PATHWAY GENES; B) PATHWAY GENES MOST SIGNIFICANTLY ASSOCIATED WITH CYTOTOXICITY (P<0.1).	138
FIGURE 2.9 HEATMAP SHOWING HIERARCHICAL CLUSTERING OF THE LCL PANEL BASED ON DIFFERENTIALLY EXPRESSED PACLITAXEL PATHWAY GENES. A) ALL PATHWAY GENES; B) PATHWAY GENES MOST SIGNIFICANTLY ASSOCIATED WITH CYTOTOXICITY (P<0.1).	139
FIGURE 3.1 SCATTERPLOT MATRIX OF DRUG RESPONSE PHENOTYPES BY EXPERIMENTAL BATCH.....	149
FIGURE 3.2 KAPLAN-MEIER CURVES FOR TIME TO PROGRESSION AND PACLITAXEL <i>IN VITRO</i> PHENOTYPES.	152
FIGURE 3.3 MANHATTAN PLOTS OF THE SINGLE SNP META-ANALYSIS GENOME-WIDE ASSOCIATION RESULTS BASED THE SIX <i>IN VITRO</i> DRUG RESPONSE MEASURES AMONG OVARIAN CANCER PATIENT LCLS.	154
FIGURE 3.4 LOCUS ZOOM PLOTS FOR REGIONS WITH $P < 10^{-6}$	155

FIGURE 3.5 PATHWAY ANALYSIS OF GENES WITH SNPs THAT DEMONSTRATED SIGNIFICANT ASSOCIATION WITH LCL CHEMOSENSITIVITY WITH CARBOPLATIN OR PACLITAXEL ALONE OR IN COMBINATION.....	161
FIGURE 4.1 PLATINUM PHARMACOKINETICS/PHARMACODYNAMICS PATHWAY	166
FIGURE 4.2 TAXANE PHARMACOKINETICS PATHWAY.....	168
FIGURE 4.3 PATIENT SELECTION AND STUDY DESIGN.	169

CHAPTER 1 LITERATURE REVIEW

1.1 Ovarian cancer

1.1.1 Introduction

The ovaries are the most important part of the female reproductive system ¹. Each ovary is divided into an outer cortex and an inner medulla ¹. The surface of the ovary is covered by a single layer of cuboidal epithelium, called germinal epithelium¹.

Ovarian cancer is a type of cancer that begins in the ovaries and often goes undetected until it has spread within the pelvis and abdomen ². It is ninth most common cause of cancer death among women in the United States. According to the Cancer Facts & Figures published by the American cancer society (ACS), 21, 290 (12.1 per 100,000 women per year) new cases and 14,180 deaths (7.7 per 100,000 women per year) due to ovarian cancer were registered in 2015 in the United States. Five year survival rate was 45.6% (**Figure 1.1**) ³. Broadly, tumors of the ovaries are classified into three major categories, i) Ovarian epithelial cancer, ii) Ovarian germ cell tumor and iii) Ovarian low malignant potential tumor.

Epithelial ovarian cancer (EOC) is one of the most common gynecological malignancies and the fifth leading cause of cancer death among women in the United States (6% of cancer deaths) ⁴. About 9 out of 10 tumors of the ovary diagnosed (90%) are of this type. EOC occur primarily in middle aged or older women while it is rare in young adults, especially before puberty ⁵. The standard treatment for patients with advanced disease is initial debulking surgery followed by carboplatin-paclitaxel combination chemotherapy. Five-year overall survival (OS) remains around 44.2%, suggesting there is a wide inter-patient variation in response ⁶.





Figure 1.1 a) Incidence of ovarian cancer in the US, 1992-2012. b) New cases, 2008-2012, All Races, Females (NCI SEER cancer statistics)

1.1.2 Risk factors

Factors like genetic mutations, hormonal therapy, age, nulliparity, obesity, and fertility drugs are associated with increased risk of ovarian cancer. Whereas oral contraceptives, multiple pregnancies, breast feeding and tubular ligation are associated with decreased risk of ovarian cancer.

- a) Genetic factors: Approximately 20% of ovarian cancer is familial and linked with BRCA1 or BRCA2 gene mutation ^{7,8}. Germline deleterious mutations in the BRCA1/BRCA2 genes are associated with 15% to 40% lifetime risk of ovarian cancer ^{9,10}. Loss of BRCA1 or BRCA2 protein expression is more common in ovarian cancer ¹¹. Prophylactic measure for this mutation is bilateral oophorectomy and it is beneficial in certain cases ¹². Other gene like TP53, PTEN/MMAC1, STK11 and mismatch repair genes MLH1, MSH2, MSH6, and PMS2, have been associated with an increased risk of ovarian cancer (NCI).
- b) Hormonal therapy: Women using more than 10 years of hormonal therapy including estrogen only, progestagen only, estrogen-progestagen combinations and tibolone are in increased risk of ovarian cancer. The relative risk (RR) of hormone therapy current users verses never users of hormone therapy was 1.20 (95% confidence interval [CI], 1.09–1.32). Risk increases with increasing duration of use ¹³. A population-based case-control study of exclusively estrogen-only therapy for at least 5 years showed odds ratio [OR], 1.6; 95% CI .A cohort study showed that use of ERT(estrogen replacement therapy)/ET (estrogen therapy) and ERT/ET followed by EPRT (estrogen –progesterone replacement therapy) more than 20 years are associated with increased risk of ovarian cancer (RR=relative risk 3.2 95%CI, 1.7-5.7) ¹⁴.
- c) Age and Nulliparity: Risk of developing ovarian cancer increases linearly from age 30 to 50 years of age and continues to increase further with age thereafter ¹⁵. Women over 55 are most commonly diagnosed with ovarian cancer. About half of the women who are

diagnosed with ovarian cancer are 63 years or older ². Furthermore, there is increased risk of ovarian cancer in older woman who have never been pregnant ^{16, 17}.

- d) Body Mass Index (BMI), Diet and Obesity: BMI (= (the weight in kgs)/ (square of height in meters)) is associated with increased mortality rate from ovarian cancer. A prospective study on more than 900,000 U.S. adults (404,576 men and 495,477 women) showed individuals with BMI ≥ 40 had higher death rates from cancers (52% for men; 62% for women) compared to 'normal' BMI (18.5 to 24.9) ¹⁸. Some studies also support the association between increased risk of cancer with height and diet, along with BMI ¹⁹⁻²¹.
- e) Oral contraceptives: Interestingly, use of Oral contraceptive is associated with reduced risk of ovarian cancer. 10%-20% reduction in ovarian cancer risk is associated with one year use of oral contraceptive, where as 50% reduce risk associated with five years use of oral contraceptive among women ^{22, 23}. Case-control (24) and cohort studies published since 2000 observed risk reduction with use of oral contraceptive more than 1 year ²⁴. A multicenter study with 330 cases and 982 matched controls, showed a statistically significant decreased risk of ovarian cancer associated with Progestational contraceptives (depot-medroxyprogesterone acetate [DMPA]) use ²⁵.
- f) Tubular ligation: Decreased risk of ovarian cancer associated with tubal ligation ^{26, 27}. Population-based case-control studies (13) showed tubal ligation is associated with a 29% reduction in ovarian cancer risk (OR, 0.71; 95% CI, 0.66–0.77) ²⁸.
- g) Breast feeding: Case control studies (30) showed breast-feeding is associated with a decreased risk of ovarian cancer (RR, 0.76; 95% CI, 0.69–0.83) ²⁹.
- h) Smoking: A meta-analysis from 51 studies with 28,114 ovarian cancer patient found a very small increased risk of ovarian cancer among current smokers compared with nonsmoker (RR, 1.06; 95% CI, 1.01–1.11) ¹³.

- i) Use of Talc: A meta-analysis of 16 studies observed an increased risk with the use of talc (RR, 1.33; 95% CI, 1.16–1.45) ³⁰. A case-control study that included 8,525 cases and 9,859 controls have proven increased risk of epithelial ovarian cancer associated with genital powder use (OR, 1.24; 95% CI, 1.15–1.33) ³¹.

1.1.3 Prognosis of Ovarian cancer

Women with low risk cancers defined as stage IA, IB, grade 1 or 2, nonclear-cell histologies do not need further adjuvant therapy. Survival of early-stage disease is significantly higher (20% to 30%) than advanced stage ovarian cancers ³². Five-year disease-specific survival is 84% for stage I compared to stage II disease, which is 76% ³². Interestingly younger women are more likely to be diagnosed with a lower-stage along with more well-differentiated tumors and have an improved outcome compared with older women. In addition, patients with a significant component of transitional cell carcinoma have a better prognosis ³³⁻³⁵. Case control studies suggest that BRCA1 and BRCA2 mutation carriers show better response compared to sporadic EOC patients ³⁶. Patients with high-risk early-stage epithelial ovarian cancer defined as stage I, grade 3; stage IC, stage II, clear-cell cancers and they require postsurgical adjuvant treatment. Additionally, late diagnosis is a major problem for ovarian cancer. Lack of awareness of patients and physicians about occurrence of nonspecific symptoms like abdominal pain and swelling, gastrointestinal symptoms and pelvic pain is another major concern. The ovarian cancer-associated biomarker CA-125 is more significant at the time of disease progression after 1-2 chemotherapy cycle in stage III and stage IV, instead of time of early stage of diagnosis ³⁷.

1.1.4 Treatment of Ovarian epithelial cancer

There are various therapies available for EOC which are selected based upon stage and tumor types. Platinum based chemotherapy is the standard-of-care for 75% of ovarian cancer patients. The

treatment of EOC is divided into three groups- Early stage, advanced stage and recurrent or persistence (as shown in **Figure 1.2**).

Standard chemotherapy worldwide for epithelial ovarian cancer are platinum agents cisplatin or carboplatin either alone or combination with cyclophosphamide or paclitaxel ³⁸⁻⁴². This regimen was evaluated and established by several clinical trials by GOG (Gynecology Oncology Group), MRC (Medical Research Council) ⁴³⁻⁴⁸. However, approximately 80% ovarian epithelial cancer patients relapse after first line platinum and taxane based chemotherapy. If CA-125 increases after first line treatment is completed, then relapse is usually suspected. ⁴⁹. Patients with recurrent disease are therefore subdivided as: i) Platinum - sensitive recurrent and ii) Platinum - refractory/resistance recurrent ⁴⁹.

Carboplatin was approved by FDA in 1987 for ovarian cancer patient who are recurrent after cisplatin therapy ⁵⁰. Currently, the most commonly used regimens for the Platinum-sensitive recurrent subtype are cisplatin or carboplatin + paclitaxel, carboplatin+ Gemcitabine, carboplatin + pegylated liposomal doxorubicin ⁵¹⁻⁵⁶. Other regimens include carboplatin + epirubicin, Cisplatin + doxorubicin + cyclophosphamide, PEGylated liposomal doxorubin + trabectedin ^{57, 58}.

If recurrence occurs within six months of completion of platinum containing therapy, it is referred to as Platinum-Refractory or Platinum-resistant Recurrence ovarian cancer. Anthracyclines (PLD), taxanes, topotecan, and gemcitabine are used as single agents for these recurrences, which are primarily contributed by platinum-resistant subsets ⁵⁹⁻⁶⁵. Paclitaxel is also commonly used in front-line induction regimens ⁶⁴. Furthermore, adverse effects of platinum-based chemotherapy in ovarian cancer as a single agent or in combinations are myelosuppression, nausea, vomiting, alopecia, and asthenia for this therapy. The most common toxicities are neutropenia, and gastrointestinal (GI) toxicity.

1.1.5 Ovarian cancer clinical trials using Platinum/Paclitaxel chemotherapy

Platinum drugs such as cisplatin plus paclitaxel combination therapy have been widely adopted as standard treatment for advanced ovarian cancer. A number of Phase I/II/ III clinical trials have evaluated the combination of carboplatin/paclitaxel chemotherapy and shown improvements in treatment outcomes in ovarian cancer patients following administration of platinating agents as chemotherapeutic agents and the use of paclitaxel as combination treatment, as summarized in **Table 1.1.**

Evidently, despite these treatment interventions, drug resistance, disease recurrence, and toxicities are major challenges for platinum-based therapy in ovarian cancer. The clinical trials presented here report that there is considerable variation in response to therapy which is a major challenge for platinum based therapy in ovarian cancer patients.

Table 1.1 Details of clinical trials evaluating platinum drugs/paclitaxel combination therapy in ovarian cancer.

Patient details	Drug response/outcome	Drug Toxicity	Pubmed ID
German Arbeitsgemeinschaft Gynäkologische Onkologie (AGO) patients (n=518)	Overall response rate=75%; Complete responses rate=41%; Partial response=34%.	Hematologic toxicity occurred more frequently than non-hematologic toxicity	9346222
Stage IV ovarian cancer patients (n=34)	Complete response rate=82%; Partial rate=95%; 1-year survival rate=94%	Nausea and vomiting grade 2=30%; Grade 2-3 gastrointestinal toxicity=20%; Grade 3-4 hematological toxicity=73%	11695811
International Federation of Gynecology and Obstetrics study on stage II to IV epithelial ovarian cancer patients (n=59)	Response rate (complete response + partial response)=72%	Quality-of-life scores improved significantly during therapy due to modest amount of toxicities.	11324769
Prospective randomized trials (phase I/II trials) by the Gynecologic Oncology Group and the European Organization for Research and Treatment of Cancer	No differences in efficacy for carboplatin/paclitaxel compared to cisplatin/paclitaxel arm	Reduce toxicity for carboplatin/paclitaxel compared to cisplatin/paclitaxel arm.	10190787
Advanced stage EOC patients (n=50)	Response rate=72%; Recurrence or progression of disease=68%		9514799
International Federation of Gynecology and Obstetrics study on Advanced epithelial ovarian cancer stages IIC, III, and IV (n=90)	patients receiving carboplatin/cisplatin + paclitaxel had a high overall response (82%).	Grade 3 and 4 neutropenia=20%; Grade 3 and 4 thrombocytopenia=4% for carboplatin group. Grade 3 and 4 neutropenia=32%; Grade 3 and 4 thrombocytopenia=7%, for the cisplatin arm	9346224
GINECO, a French cooperative clinical trials group for gynaecological cancer research (multicenter phase II carboplatin/paclitaxel study): (n=50).	Overall response rate=43%.	Grades 3 and 4 neutropenia=30%; Transitory peripheral neuropathy=45%.	9346219
Phase I study of carboplatin/paclitaxel in patients with stage IIC to IV ovarian cancer: (n=30)	Overall response rate=57%.	Grades 1 and 2 peripheral neurotoxicity=53%; patients developed Grade 4 neutropenia=31%.	9045330




Figure 1.2 Overall treatment strategies for epithelial ovarian cancer (EOC)

1.2 Lung cancer: Non-small cell lung cancer (NSCLC)

1.2.1 Introduction

In addition to ovarian cancer, platinum agents and paclitaxel are also used to treat patients diagnosed with lung cancers. This section is a brief overview of non-small cell lung cancer (NSCLC).

Lung carcinoma is the most common malignancy worldwide, and the leading cause of cancer deaths. According to the most recent estimates, the global incidence of lung cancer is more than 1.6 million cases/year, resulting in more than 1.3 million deaths/year (>18% of all cancer deaths)⁷⁴. In the USA, the incidence of lung cancer was estimated to be 221,200 cases in 2013, leading to nearly 158,040 deaths (27.5% of all cancer deaths)³. Five-year relative survival with for lung cancer patient was 15.7% (from 1995 to 2001)⁷⁵. Lung cancer can either arise directly from lung tissue or as a result of metastasis from other parts of the body. Broadly, there are two main types of primary tumor described as either small-cell or non-small-cell lung carcinomas (**Figure 1.3**).

Non-small cell lung cancer (NSCLC) is the most common type of lung cancer constituting around 80% of lung cancers. It is histologically heterogeneous and constitutes squamous cell carcinoma, large cell carcinoma and adenocarcinoma (based on data obtained from National Cancer Institute-Non-small cell lung cancer treatment). NSCLC are therefore classified by the World Health Organization (WHO)/International Association Lung Cancer (IASLC) based on histology. There are three main subtypes, i) squamous cell carcinoma (25% of lung cancers), ii) adenocarcinoma (40% of lung cancers) and iii) large cell carcinoma (10% of lung cancers). Squamous cell carcinoma is located centrally, in the larger bronchi of the lung and strongly related with smoking. Squamous cell carcinoma is further subdivided into papillary, clear cell, small cell and basaloid tumor based upon histology of tumor. Similarly, adenocarcinoma is also subdivided into acinar, papillary tumor,

solid adenocarcinoma with mucin, adenocarcinoma with mixed subtypes and bronchioloalveolar carcinoma. Adenocarcinoma has the frequent histologic heterogeneity.

The 5-year relative survival of patients with advanced NSCLC remains approximately 4 % ⁷⁶. One major drawback of lung cancer is that the majority of lung cancer patients are diagnosed in advanced stages based on small biopsy and cytology ⁷⁷. Determination of stage is very important for prognosis and selection of therapy. NSCLC staging is based on histology of tumor, resection margins (surgical margin) of it and also lymph node involvement (status and location of lymph node). Based upon these NSCLC stage is determine by TNM classification according to American Joint Committee on Cancer (AJCC) 2010, whereas T = size and extension of tumor, N = lymph node involvement and M = extension of metastasis ⁷⁸.




Figure 1.3 Incidence and Mortality of lung cancer; Non-small cell lung cancer (NSCLC) (NCI SEER cancer statistics).

1.2.2 Risk Factors

Several risk factors are associated with development of lung cancer. Major risk factors include cigarette, pipe, or cigar smoking. The risk for lung cancer is on average tenfold higher in smokers compare to nonsmokers. The risk increases with the quantity of cigarettes, duration of smoking, and starting age ⁷⁹. Exposure to second-hand smoke, radon, arsenic, asbestos, chromates, chloromethyl ethers, nickel, polycyclic aromatic hydrocarbons, radon and air pollution are also considered as potential risk factor of lung cancer ⁸⁰.

1.2.3 Symptoms and Diagnosis

The most common symptom is worsening cough or chest pain. Other presenting symptoms are hemoptysis (coughing up blood), malais (discomfort, illness), weight loss, dyspnea/ breathlessness, hoarseness (abnormal voice change). Symptoms from distant metastases can show neurological defect or personality changes from brain metastases or pain from bone metastases. Physical examination also shows supraclavicular lymphadenopathy, pleural effusion or lobar collapse, unresolved pneumonia, chronic obstructive pulmonary disease or pulmonary fibrosis ⁸¹.

1.2.4 Prognostic factors

Prognosis of disease depends on current physical status, current disease status with stage, and prior physical status. Depending on these three prognostic factors alone, median survival varies between 6 weeks and over a year. Additionally, other factors such as tumor size, histologic type, and cell type, age and sex appeared to be important when considered alone. Adverse prognosis is related with large tumor size >3 cm, histology-nonsquamous, vascular invasion and if tumor metastasize in lymph nodes. Prognosis is adversely affected by poor performance status and weight loss of more than 10% in inoperable condition patients ⁸²⁻⁸⁷.

Biomarkers including genetic mutations and gene expression are important prognostic factors for lung cancer or Non-Small Cell Lung Cancer. These include identification of significant mutations

in patient groups and strategies to molecular targeted therapy are therefore important in improving the survival of these subsets of patients ⁸⁸. Relevant genes include epidermal growth factor receptor (EGFR), Kirstein rat sarcoma viral oncogene (KRAS), Human epidermal growth factor receptor2 (HER2), V-raf murine sarcoma viral oncogene homolog B1 (BRAF), PI3K catalytic protein alpha (PI3KCA), gene rearrangements in anaplastic lymphoma kinase (ALK) as well as somatic mutations and gene duplications in MAP2K1/MEK, a hepatocyte growth factor receptor, ⁸⁹⁻⁹¹.

1.2.5 Treatment of Non-Small cell lung cancer (NSCLC)

Treatment of NSCLCs depends on staging (as shown in **Figure 1.4**). Standard treatment options are surgery, chemotherapy and radiation therapy. Platinum-based chemotherapy is the standard for NSCLC ⁹². Commonly use drug regimens are cisplatin/etoposide (EP5050) or weekly carboplatin/paclitaxel ⁹³⁻⁹⁵. Additionally, paclitaxel is also used as combination with cisplatin or carboplatin ^{92, 96, 96, 97, 97-110}. Platinum-based radiation chemotherapy (Platinum + etoposide or cisplatin + mitomycin + vindesine) is also used as a treatment option and may improve survival of patients with locally advanced NSCLC. However, toxicity is also high for combination therapy than single therapy ^{100, 111-115}. Many clinical trials have shown Cisplatin-based combinations plus radiation therapy shows 10% reduction in the risk of death compared with radiation therapy alone ¹¹⁶⁻¹¹⁹. Maintenance therapy with single agent or initial chemotherapy is recommended for patients with stable or responding disease after four cycles of platinum-based combination chemotherapy ^{101, 120, 121}.

In spite of these treatment options, 5-year survival rate of NSCLC is very poor (3% to 7%) for majority of patients ^{122, 123}.

1.2.6 NSCLC clinical trials using Platinum/Paclitaxel combination therapy

Carboplatin (PC) plus paclitaxel (Taxol) is a standard treatment of advanced non-small-cell lung cancer (NSCLC). In **Table 1.2** we evaluate the results from clinical trials performed on non-small-

cell lung cancer (NSCLC) patients using combination of paclitaxel (Taxanes) and carboplatin Chemotherapy. Similar to our observation regarding clinical trials in ovarian cancers, variations in drug response and drug-related toxicities are major cause of concerns for patients diagnosed with NSCLC undergoing platinum/paclitaxel-based combination chemotherapy. Understanding the genomics and transcriptomics governing these inter-individual variations in treatment outcome may provide tools to guide treatment and tailor therapies specific to each patient with the goal of achieving maximum therapeutic effects.

Table 1.2 Summary of clinical trials of carboplatin/paclitaxel combination therapy in NSCLC.

Patient details	Drug response/outcome	Drug Toxicity	PubMed ID
Patients with stage IIIB-stage IV NSCLC (n=90)	Overall response=40%. One-year survival=50.7%	Grades 3/4 leukopenia=13.3%,; anemia=15.5%; thrombocytopenia= 11.1%.	11863090
Stage IIIB or IV NSCLC (n=100) patients.	Objective response=36%,.	Grade 3/4 peripheral neuropathy=18%.	9007133
Advanced stage NSCLC patients (n=33).	Complete responses=41%; partial responses=41%.	Grade 3/4 esophagitis=51%.	9007133
Phase II trial at the the Fox Chase Cancer Center and its network on patients with advanced NSCLC	Response rate =55%; median event-free survival=24 weeks; median survival=47 weeks,	Peripheral sensory neuropathy=75%; myelosuppression=44% (grade 4 granulocytopenia 38% and grade >=3 thrombocytopenia 50%-70%).	9331128
Phase II trial conducted by the Eastern Cooperative Oncology Group with stage IV non-small cell lung cancer (NSCLC).	Overall response rate=27%. Median survival=38 weeks; Survival rate at 1-year=32%	Grade 3 or 4 granulocytopenia=47%; Thrombocytopenia=3%. Most common non-hematologic toxicities were nausea, emesis, neuropathy, arthralgia and myalgia.	9007120
Phase II trial in patients with advanced NSCLC (n=54)	Response rate= 50%.	Myelosuppression was observed as principal toxicity. Grade 3 or 4 granulocytopenia=70%; thrombocytopenia=13%; anemia=9%; fatigue=9%; hemorrhagic cystitis=1% after the first cycle.	7543559
Multi-center study of two dose levels of paclitaxel and carboplatin in locally advanced and metastatic non-small cell lung cancer (NSCLC) (n=130, 99 in the high dose (HD) and 31 in the low dose (LD) cohort)	Overall best clinical response rate =23.8%; one complete (CR); 30 partial responses (PR). One-year survival=34%.	Myelosuppression was the most prominent side-effect for both cohorts. Grade 3-4 leucopenia =HD 34.4% and LD 19.3%; neutropenia (HD 59.2, LD51.6%. The most prominent non-hematologic toxicities were alopecia and polyneuropathy.	11396237

Stage	Treatment	5-yr Survival
IA & IB	1) Surgery (Lung tumor and lymph node surgery) 2) Surgery not approved or declined - Radiation therapy 3) Chemotherapy if cancer still present(1B) 4) Drugs: Carboplatin, Paclitaxel	45-49%
IIA & IIB	1) Surgery → adjuvant chemotherapy +/- radiation (Chemoradiation therapy) 2) Surgery not approved or decline - Radiation / Chemoradiation therapy 3) Drugs: carboplatin, Paclitaxel	30%
IIIA (resectable)	1) Surgery → adjuvant chemotherapy /Chemoradiation therapy 2) Surgery not approved or declined - Chemoradiation therapy 3) Drugs: Carboplatin, Paclitaxel	14%
IIIB & IV (unresectable)	1) Radiation / Chemoradiation (Stage III) 2) Chemotherapy or Targeted therapy (Tyrosine-kinase inhibitor-TKI).	1-5%

Figure 1.4 NSCLC Treatment Summary by Stage (NCCN guidelines; Version 1.2016).

1.3 Platinum drugs

1.3.1 Introduction

This section provides discussion of the key agents that are the focus of this thesis. Platinum drugs are currently used to treat various types of cancers including ovarian, non-small cell lung cancer (NSCLC), testicular cancer, lymphomas and myelomas. **Figure 1.5** shows the major members of this class of platinating agents.




Figure 1.5 Chemical structure of Platinum drugs

Cisplatin was the first platinum drug to be approved by FDA. Therapeutic potential of cisplatin was discovered by Barnett Rosenberg in 1969¹³⁰. Unfortunately, cisplatin chemotherapy has dose-limiting side effects including nephrotoxicity, ototoxicity, myelosuppression and neurotoxicity¹³¹. To overcome cisplatin toxicity, the second generation platinating agent carboplatin (Diammine[1,1-cyclobutanedicarboxylato(2-)-O,O'] platinum(II); Empirical formula: C₆H₁₂N₂O₄Pt; Molar mass: 371.249 g/mol; Trade name: Paraplatin) was developed and approved by the FDA in 1989 for the treatment of advanced ovarian cancer¹³². Additionally, to reduce toxicity carboplatin is administered by a 30-minute intravenous infusion of 300 to 500 mg/m² based on a patient renal function according to Calvert Formula:

$$[\text{Total Dose (mg)} = (\text{target AUC}) \times (\text{GFR} + 25)].$$

Neurologic toxicity due to carboplatin based therapy is low/very rare (~ 5%).

However, although carboplatin is routinely used for treatment of ovarian and other cancers, one of the biggest challenges faced by clinicians is the variation in clinical response,^{133, 134} and toxicities including myelosuppression (Thrombocytopenia ~ 25%, Neutropenia ~ 16%, leukopenia ~ 16%, Anemia ~ 71%), gastrointestinal toxicity (vomiting, nausea and diarrhea ~ 65%,)¹³³ as well as development of drug resistance¹³⁴. Therefore, a large number of the cancer patients relapse after first line based chemotherapy due to wide inter-patient variation^{131, 135}.

Differential expression and/or activity of genes involved in the platinum drug pathway due to presence of single nucleotide polymorphisms (SNPs) may have an impact on treatment outcome and toxicity in cancer patients being treated with platinum compound. Here we discuss the most relevant genetic variations in the genes involved in the platinum-drug pathway and to evaluate their association with chemotherapy.

1.3.2 Mechanism of action of Platinum drugs

Platinum drugs are highly polar hydrophilic molecules. They do not easily diffuse across hydrophobic/lipophilic lipid membranes. Therefore, following intravenous administration of carboplatin or cisplatin, they remain inactive in their native form in the high chloride environment of plasma¹³¹. Upon transport of platinating agents by ion pumps and transporters, they enter the cell where, owing to a much lower concentration, the carboxyl ligands are slowly displaced by water (aquation), at the rate of 10^{13} molecules/sec, leading to the development of aqua ligands. The positively charged aqua ligands are then displaced and the platinum atom binds to purine bases on DNA forming DNA adducts as mono adducts, 1,2-intrastrand DNA cross-links (1-2GpG, 1-3GpG, 1-2GpA) and inter-strand crosslinks. These DNA crosslinks are attributed to the formation of DNA helix-distorting adducts that result in strand breakage activating DNA repair mechanisms that interfere with cell division, which ultimately results in apoptosis¹³¹. The mechanism of action of platinum drugs is shown in **Figure 1.6**.




Figure 1.6 Mechanism of action of platinum drugs.

Upon entering the cell platinum drugs lose chloride or oxalate ions, and gains two water molecules. The platinum atom binds to N⁷ of purine in the DNA forming DNA adducts as mon-adducts and through 1,2-**intra-strand** (involving adjacent guanines on the same strand) and **inter-strand** (involving binding to guanines on opposite DNA strands).

1.3.3 Metabolic pathway of platinum drugs

1.3.3.1 Genes of importance in drug pharmacokinetics (PK)

Figure 1.7 shows detailed metabolic pathway of platinum drugs including the pharmacokinetics (PK) and the pharmacodynamics (PD).

Platinum drugs enter the cell as a “passenger” along with normal copper transport by binding to the N- terminus of Copper Transporter 1 (CTR1) followed by CTR1 monomer crosslinks without blocking copper binding ^{136, 137}. The Copper Transporter 1 (CTR1), encoded by the gene Solute Carrier Family 31(SLC31), thus plays an important role in the influx of platinum drugs into the cell ¹³⁸⁻¹⁴⁰. It is a 28kDa protein consisting of 190 amino acids. It has 3 trans-membrane domains. Mammals and yeasts lacking CTR1 have been shown to have lower accumulation of platinum inside the cell. A study demonstrated that the loss of CTR1 function is associated with 10-15% decrease in cisplatin entering the cell resulting in a two fold increase in cell survival ¹⁴¹. Low or reduced expression of CTR1 has been correlated with resistance to carboplatin and cisplatin and overexpression has been associated with increased carboplatin and cisplatin uptake ¹⁴²⁻¹⁴⁶. Furthermore, studies have also shown that CTR1-knockout results in resistance to platinum drugs. CTR1 lacking cells accumulates 50% less cisplatin and eightfold higher resistance compare to the CTR1+ cells ¹⁴⁷. CTR1-/- tumor xenografts in mouse models also are not responsive to platinum drugs. Loss of CTR1 decreases the platinum drug binding to the transporter and reduces influx of cisplatin by 81%, oxaliplatin 68% and for carboplatin almost complete elimination of ¹⁴⁸influx (Larson et al. 2009). However, once inside the cell, platinum drug also initiates the rapid degradation of CTR1 ion through ubiquitination and proteosomal action mediated by copper chaperone antioxidant protein (ATOX1). Proteasome inhibitors like bortezomib have been shown to block this process of CTR1 degradation which leads to increase in the uptake and transport of platinum-containing drugs. The molecular mechanisms of platinum compound involved

endocytosis of CTR1 is not completely clear yet ^{148,149}. Some evidences show that CTR2 is involved in platinum compound/drug accumulation and that its expression correlates with the sensitivity of ovarian carcinoma cell lines to cisplatin ^{139,150}. Additionally SLC21A8 expression was significantly associated with 3 platinum-based chemotherapeutics agents such as cisplatin, carboplatin, and (trans-1-1,2-diaminocyclo-hexane)platinum-II (DACH-Pt), which is structurally related to the another platinating agent oxaliplatin ($p=0.025$) ¹⁵¹.

Copper transporting ATPase (ATP7A and ATP7B) provides efflux and resistance to platinum containing drugs along with copper transport ¹⁵²⁻¹⁵⁵. Cisplatin and other platinum drugs like carboplatin and oxaliplatin molecules bind to the first four metal N-terminal binding sites of ATPase copper transporter beta (ATP7B) transporter and stimulate catalytic phosphorylation of ATP7B, which eventually facilitated drug efflux ^{140, 155-157}. ATP7B transfection in human epidermoid carcinoma cell line KB-3-1 resulted in resistance to both cisplatin and carboplatin ^{158, 159}. *In vivo* studies in breast carcinoma ¹⁶⁰ and ovarian cancer cell lines ^{161, 162} have shown association of expression of these genes with cisplatin and/or carboplatin resistance. ATP7B is the primary candidate for platinum drug resistance where ATPase copper transporter alpha (ATP7A) play minor role. ATP7B knockdown ovarian cells show increase cisplatin sensitivity and increased efficacy with cisplatin therapy. However, the mechanism between this association is not completely clear. High ATP7B gene expression is significantly associated with cisplatin resistance in ovarian carcinoma cell lines. In 2008/MNK ovarian cancer cell lines, increased ATP7A expression is predictive of poor survival ($p<0.00057$) to the cisplatin, carboplatin and Oxaliplatin ¹⁵⁴. Additionally, in ovarian cancer cell line model ATP7B genes were expressed at higher levels in platinum-resistant cells compared with sensitive cells. ATP7B silencing showed 2.5-fold reduction of cisplatin IC₅₀, increased DNA adduct formation and reducing tumor growth along with increased tumor cell apoptosis, and reduced angiogenesis ¹⁵². However, overexpression of ATP7A and ATP7B in Me32a fibroblast resulted in increased resistance to cisplatin but not to carboplatin

¹⁶³. Homologous transporters ATP7A and B are 160-170-KDa membrane proteins with eight transmembrane and several cytosolic domains. The n-terminal cytosolic domain contain six copper binding sites where as ATP binding domain is located between transmembrane domain six and seven ^{164, 165}. ATP7A and ATP7B are primarily involved in the copper transport from cytoplasm into trans-Golgi network. As the copper content of the cell increases, ATP7A moves from trans-Golgi network to the plasma membrane while ATP7B relocates to intracellular vesicular compartments, to facilitate the export pathway ^{166, 167}.

ATP-Binding Cassette (ABC) transporter superfamily of proteins with 48 members are among most ancient and largest protein families since they have been found present in all living organisms, from prokaryotes to humans. Classification of proteins as ABC transporters is based on the sequence and organization of their ATP-binding cassette (ABC) domains ¹⁶⁸⁻¹⁷⁰. Glutathione conjugates of platinum compound can be excreted from the cells by members of the ABCC transporter family ¹⁷¹. Therefore, the most important ABC transporters studied in context of transport of platinum-containing drugs are ABCC2 and ABCG2 – which are very frequently associated with resistance of cancer cell to multiple drugs, also known as multidrug resistance (MDR) ¹⁷². ABCC2 - ATP-Binding Cassette, Sub-Family C (CFTR), Member 2 or Multidrug Resistance- Protein 2 (MRP2) and ABCG2 - ATP-Binding Cassette, Sub-Family G (WHITE), Member 2 are members of ATP-binding cassette (ABC) transporter superfamily that regulate secretion of organic anions and efflux of platinum compounds ¹³⁸. Cytotoxicity of cisplatin, carboplatin and oxaliplatin was significantly lowered in MDCKII cells transfected with ABCG2 transporter. The IC(50) values in MDCKII-ABCG2 were 25.7 for cisplatin, 164 for carboplatin and 165 micro M for oxaliplatin whereas IC(50) for MDCKII cells were 15.4 for cisplatin, 133 for carboplatin and 50.3 micro M for oxaliplatin ¹⁷³. Cisplatin-resistant melanoma cells showed a distinct overexpression of MRP2/ABCC2 on mRNA and protein level ¹⁷⁴. Platinum drug resistance Ovarian carcinoma line A2780RCIS, the adrenocortical carcinoma line D43/86RCIS and the melanoma line MeWoCIS1

overexpressed of MRP2/ABCC2. The level of DNA platination corresponded inversely to the level of MRP2 expression in these cell lines. Therefore, functional inhibition of MRP2 might be a promising treatment strategy for resistance to platinum-based anticancer drugs ¹⁷⁵. Multidrug resistance-associated protein 1 (MRP1) is a protein which is encoded by the ABCC1 gene and it play an interesting role in platinum drug resistance through the GS-X pump. The GS-X pump actively effluxes the glutathione S-platinum (GS-Pt) complex to detoxifying cells ¹⁷⁶. Additionally, ABCC4 overexpression significantly associated with decreased cisplatin sensitivity in lung cancer patients ^{138, 171}.

Several genes, such as Myeloperoxidase (MPO), Superoxide Dismutase 1 (SOD1), GSTM1, NAD(P)H Quinone Dehydrogenase 1 (NQO1), GSTP1, and MT are responsible for lowering the intracellular concentration of platinum drugs. Increased concentrations of Glutathione (GSH), Metallothioneins (MT) induce resistance against cisplatin. Cytosolic glutathione S-transferase (GST) supergene family includes genes belonging to the subfamilies [alpha] (GSTA), μ (GSTM), [pi] (GSTP) and [theta] (GSTT). GSTs are phase II metabolic isozymes involved in detoxification of many xenobiotic agents like anticancer drugs and carcinogens through conjugation with reduced glutathione (GSH) ¹⁷⁷. This class of enzymes detoxifies carcinogens, therapeutic drugs, and environmental toxins by conjugation with glutathione. Genetic variations in GST genes can affect toxicity and efficacy of drugs. In ovarian tumor cell lines Glutathione S-transferase P1 knockdown showed 2.3- and 4.83-fold change in cisplatin and carboplatin chemosensitivity (IC₅₀) respectively ¹⁷⁸⁻¹⁸⁰. On the other hand, Glutathione S-transferase P1 knockdown in mesothelioma cells exhibited a decrease in GST enzyme activity and GST π protein levels along with an increase in both glutathione levels and sensitivity to cisplatin and oxaliplatin ¹⁸¹. Similarly, GST-pi mRNA was significantly increased in cisplatin resistant gastric cancer and bladder cancer cells ^{182, 183}. SOD is involved in superoxide generation and plays an important role in Platinum drugs detoxification in

cell. Inhibition of the SOD1 activity has been shown to enhanced the cisplatin sensitivity in the ovarian cancer resistant cells ¹⁸⁴.




Figure 1.7 Metabolic pathway of Platinum drugs

1.3.3.2 Genes of importance in drug pharmacodynamics (PD)

1.3.3.2.1 Pt-adduct repair pathway

Upon entering the cell, platinum drug binds to DNA and forms adducts¹⁸⁵. High mobility group (HMG) proteins recognize and bind to DNA intra-strand crosslinks and prevent replicative bypass repair and nucleotide excision repair (NER) leading to initiation of apoptosis^{131, 186}. HMGB1 thus inhibits DNA repair and causes more cisplatin DNA damage in murine model. Knocking out HMGB1 in cell lines showed increased DNA repair (¹⁸⁷). Expression levels of HMGB1 have also been significantly associated with the potency of cisplatin and carboplatin in cancer cells¹⁸⁸. Binding of HMGB1 to DNA-Pt adducts is very specific and the amount of binding is correlated with cytotoxicity of platinum compounds¹⁸⁵. HMGB2 have been shown to enhance the cisplatin sensitivity of cells by inhibiting repair of the DNA lesion caused by cisplatin. Additionally, Polymerase 1 (PARP1) is a protein that also recognizes DNA lesions and plays an important role in the activation of the Base excision repair (BER) pathway¹⁸⁹⁻¹⁹³.

1.3.3.2.2 Mismatch repair pathway

Single stranded DNA damage due to Pt-DNA adducts are recognized and repaired by the DNA mismatch repair pathway (**Figure 1.8**)¹⁹⁴. Genes involved in the mismatch repair pathway include MutS Homolog 2 (MSH2), MutS Homolog 3 (MSH3), MutS Homolog 6(MSH6), MutL Homolog 1(MLH1), MutL Homolog 3 (MLH3), Postmeiotic Segregation Increased 2 (PMS2), Exonuclease1 and PCNA. Lack of detection of Pt-DNA adducts by the mismatch repair genes MSH, MLH and PMS2 (MSH2, MSH3, MSH6, MLH1, MLH3, PMS1 and PMS2) may fail to trigger apoptosis which result in resistance to the platinum drugs^{194, 195}. A number of *in vitro* studies have suggested that MMR gene expression deficiency is a cause of platinum resistance. Furthermore, loss of hMLH1 in ovarian tumor cells was found to be correlated with cisplatin resistance¹⁹⁶. The human colon cancer cell line HCT116+ch2 showed 2.1-fold resistance to cisplatin and 1.3-fold resistance

to carboplatin compared to the subline expressing a wild-type copy of hMLH1. Concurrently, human endometrial cancer cell line HECS9 deficient in hMSH2 showed 1.8-fold resistance to cisplatin and 1.5-fold resistance to carboplatin when compared to a subline with wild-type hMSH2¹⁹⁷. Another study reported that hMLH1 mRNA and protein expression levels were significantly decreased in the cisplatin-resistant head and neck squamous cell carcinoma cell lines compared with a cisplatin-sensitive cell line¹⁹⁸. Furthermore, dMSH2 knockout embryonic stem cell line was 2.1-fold more resistant to cisplatin and 1.7-fold more resistant to carboplatin when compared with wild-type wt-2 cell line while the PMS2^{-/-} mouse fibroblasts were 1.9-fold more resistant to cisplatin and 1.5-fold more resistant to carboplatin when compared to the isogenic PMS2^{+/+} fibroblasts¹⁹⁹. MSH3 deficiency increase sensitivity to both cisplatin and oxaliplatin at clinically relevant doses. MSH3 deficiency contributes to the cytotoxicity of platinum drugs through deficient DSB repair²⁰⁰. Another study reported that, loss of MLH1, MLH2, MSH2, MSH3, MSH6 in isogenic strains of *Saccharomyces cerevisiae* led to increased resistance to cisplatin and carboplatin. Loss of PMS2 gene was found to be associated with an increased sensitivity, ranging from 2-6-fold to platinum compounds cisplatin and oxaliplatin in another study²⁰¹.




Figure 1.8 Mismatch repair (MMR) pathway.

The MMR pathway recognizes and repairs single stranded DNA damage due to Pt-DNA adducts. Major genes involved in this pathway include the MutS and MutL Homolog genes, PMS2, Exonuclease1 and PCNA.

1.3.3.2.3 Nucleotide excision repair pathway

The nucleoside excision repair (NER) pathway genes Xeroderma pigmentosum group (XPA), XPC, XPB/ Excision repair cross-complementing group 1 (ERCC3), XPD (ERCC2), XPG (ERCC5), XPF (ERCC4), ERCC1, DNA polymerases and DNA ligases are primarily involved in the identification and repair of platinum-DNA adducts (1, 2-intrastrand DNA cross-links) ²⁰² (**Figure 1.9**). The human platinum-resistant ovarian carcinoma cell line SKOV3/DDP showed expression levels of ERCC1 were significantly suppressed after stably transfecting with shERCC1-recombinant plasmid, whereas increased sensitivity to cisplatin ($p < 0.01$). In another study siRNA-mediated ERCC1 silencing in the MCF-7, HeLa S3 and HCT116 cells showed both the ERCC1 mRNA and protein expression were significantly inhibited, which led to a decrease in repair activity of cisplatin-induced DNA damage along with cell viability against platinum-based drugs such as cisplatin, carboplatin, and oxaliplatin ²⁰³. On the other hand, ovarian cancer, which is frequently more resistant to cisplatin, shows increased expression of NER proteins including ERCC1 and XPC ²⁰⁴. XPA promotes resistance in tumor cells to platinum based chemotherapeutic drugs. XPA was expressed at higher levels in cisplatin-resistant melanoma cells than in cisplatin-sensitive ones. Furthermore, the knockdown of XPA not only increased cellular apoptosis and also inhibited cisplatin-induced autophagy, which increase cisplatin sensitivity. XPA promoted cell-protective autophagy by polymerase 1 (PARP1) activation in a DNA repair-independent manner and cell became cisplatin resistance ²⁰⁵. Additionally, expression of XPA was higher in NSCLC tissues than that in normal lung tissues. Furthermore, silencing XPA gene increased the apoptosis and cisplatin sensitivity ²⁰⁶.

XPB mRNAs have been observed to be expressed >2-fold higher in clear cell epithelial ovarian tumors ²⁰⁷. Higher mRNA levels of XPB were observed in clinically resistant tumor to platinum-based chemotherapy. In a cohort of 27 patients, mRNA levels of XPB was 5-fold higher in platinum-resistant tumors ($P = 0.001$). Furthermore, these platinum-resistant tumors also show

higher mRNA levels of ERCC1 and XPA ²⁰⁸. ERCC1 is an excision nuclease within the nucleotide excision repair pathway that forms a heterodimer with XPF. ERCC1-XPF complex inhibitor reduced the level of this protein in ovarian cancer cells, inhibited Nucleotide Excision Repair and sensitized cells to cisplatin ²⁰⁹. Non-small cell lung cancer, ovarian and breast cancer cell lines showed increased cisplatin cytotoxicity and efficacy when ERCC1/XPF was down-regulated by RNAi ²¹⁰. The expression of XPF was upregulated in cisplatin resistance KB carcinoma cells (KCP-4) ²¹¹. Osteosarcoma cells (U2OS) by XPF and XPG knockdown exhibited increased sensitivity to platinum drugs compared to the parental cells up to fivefold ²¹². XPF is also required for the repair of DNA lesions through the Nucleotide Excision Repair and Interstrand Crosslink Repair pathways. Xeroderma pigmentosum D (XPD) protein expression levels may also influence response to platinum-based chemotherapy in epithelial ovarian cancer (EOC). In resistance group of serous ovarian cancer, XPD protein expression levels were significantly higher than sensitivity group. Upregulation of ERCC1 and XPD protein expression was associated with resistance process to platinum-based chemotherapy in advanced EOC ²¹³. XPD expression was significantly related to resistance to platinum-containing treatment ($p = 0.043$) ²¹⁴.




Figure 1.9 Nucleotide excision repair (NER) pathway

The NER pathway is involved in the identification and repair of platinum-DNA adducts (1, 2-intrastrand DNA cross-links). Major genes involved in this pathway include the Xeroderma pigmentosum group genes A-G, DNA polymerases and DNA ligases.

1.3.3.2.4 Homologous recombination and non-homologous end joining

Homologous recombination and non-homologous end joining (NHEJ) DNA repair processes are also involved in the repair of double-stranded breaks due to DNA-Pt adducts (Thacker and Zdzienicka. 2004). The genes XPF and ERCC1 gene are involved in homologous recombination while X-ray repair cross complementing group/ X-Ray Repair Complementing Defective Repair in Chinese Hamster Cells 1 (XRCC1), XRCC2, XRCC3, XRCC4, XRCC5, XRCC6 and XRCC7 are involved in non-homologous end joining process ²¹⁵. Protein expression of XRCC1 was significantly increased in cisplatin-resistant gastric cancer cells and caused cisplatin resistance to the cells. Furthermore, Inhibition of XRCC1 expression increased sensitivity of resistant cells to cisplatin ²¹⁶. High XRCC1 expression was found to be associated with platinum resistance ($p < 0.0001$) in ovarian cancer ²¹⁷.

1.3.3.2.5 Translesional bypass replication repair

Finally, when all DNA repair mechanisms fail, translesional bypass replication or SOS repair pathway is activated to overhaul the DNA damages by pt-DNA adducts. Several genes including Polymerase (DNA Directed), Beta or POLB (DNA pol β), POLH (DNA pol η), POLK, POLM, DNA polymerase zeta, REV1, REV3 and REV7 are involved in this pathway ²¹⁸. DNA pol β and DNA pol η Overexpression lead to cisplatin resistance. Polymorphism of these genes also associated with clinical response to the platinum drugs in cancer patients. REV3 is an essential component for of Pol ζ and both play an important role in SOS repair mechanism. Reduced levels of Rev3 exhibited sensitivity to cisplatin and significantly associated with better OS of platinum drug treated mice ²¹⁹. Expression of hRev3 mRNA in the cisplatin resistant head and neck squamous cell carcinoma cell line was approximately 1.4-fold higher than in the sensitive cells. Furthermore, hRev3 knockdown with siRNAs increases sensitivity to cisplatin in this cell line ¹⁹⁸. Reduced expression of the DNA polymerase zeta (catalytic subunit REV3) made fibroblasts more sensitive to cisplatin ^{218, 220}.

1.3.4 Pharmacogenomics of platinum drug pathway genes

Pharmacogenomics is a field of research that combines pharmacology and genomics to study the effect of genes and genetic variations on the response to drugs with an ultimate goal of developing safe and efficacious ‘best-fit’ medications based on an individual’s genetic make-up ²²¹.

We performed a comprehensive review of the research on pharmacogenomics of platinum drug pathway genes conducted through a systematic search of papers reporting association between gene expression and/or genetic polymorphisms in Platinum drug pathway genes and clinical response to platinum-drug based therapy. The genes included in the ensuing review are the most important genes that have been reported to be involved in the transport (influx and efflux), DNA adducts repair, detoxification/xenometabolism (PK) and PD of platinum-based drugs. The criteria for identifying the genes to be selected for the study was based on the information available from the resources providing information on key pathways, e.g. KEGG (Kyoto Encyclopedia of Genes and Genomes), GO (NCBI’s Gene Ontology database), PharmGKB (The Pharmacogenomics Knowledgebase) and PGRN’s (Pharmacogenomics Research Network) drug-specific pathway information. Care was taken to ensure uniformity with respect to gene names and SNP IDs based on scientifically accepted notations and nomenclatures. The following sections will be divided into studies conducted on genetic variations within genes involved in platinum drug pharmacokinetics (transporters, drug metabolizing enzymes) and pharmacodynamics including a summary of the major studies and the conflicting evidences, if any.

1.3.4.1 Genes involved in transport and metabolism of platinating agents (PK genes)

1.3.4.1.1 Drug efflux and influx Transporter

CTR1/SLC31 (Chromosomal location 9q32; length 42965bp; 5 exons) is a predictor of platinum-based chemotherapy response and also plays a role as prognostic factor in NSCLC ^{222, 223}. Ctr1 is predictor of platinum-based chemotherapy response and also plays a role as prognostic factor in

NSCLC ²²³. On the other hand, high SLC31A1/CTR1 expression of was associated with poor survival of (HNSCC) head and neck squamous cell carcinoma patients ²²⁴. In African Americans CTR1 expression is significantly reduced ($P=0.001$) along with tissue platinum concentrations ($P=0.009$) and tumor shrinkage ($P=0.016$) compared to Caucasians ²²⁵. Additionally, CTR1 expression is a predictive/prognostic factor for a good outcome in stage III endometrial cancer patients treated with adjuvant carboplatin based combination chemotherapy. The patients with tumors showing high expression levels of CTR1 associated with longer DFS ($P = 0.058$) and significantly longer OS ($P = 0.003$) ²²⁶. Another recent study showed CTR1 expression was significantly associated with improved chemotherapeutic responses. The median survival time was 15 months in patients with low CTR1 expression, but >66 months in patients with high CTR1 expression ($P<0.001$)²²⁷.

Table 1.3 summarizes genotype-phenotype associations between polymorphisms within platinum drug influx and efflux transporter genes and treatment outcome of platinating agent-containing cancer chemotherapy.

The CTR1 SNP rs10981694 A>C was associated with cisplatin ototoxicity in NSCLC patients. C-Variants are poor tolerance to ototoxicity ($p, 0.05$) ²²⁸. Genetic polymorphisms rs7851395 A>G and rs12686377 T>G were significantly associated with platinum resistance in NSCLC patients treated with platinum-based anticancer drugs. Patients with a GT haplotype showed increased susceptibility to platinum resistance ($P < 0.05$), whereas an AG haplotype contributed to longer survival ($P < 0.05$) in Chinese NSCLC patients $\{\{\}\}$. Further, deletion of the CTR2 gene leads to increased cisplatin and carboplatin accumulation in cells ¹⁵⁰.

Genomic Location for ATP7A gene is Chromosome Xq21.1 and it is 139,740 bases long with 23 exons and 3 splice variants. ATP7B is located at chromosome 13q14.3. This gene is 121,014 bases in length with 34 exons. ATP7A and ATP7B play important roles in platinum drugs efflux.

Resistance cell line to carboplatin, nedaplatin and oxaliplatin showed up-regulated expression of ATP7A, ATP7B transporter ²²⁹. In NSCLC patient's low ATP7A expression was associated with an improved prognosis. The median survival time was 20 months in patients possessing tumors with high ATP7A expression, but >66 months in patients possessing tumors with low ATP7A expression ($P < 0.001$). ATP7B expression was also significantly associated with tumor cell differentiation ²²⁷. ATP7B mRNA and protein expression in colorectal tumors is associated with clinical outcome to oxaliplatin/5FU. Patients with the lowest ATP7B expression significantly associated with longer time to progression ($p = 0.0009$) and increased risk of progression ($p = 0.002$) along with the maximum benefit from platinum based therapy ¹⁵³. ATP7A-positive patients NSCLC patient showed significantly poorer histological grade ($p = 0.039$) and poorer response to platinum-based chemotherapy ($p = 0.001$) compared with ATP7A-negative patients. Therefore, ATP7A expression may be used as an independent prognostic factor for OS ($p = 0.045$) for NSCLC patients treated with platinum based therapy ²³⁰. In a recent gene profiling study (Taqman low-density array-TLDA) on cisplatin sensitive cell lines showed ATP7B gene expression was decreased 1.95-fold ($p = 0.03$) but there was no change in gene expression of ATP7A or SLC31A1 ²³¹.

Only one study so far has investigated the association of ATP7a and ATP7B transporter SNPs with platinum drug response by sequencing of these genes in Japanese cancer patients treated with oxaliplatin/5-fluorouracil/leucovorin and paclitaxel/carboplatin. 38 genetic variations, including 30 novel ones (seven nonsynonymous) in ATP7A and 28 novel and 33 known genetic variations (including 13 nonsynonymous SNPs) in ATP7B were detected. These SNPs possibly influence the efficacy and toxicities of oxaliplatin and carboplatin treatment by changing the drug concentrations within tumor cells ²³².

The ABCC2 gene is located on chromosome 10 (10q24.2) and it is 69,979 bases long. This gene has 34 exons with 3 splice variants; whereas ABCG2 or Breast cancer resistance protein (BCPR)

is located on chromosome 4 (4q22.1) and 141,154 bases long with 19 exons and 4 splice variants. ABCC2 and ABCG2 are involved in carboplatin efflux^{138, 233}. Multiple studies have shown association of these transporters with survival in ovarian cancer patients. Low ABCC2 gene expression in tumor tissues from patients with ovarian cancer was associated with prolonged Progression Free Survival (PFS)²³³. Additionally, localization of ABCC2 in the nuclear membrane was associated with cisplatin resistance and clinical outcome in ovarian carcinoma and overexpression of ABCC2 at mRNA and protein levels was shown to be associated with decreased formation of intra-strand DNA cross-links and resistance to cisplatin in melanoma cells²³⁴. In gastric cancer ABCG2 over-expression was shown associated with longer OS for platinum-containing chemotherapy for²³⁵. In HNSCC (head and neck squamous cell carcinoma) patient's expression of ABCC2, and ABCG2 indicates poor survival of patients treated with platinum based therapy. High ABCB1, ABCC1 expression was associated with both favorable progression-free survival (PFS, $p = 0.0357$, $p = 0.0183$) and OS ($p = 0.0535$, $p = 0.038$)²²⁴.

The non-synonymous polymorphisms rs2273697 (1249G>A/Val417Ile) in ABCC2 and rs2231142 (C421A/Gln141Lys) in ABCG2 were correlated with treatment response and survival in ovarian cancer patients treated with platinum-based chemotherapy. SNP rs2231142 was associated with longer median PFS²³⁶. Another study reported, polymorphism ABCG2 rs2231142 (C421A/Q141K) significantly associated with worse OS for lung cancer patients treated with platinum-based drugs (HR: 1.60; 95% CI 1.04-2.47; $n = 256$)²³⁷. ABCG2 SNPs rs2725264 and rs4148149 were associated with OS ($P=0.041$, $P= 0,014$) in 129 NSCLC patients treated with first-line platinum-based chemotherapy. The median survival duration for the rs2725264 T/T, T/C, and C/C genotypes was 35.75 , 34.25 and 14.89 respectively²³⁸. The ABCC2 polymorphism, rs8187710 (4544G>A), is associated with OS ($p=0.036$) in advanced NSCLC patients treated with platinum-based therapy. Allele A was associated with adverse OS ($p=0.009$) of IV NSCLC patients²³⁹. Polymorphisms in ABCC2 rs717620 (C-24T) and rs3740066 (C3972T) was associated with

treatment response. Homozygotes of -24C was associated with a better treatment response ($P=0.032$). Whereas patients with 3972T had increased risk of severe thrombocytopenia toxicity with female patient group ($P=0.034$), with overall toxicity ($P=0.02$) and hematologic toxicity ($P=0.002$) among platinum-treated non-small cell lung cancer patients ^{240, 241}.

ATP-Binding Cassette, Sub-Family B (MDR/TAP) Member 1 (MDR1/ABCB1) is a 209,691 bases long gene located at chromosome 7q21.12. The MDR1/ABCB1 synonymous coding polymorphism rs1045642 (C3435T/Ile1145Ile) was associated with platinum-based chemotherapy response in lung cancer where the C/C genotype was more sensitive to platinum-based chemotherapy than patients with C/T and T/T. Frequency of this SNP is highest in Asian and Caucasians populations and the lowest in African populations ²⁴²⁻²⁴⁶. rs2032582 (G2677T) is a nonsynonymous SNP located in exon 21 of the ABCB1 gene. ABCB1 2677G>T/A polymorphism were significantly associated with grade 3 or 4 hematological and gastrointestinal toxicities in epithelial ovarian cancer patients and in non-small-cell lung cancer patients treated with platinum and taxen compounds ^{247, 248}. Variant allele (T) of the ABCB1 SNP rs1128503 was associated with the risk of anemia in ovarian cancer patients treated with platinum compound. ($C > T$; $p = 0.023$, OR = 1.71, 95% CI = 1.07-2.71) ²⁴⁹.

ABCC1 gene is located at chromosome 16p13.11 (length: 193,498 bases) and ABCC4 is located at chromosome 13q32.1 (length: 281,618 bases). Higher mRNA expression of ABCC1 significantly correlate with short period of progression free survival in advanced ovarian carcinoma patients treated with platinum based chemotherapy ²⁵⁰. rs1729786, a tagSNP in the gene ABCC4 was found significantly associated (OR=0.68; 95% CI=0.50–0.92; unadjusted $p=0.037$) with chemotherapy-induced peripheral neuropathy (CIPN) among 950 lung cancer patients treated with platinum based therapy ²⁵¹.

Table 1.3 Summary of SNPs in drug transporter genes associated with response to platinating agents.

SNP ID	Gene symbol	Gene Function	Cancer Type	Associated Phenotype	Associated Genotype	PubMed ID (s)
rs10981694	SLC31	Influx transporter	NSCLC	Cisplatin ototoxicity	rs10981694 (A>C) C Variants show poor tolerance to ototoxicity	22516052
rs7851395, rs12686377	SLC31	Influx transporter	NSCLC	Platinum resistance, survival	AG haplotype of rs7851395 A>G and rs12686377 T>G contributed to longer survival. Patients with GT haplotype show increased susceptibility to platinum resistance	22725681
rs2273697	ABCC2	Efflux transporter	Ovarian cancer	Treatment response, survival	1249G>A or Val417Ile	22112610
rs2231142	ABCG2	Efflux transporter	Ovarian cancer	Median PFS	C421A or Gln141Lys	22112610
			Lung cancer	OS	rs2231142 significantly associated with worse OS (HR: 1.60; 95% CI 1.04-2.47)	19107936
rs2725264, rs4148149	ABCG2	Efflux transporter	NSCLC	OS	rs2725264 vs OS (P=0.041); rs4148149 vs OS (P= 0.014)	23689644
rs8187710, rs717620	ABCC2	Efflux transporter	NSCLC	OS , treatment response	rs8187710 vs OS (p=0.036). Allele A was associated with adverse OS (p=0.009); rs717620 (C-24T), 24CC was associated with a better treatment response (P=0.032).	26816351, 19568750
rs3740066	ABCC2	Efflux transporter	NSCLC	Toxicity	rs3740066 (C3972T) 3972T had increased risk of severe thrombocytopenia toxicity (P=0.034), overall toxicity (P=0.02) and hematologic toxicity (P=0.002)	20943283

rs1045642	ABCB1	Efflux transporter	Lung cancer	Chemosensitivity	rs1045642 (C3435T) C/C genotype more sensitive to platinum-based chemotherapy than patients with C/T and T/T.	24687344, 22932088, 22766400, 22704851, 22296372
rs2032582	ABCB1	Efflux transporter	Ovarian cancer	Toxicity	rs2032582 (G2677T), vs grade 3 or 4 hematological and gastrointestinal toxicities	19203783, 20189873
rs1128503	ABCB1	Efflux transporter	Ovarian cancer	Anemia	C > T; p = 0.023, OR = 1.71, 95% CI = 1.07-2.71	25881102
rs1729786	ABCC4	Efflux transporter	Lung cancer	CIPN	rs1729786 vs Chemotherapy-induced peripheral neuropathy (p=0.0377)	25586538

1.3.4.1.2 Xenobiotic metabolism genes

Polymorphisms in GST genes are associated both with cancer susceptibility and anticancer drug resistance²⁵². GSTP1 gene is located on chromosome 11 (11q13.2). This gene has 7 exons along with 7 splice variants. Low GSTP1 expression level was found to be associated with better response in NSCLC patients ($P < 0.003$)²⁵³. Polymorphisms in GST genes are associated both with cancer susceptibility and anticancer drugs resistance²⁵². Genetic variations in GST genes including rs1695 (GSTP1 I105V, shown in **Figure 1.10**) have been shown to be associated both with resistance, toxicity and efficacy of platinum-based drugs^{247, 252, 254, 255}. rs1695 (GSTP1 Ile105Val, A313G) was associated with clinical outcome in NSCLC patients. GSTP Val/Val exhibited a shorter survival time, and had a 1.89-fold greater risk of death than patients with the Ile/Ile genotype. Similar association was also reported by several studies^{241, 256-263}. Additionally, Patients with homozygous mutant GSTP1 GG genotype were at risk for severe platinum-associated polyneuropathy ($P = 0.01$) in NSCLC patients²⁵⁵. rs1695 is also associated with neutropenia in NSCLC patients. Toxicity was less for patients with the 105Val allele ($p = 0.020$)²⁶⁴. Similarly, rs1695 was also found to be associated with PFS ($p = 0.004$) in ovarian cancer patients^{265, 266}. rs1695 was also found to be a significant risk factor for grade 3 or 4 hematological (adjusted OR, 3.08; 95% CI, 1.12-8.43) and gastrointestinal toxicities (adjusted OR, 9.74; 95% CI, 1.59-15.85)²⁶⁷. Additionally, high expression of GSTP1-1 was significantly associated with a poor DFS and OS ($P = 0.047$ and $P = 0.033$, respectively)²³⁴, as well as poor PFS in ovarian cancer patients²⁶⁸. Furthermore, the GSTP1-A-allele was also associated with better prognosis ($P = 0.032$) and GSTP1-AG genotype was associated with necrosis in the tumor's post-chemotherapy histology ($P = 0.001$) testicular germ cell tumor²⁶⁹. GSTP1 expression and rs1695 GA/GG was also significant associated with poor OS ($P=0.004$) in gastric cancer in Chinese population²⁷⁰⁻²⁷². rs1695 was also found to be contributed in colorectal cancer patients treated with oxaliplatin. Patients with Val/Val genotypes significantly associated with better and longer survival of colorectal cancer^{273, 274}. rs1695 was also associated

with toxicity for cisplatin based radiochemotherapy ($P = 0.005$)²⁷⁵ esophageal cancer. Furthermore, rs1695 was also contributed in oxaliplatin-related neuropathy. Patients with homozygous GSTP1 105Ile allele showed more toxicity than patients homozygous or heterozygous for the GSTP1 105Val allele ($P = 0.02$) in gastrointestinal cancer patients treated with oxaliplatin-based chemotherapy²⁷⁶. Another SNP rs1138272 (Ala114Val) was found to be associated with platinum based chemotherapy treated NSCLC patients. Genotype (Ala/Val or Val/Val) had significantly better survival compared with patients with (Ala/Ala; $P = 0.037$). Median survival was 16.1 months compare to 11.4 months for Ala/Ala²⁷⁷.

Glutathione S-Transferase Mu 1 (GSTM1) gene is located on chromosome one (1p13.3). The most important polymorphism in GSTM1 is a partial gene deletion in GSTM1 gene which is called GSTM1 null genotype, resulting in complete absence of GST M1 enzyme activity. 5-year survival and time to progression was better for GSTM1-null genotype patients compared with GSTM1-wt genotype patients ($P = 0.001$ and $P = 0.013$, respectively) in ovarian cancer patients receiving platinum based chemotherapy. GSTM1-wt genotype was also important predictors of risk of death ($P = 0.036$)²⁷⁸. Additionally, thrombocytopenia, anemia and neuropathy were less among patients with the GSTM1-null or GSTM3 intron 6 AGG/AGG genotype in ovarian cancer patients receiving platinum based therapy^{265, 279}. GSTM1 null polymorphism was also associated with treatment response in testicular germ cell tumors, NSCLC patients and lung cancer patients where GSTM1-null were superior responders to platinum drugs than GSTM1-wt ($P = 0.014$)^{258, 261, 269, 280}.

Glutathione S-Transferase Theta 1 (GSTT1) is located on chromosome 22 (22q11.23) and 8,179 bases long. GSTT1 overexpression was found to be influenced decreased cisplatin sensitivity in lung cancer patients treated with platinum compound (¹⁷¹). Non-null GSTT1 genotype was a poor prognostic factor for overall response in ovarian cancer^{247, 281} and toxicity^{247, 266, 279}. Similar association was also observed in gastric cancer patients where response rate was higher in GSTT1 (+) genotype compared with GSTT1 (-) genotype²⁷¹.

Glutathione S-Transferase Alpha 1 (GSTA1) gene is located on chromosome 6 (6p12.2). The GSTA1 promoter region polymorphism rs3957357 (-69 C>T) polymorphism correlated with the OS in ovarian cancer patients receiving platinum drugs chemotherapy. Patients with T/T genotype survived longer than C/C carriers ($P=0.044$)²⁶⁵. GSTA1*B polymorphism comprise with -52G>A (rs3957356), -69C>T (rs3957357) and -567T>G (rs4715332) were also associated with an increased OS in patients with different malignancies treated by platinum compound²⁸².

Glutathione S-Transferase Mu 3 (GSTM3), Glutathione S-Transferase Mu 4 (GSTM4) and Glutathione S-Transferase Mu 5 (GSTA5) genes are located on chromosome 1. The GSTM3 SNP rs1799735 or GSTM3*B induces high GSTM3 expression and results in beneficial effects in cisplatin chemotherapy^{265, 266, 283} in ovarian cancer. Two tag SNPs in GSTA5 (rs4715354) and GSTM4 (rs560018) were associated with survival in lung cancer patients treated with platinum based drugs. Furthermore, rs560018 was significantly associated with IC50 in lymphoblastoid cell lines generated from 100 healthy Caucasian-American subjects ($p=0.019$)¹⁷¹.

SOD1 and SOD2 are also involved in platinum drug xenometabolism. The SOD2 missense/nonsynonymous coding SNP rs4880 (47T/C or Val16Ala) CT + CC combined genotype was shown to be significantly associated with decreased median OS time of 23 months when compared to the TT genotype ($P=0.002$) in gastric cancer patients receiving platinum based combination therapy²⁷⁰. SNP rs4880 was also found to be associated with ototoxicity in cisplatin treated pediatric medulloblastoma patients Texas Children's Cancer Center (1987-2010). The rs4880 T > C substitution results in a Val > Ala amino acid change at position 16 of the SOD. The C-allele of the rs4880 variant was significantly associated with increased SOD2 activity which leads to ototoxicity (odds ratio = 3.06, 95% confidence interval: 1.30-7.20)²⁸⁴.

Myeloperoxidase (MPO) is another component of the detoxification pathway of platinum drugs. The mutant genotypes of the MPO promoter SNP -463G>A was shown associated (A allele) with lower carboplatin-induced hematological toxicity in cancer patients²⁸⁵. Another missense SNP

rs1800566 (C609T/Pro187Ser) in the gene NAD(P)H Quinone Dehydrogenase 1 (NQO1) was significantly associated with higher prevalence of TT genotype ($P < 0.001$). TT genotype carriers had less chance to respond with chemotherapy than compared to CC genotype carriers ($P = 0.003$) in advanced NSCLC receiving platinum-based chemotherapy ²⁸⁶. **Table 1.4** lists the salient variations in xenobiotic metabolism genes associated with pharmacogenomics of platinum drug-based chemotherapy.

Table 1.4 Summary of SNPs in drug metabolism genes associated with Platinum-based chemotherapy.

SNP ID	Gene symbol	Cancer Type	Associated Phenotype	Associated Genotype	PubMed ID (s)
rs1695	GSTP1	NSCLC	Clinical outcome, polyneuropathy	rs1695 (GSTP1 Ile105Val, A313G) Val/Val exhibited a shorter survival time, and had a 1.89-fold greater risk of death than patients with the Ile/Ile genotype. Patients with GG genotype were at risk for severe platinum-associated polyneuropathy (P = 0.01)	023167352, 24958519, 24729086, 22761669, 22009704, 21766492, 19568750, 17409936, 11844594, 22031394, 17409936
		Ovarian cancer	PFS, Toxicity	rs1695 vs PFS (p = 0.004); rs1695 vs grade 3 or 4 hematological (adjusted OR, 3.08; 95% CI, 1.12-8.43); vs gastrointestinal toxicities (adjusted OR, 9.74; 95% CI, and 1.59-15.85)	22188361, 19786980, 19203783
		Testicular germ cell tumor	Prognosis	A allele was associated with better prognosis (P = 0.032) and AG genotype was associated with necrosis in the tumor's post-chemotherapy histology (P = 0.001)	19741569
		Gastric and Colorectal cancers	Tumor response, PFS, and OS	Patients with Val/Val genotypes significantly associated with better and longer survival.	27154175, 25901207, 23020798, 9626464, 22994779, 19922504,
		Esophageal cancer	Toxicity	Rs1695 vs toxicity for cisplatin-based radiochemotherapy (P = 0.005)	21286719
		Gastrointestinal cancer	Neuropathy	Ile/Ile showed more toxicity than Val/Val and Val/Ile genotype (P = 0.02) in patients treated with oxaliplatin-based chemotherapy	16707601
rs1138272	GSTP1	NSCLC	Survival	Genotype (Ala/Val or Val/Val) had significantly better survival compared with patients with (Ala/Ala) (P = 0.037).	16342067

rs3957357	GSTA1	Ovarian cancer	OS	Patients with rs3957357 (C>T) T/T genotype survived longer than C/C carriers (P=0.044) .	19786980
rs4880	SOD2	Gastric cancer, medulloblastoma	OS	rs4880 CT/CC was significantly associated with decreased median OS time compared to the TT genotype (P=0.002).	25901207, 26400460
MPO-463G>A	MPO		Toxicity	Promoter SNP -463G>A was associated with lower carboplatin-induced hematological toxicity	24083736
rs1800566	NQO1	NSCLC	Response	rs1800566 TT genotype carriers showed less response compared to CC genotype carriers (P = 0.003)	24464627




Figure 1.10 Genetic variants in GSTP1 associated with survival/outcome and/or toxicity in patients treated with Platinum drug-based chemotherapy (Green boxes denote coding SNPs).

1.3.4.2 Genes involved in DNA damage and repair (PD) pathway and their impact on platinating agent response

Several genes are involved in the drug pharmacodynamics pathway. HMGB and PARP are involved in the identification of pt-DNA adducts which is followed by repair of these DNA damages (pt-DNA adducts) through NER repair genes, Mismatch repair genes, double strand break repair genes and finally translesional replication repair (SOS) pathway genes when all other repair pathway fails. Polymorphisms in these DNA repair genes may modulate not only DNA repair capacity, but also clinical outcome of treatment with DNA damage-inducing anticancer drugs including platinating agents (**Tables 1.5 and 1.6**).

1.3.4.2.1 Pt-DNA adduct repair

The HMGB1 gene is located at chromosome 13q12.3. It is 160,894 bases long including 8 exons and 8 splice variants. In epithelial ovarian cancer (EOC) patients treated with platinum based therapy HMGB1 expression was an independent predictor for PFS ($P = 0.024$). High HMGB1 expression associated significantly with poorer prognosis (PFS)²⁸⁷. In Chinese lung cancer patients, rs1412125 and rs2249825 of HMGB1 were significantly associated with the platinum-based chemotherapy response²⁸⁸. Poly (ADP-Ribose) Polymerase 1 (PARP1) is a 47,410 bases long gene; located in chromosome 1q42.12 with 23 exons and 10 splice variants. PARP1 is a 113 kDa protein that contains a DNA-binding domain with two zinc fingers and an auto-modification domain. The non-synonymous polymorphism in PARP1 rs1805412 (Val762Ala/T2285C) influenced the prognosis of advanced NSCLC patients treated with platinum-based chemotherapy. PARP1 2285CC genotype was a significantly associated with poor PFS (CC vs. CT/TT: adjusted HR = 1.90, 95 % CI = 1.02-3.52)²⁸⁹.

1.3.4.2.2 Mismatch repair pathway genes

MSH2 and MSH6 are located at chromosome 2p21 and 2p16.3 respectively, whereas MSH3 is located at chromosome 5q14.1. Downregulation of hMSH2 expression has been shown associated with improved survival, while high expression was correlated with poor survival in ovarian cancer patients following cisplatin treatment ¹⁵⁴. Furthermore, MSH2 expression was associated with a shorter time to tumor recurrence, resistance to chemotherapy and death in testicular cancer ²⁹⁰. On the other hand, human MSH6 (hMSH6) defects result in 1.5-4.8 fold increase in cisplatin resistance and 2.5-6 fold increase in replicative bypass of cisplatin adducts ²⁹¹. MSH6 protein levels were also correlated to progression during Platinum-based chemotherapy ($P = .0281$) in patients with malignant pleural mesothelioma (MPM) ²⁹².

Among the mismatch repair gene polymorphisms evaluated for correlation of with response to cisplatin- or carboplatin-based chemotherapy, gIVS12-6T/C (hMSH2) was associated with treatment response ($p=0.0173$) in peripheral lymphocytes from advanced NSCLC patients belonging to the Chinese population ²⁹³. Significant correlation was found between the MSH3 5'UTR SNP rs1105524 and PFS. The G/A and A/A genotypes (median survival time: 14.27 months; 95%CI = 9.80-18.75) showed shorter survival than patients with the G/G genotype (median survival time: 26.37 months; 95%CI = 15.03-37.71) ($P = 0.04$) ²⁹³.

MLH1 is located at chromosome 3p22.2 with 21 exons while MLH3 is located at chromosome 14q24.3 and it has 14 exons. Decreased hMLH1 expression was significantly correlated with improved survival, while high expression was correlated with poor survival in ovarian cancer patients following cisplatin treatment ²⁹⁴. A study by Scartozzi et al observed that among 38 patients with advanced ovarian cancer treated with cisplatin-based chemotherapy, 19 (56%) showing loss of hMLH1 expression with a median survival of 55 months while 15 (44%) showing normal hMLH1 expression had a lower median survival of 12 months ($P=0.014$) ²⁹⁵. MLH1 expression

was also significantly associated with PFS ($P = .0205$) following platinum-based chemotherapy in malignant pleural mesothelioma²⁹². In addition, MLH1 expression was associated with a shorter time to tumor recurrence, resistance to chemotherapy and death in testicular cancer²⁹⁰. Non-functional hMLH1 caused 1.5-4.8-fold increased cisplatin resistance and 2.5-6-fold increased replicative bypass of cisplatin adducts²⁹¹. The non-synonymous coding MLH1 polymorphism rs1799977 (Ile219Leu/A655G) was shown associated with OS of diffuse large B-cell lymphoma (DLBCL). AG/GG genotype displayed an increased death risk (hazard ratio [HR] = 3.23; $P < .001$) compared with patients carrying the AA genotype²⁹⁶.

The PMS1 gene is located on chromosome 2 (2q32.2), while PMS2 is found in clusters on chromosome 7 (7p22.1). Deficiency of PMS2 associated with cisplatin resistance in ovarian cancer patients²⁹⁷.

1.3.4.2.3 Nucleoside excision repair

XPA is located on chromosome 9 (9q22.33) and it has 10 exons along with 4 transcripts. High XPA level predicts a poor prognosis in locally advanced nasopharyngeal carcinoma (NPC) patients treated with platinum-based chemo-radiotherapy²⁹⁸. Malignant ovarian cancer tissues from 13 patients, whose tumors were clinically resistant to therapy showed greater levels of total XPA mRNA and activity ($P = 0.011$) compared to clinically sensitive individuals to platinum therapy ($n = 15$)^{208, 299, 300}. XPA expression in effusion tumor cells from metastatic ovarian carcinoma patients was shown associated with complete response to chemotherapy ($P = 0.03$): XPA expressed in greater than 25% of tumor cells from recurrent patients showed better PFS ($P < 0.001$) and OS ($P = 0.04$)^{301, 302}. XPA rs1800975 (A23G), a promoter polymorphism, is associated with favorable prognosis of NSCLC receiving platinum based chemotherapy. XPA23 A/G+G/G genotypes were associated with significantly longer progression-free survival (PFS) ($P = 0.001$) and OS ($P = 0.001$)^{303, 304}. Another important finding was, association between this rs1800975 and prognosis in 250

inoperable non-small cell lung cancer (NSCLC) patients treated with platinum-based chemotherapy. XPA 23 GA/AA genotypes were associated with poor survival (HR 1.55, $p = 0.011$ overall and HR 1.72, $p = 0.008$) and increased risk of death and progression (HR 1.73, $p = 0.013$ and HR 1.65, $p = 0.016$, respectively) ³⁰⁵. Furthermore, rs1800975 was associated with relative risk (RR) of NSCLC patients receiving platinum based therapy. Homozygous G/G genotype showed a higher RR than patients with G/A or A/A genotype (74.1% vs 50.0%; OR: 0.09; 95% CI: 0.00–0.64; $p = 0.013$) ³⁰⁶. rs1805160, XPA (Arg228Gln) was found significantly associated with recurrence from ovarian cancer in 125 Caucasian patients (HR (AA)= 12.06 (95% CI 1.22–118.76); HR(AG+GG) =9.11 (95% CI 1.12–73.90)) ³⁰⁷. In addition, the XPA missense polymorphism rs104894131 (Cys108Phe) was associated with severe hematological toxicity in 55 NSCLC patients treated with platinum compound in combination with vinorelbine. ³⁰⁸.

XPC is located in chromosome 3 (3p25.1). It is 33,637 bp long and has 18 exons and 7 splice variants. XPC intronic variant rs2607755 was significantly associated with OS and disease free survival of platinum compound treated African Americans head and neck cancer patients (OS HR = 0.62 (0.45, 0.86), DFS HR = 0.51 (0.30, 0.86)) ³⁰⁹. Epithelial ovarian cancer cohort study showed that median PFS of patients carrying the Lys/Lys and Lys/Gln+Gln/Gln genotype of the XPC missense SNP rs2228001 (Lys939Gln/C2704A) polymorphism were 25 and 12 months ($P=0.039$), OS of patients was 31.1 and 27.8 ($P=0.048$). Additionally, patients with the Gln allele had an increased risk of death (HR=1.75; 95% CI=1.06-2.91) compared Lys/Lys genotype ³¹⁰. This polymorphism also showed similar association in NSCLC patients. The heterozygous A/C genotype carriers had a poorer response rate than the wild A/A genotype carriers ($P = 0.023$) ³¹¹. Additionally, rs2228001 was associated with higher toxicity in NSCLC patients treated with platinum based therapy. T allele was associated with higher hematological toxicity (OR: 7.50, 95 % CI: 0.89-63.17, $p = 0.036$) ³⁰⁸. rs2228001 CC genotype was also associated with ototoxicity in cisplatin-treated osteosarcoma patients (P -value= 0.042) ³¹². Furthermore, rs2228001 (Lys939Gln) was significantly associated

with grade 3 or 4 neutropenia ($p = 0.0026$) and grade 3 or 4 hematological toxicity in bladder cancer patients receiving platinum based chemotherapy. Gln/Gln or Lys/Gln + Gln/Gln genotypes of XPC had higher toxicity compared with Lys/Lys (OR: 10, $p = 0.0070$ vs OR: 6.3, $p = 0.0069$; respectively) ³¹³. rs3731108 (AG)/AA genotype versus the GG genotype was associated with prolonged a PFS of 21.3 months versus 13.4 months ($P = .03$) with stage III and IV papillary serous ovarian cancer. Similarly, rs1124303 (GT)/GG genotype versus the TT genotype was associated with a prolonged PFS of 22.8 months versus 14.9 months ($P = .03$) in same patient's cohort ³¹⁴(Fleming ND.et al 2014 PMID: 21751198). Another SNP, XPC rs2228000 (Ala499Val) significantly associated with lung cancer survival ($P = 0.002$). Among 185 stages III-IV NSCLC patients treated with platinum-based chemotherapy ³¹⁵.

XPB/ERCC3 is located at chromosome 2q14.3 with 15 exons and 9 splice variants. XPB/ERCC3 is also associated with platinum drug response in different cancers. For an example, rs3738948 was associated with platinum drug response in NSCLC patients. A/A genotype achieved poor response (68.3%) compare to A/G and G/G genotype (100% and 89.3%) ($p = 0.008$) ³⁰⁶.

XPB/ERCC2 (chromosome 19q13.32; 24 exons and 12 splice variants missense variant rs13181 (2251A>C or Lys751Gln) was associated with nephrotoxicity in platinum drug treated NSCLC patients ³⁰⁸. G allele and T allele carrier patients showed significantly lower nephrotoxicity ($p = 0.017$ and $p = 0.029$) than in other patients ³⁰⁸. The XPD synonymous SNP Asp312Asp rs1799793 (G312A), and XPD Lys751Gln (rs13181) were significantly associated with NSCLC patient's response to platinum drugs therapy ³¹⁶⁻³¹⁹. Patients with Lys/Gln genotype was 0.400 times chemotherapy sensitivity compare with Lys/Lys genotype ($P < 0.05$). Furthermore, Lys/Lys vs Lys/Gln had significant differences in PFS. Whereas, Asp/Asp vs Asp/Asn genotype also showed similar association ³¹⁶⁻³¹⁹. In 93 NSCLC patients significant difference was observed in OS between XPD Asp312Asp and Asp312Asn individuals (20.0 vs 12.4 months, $P=0.04$) ³²⁰. Similar result was observed in another study in NSCLC patient's cohort, where median RFS (relapse-free survival

duration) CC, CT heterozygous and TT mutant allele genotypes were 28.3 months, 46.9 months and 80.1 respectively. Furthermore, median RFS was longer in mutant group ($P = 0.018$)³²¹. Another study also reported that, XPD 312 C/T+T/T and XPD 711 C/T+T/T (rs1052555) associated with poor responses to platinum based chemotherapy in 496 and 375 NSCLC patient's cohort. Furthermore, median PFS and OS of patients of XPD 312 C/T+T/T genotype and XPD 711 C/T+T/T genotype was significantly lower compare to wild-type homozygous patients³²²⁻³²⁴. XPD Lys751Gln (A>C) polymorphism was also significantly associated with decreased PFS in Asians NSCLC patient's population treated with platinum based drug ($P=0.879$)³²⁵. Variant genotypes of XPD Asp(312)Asn, Asp(711)Asp and Lys(751)Gln formed haplotype and were significantly associated with poorer NSCLC survival. The most common haplotype GCA (Asp(312)Asn, Asp(711)Asp and Lys(751)Gln) exhibited significant risk effect on NSCLC survival ($P = 0.001$). ERCC2 Lys751Gln was also associated with survival outcome of metastatic breast cancer patients treated with carboplatin-containing chemotherapy³²⁶. These two SNPs were also associated with Grade 4 neutropenia and nephrotoxicity in advanced NSCLC, ovarian cancer and advanced gastric cancer (AGC) patients treated with platinum based chemotherapy^{263, 265, 313, 327}. In 73 metastatic colorectal cancers patient cohort, rs13181 Lys/Lys genotype responded well ($P = 0.015$). The median survival for those with the Lys/Lys genotype of rs13181 was 17.4 compare to 12.8 and 3.3 months for patients with the Lys/Gln and Gln/Gln respectively ($P = 0.002$). Lys751Lys and the Asp312Asp polymorphism are linked and they form a haplotype ($P < 0.001$)^{328, 329}. Similar result was observed in another study, ERCC2 rs13181T>G, the G allele was associated with reduced response and poor PFS and OS in Caucasians³³⁰. There is evidence that, these two polymorphism affected response and prognosis of bladder cancer patients treated with platinum based chemoradiotherapy³³¹. XPD 751 A/A also showed a significant longer survival in the squamous cell carcinoma ($P=0.034$) and ovarian cancer treated with platinum based chemotherapy^{249, 332}. These two polymorphisms were associated with event-free survival (EFS) in 130 patients with

high-grade osteosarcoma receiving platinum based therapy³³³. Similar result was reported from another study, where ERCC2 rs13181 G allele was significantly associated with shorter event-free survival (P-value=0.021)³¹². Colorectal cancer patients with XPD751 Lys/lys genotype were associated with 3.8-fold increase in failure of chemotherapy compared to Lys/Gln. The median time to progression (MTTP) were 11.3 months vs 2.9 months for patients with Lys/Gln and Gln/Gln genotypes (P < 0.05)³³⁴. Furthermore, XPD Lys751Gln rs13181 (A751C) was significantly associated with Event-free survival and OS in patients receiving oxaliplatin. Patients with C allele had a shorter median event-free survival and OS (6 months vs 11.6 months p = 0.008, 15.6 vs 25.3 months p=0.016)³³⁵⁻³³⁷. In addition, GA+AA genotypes of XPD rs1799793 were associated with improved response to platinum chemotherapy of gastric cancer patients and a significantly decreased risk of mortality³³⁸. AA genotype of ERCC2 rs1799793 was also associated with a better response to chemotherapy, longer OS in gastric cancer patients³³⁹. A synonymous SNP in XPD gene rs238406 Arg(156)Arg (homozygous minor allele) was associated with increased risk of grade 3 or 4 leukopenia (P=0.005) in non-small cell lung cancer patients in a Chinese population receiving platinum based chemotherapy. Further, Arg(156)Arg (rs238406) polymorphism was also significantly associated with grade 3 or 4 hematologic toxicity (; P = 0.009), and severe leukopenia (P= 0.005)^{340, 341}. Furthermore, haplotype "CG" of (Arg(156)Arg-Asp(312)Asn) associated with grade 3 or 4 hematologic toxicity³⁴⁰. ERCC2/XPD rs238406 SNPs were poorer DFS and OS of 572 esophageal squamous cell cancer (ESCC) patient's treated with platinum drugs. Hazard ratio for TT vs. GG+GT for DFS and OS was HR = 1.43, P = 0.020 and HR = 1.52, P = 0.008³⁴². The XPD SNP rs50872 was associated with median survival duration, poor survival and grades 3 and 4 infections in NSCLC patient's population receiving platinum based treatment. Median survival for G/G, A/G, and A/A genotypes was 35.75, 36.07 and 16.75 months, respectively (p<0.001). Additionally, rs238405 and rs238416 was significantly related to OS²³³.

XPF/ERCC4 is a 32,204 bases long gene located at chromosome 16p13.12. It has 14 exons along with 8 splice variants. Low XPF expression in HNSCC patients is associated with better response to induction chemo-radiotherapy, while high XPF expression correlates with a worse response ($p=0.02$) in advanced head and neck cancer receiving platinum therapy ³⁴³. The XPF/ERCC4 (rs12926685) CT+CC genotype versus the TT genotype was associated with a prolonged PFS of 16.7 months versus 12.4 months ($P = .03$) in 139 patients with stage III and IV papillary serous ovarian cancer treated with platinum based therapy ³¹⁴. ERCC6-Q524* and ERCC4-A583T were related with favorable response to first-line platinum-based chemotherapy in advanced-stage high-grade serous EOC. ERCC4-A583T is missense mutation somatic mutation associated with heterozygous loss or complete loss of functional ERCC4, while ERCC6-Q524 is a nonsense mutation ³⁴⁴. Synonymous SNP, rs1799801 in the ERCC4 gene significantly correlated with the relative risk (RR) in advanced NSCLC patients receiving platinum based therapy. Patients with the T/T genotype showed a significantly lower RR than compare to T/C or C/C genotype (25.0% vs 44.7%; OR: 3.30; 95% CI: 1.19–10.00; $p = 0.021$)³⁰⁶.

ERCC5/XPG is located on chromosome 13 (13q33) with 15 exons and 8 splice variants. ERCC5 rs1799977 (His46His), a synonymous polymorphism, were associated with clinical outcomes of 378 advanced non-small cell lung cancer (NSCLC) patients receiving platinum-based chemotherapy regimens. NSCLC patients carrying the ERCC5 46T/T genotype had increased PFS and OS, with 0.52 (0.31-0.96) ^{345, 346}. In 228 advanced NSCLC patient's cohort among three tag-single nucleotide polymorphisms rs2094258, rs751402, and rs2296147 in the ERCC5 promoter region. The rs751402 AA genotype was associated with a better treatment response (AA vs. AG+GG: odds ratio (OR)=2.74, 95% confidence interval (CI) 1.04-7.26, $P=0.036$). Additionally, in the subgroup of patients with squamous cell carcinoma rs751402 AA genotype showed stronger association (AA vs. GG: OR=6.40, 95% CI 1.15-35.50, $P=0.043$; AA vs. AG+GG: OR=6.12, 95% CI 1.15-32.52, $P=0.019$). This result also suggested that ERCC5 rs751402 AA genotype increased

the chemotherapy response in advanced NSCLC, especially in patients with squamous cell carcinoma ³⁴⁷. Furthermore, ERCC5/XPG rs2094258 and rs873601 also showed poorer DFS and OS among patients with resected esophageal squamous cell cancer (ESCC) patients. rs2094258, CT+TT vs. CC: for DFS hazards ratio = 1.68 and P = 0.012; and for OS hazard ratio = 1.99 and P = 0.0001), whereas for rs873601 GA+GG vs. AA, DFS hazard ratio = 1.59 and P = 0.024; OS hazard ratio = 1.91 and P = 0.0005 ³⁴². Another study reported similar associations where CT+TT genotype of rs2296147 had a significantly longer median PFS and OS than CC genotype. Similarly, rs2094258 AG+GG genotype had a longer median progression time and OS time than AA genotype (³⁴⁸. ERCC5 (D1104H) rs17655 was related with more infection (P=0.017) in 388 stage IIIB and IV NSCLC patients treated with platinum-based chemotherapy ³⁴⁹. Additionally, ERCC5 (D1104H) rs17655 was associated with OS (P = 0.004) for lung cancer patients and with PFS for ovarian cancer receiving platinum-based chemotherapy ^{350, 351}. The polymorphism rs751402 located in the 5' UTR of ERCC5 generates a ORF (G= no ORF vs A= ORF) that controls the expression of ERCC5. AA expresses a significantly lower level of the uORF that results in higher levels of ERCC5 protein. cancer patients who carried the A allele were more resistance platinum chemotherapy due to relatively high rates of repair of platinum–DNA adducts within tumors ³⁵². Further, platinum drug treated NSCLC patients carrying the 5' UTR variant rs2296147 TT and the 3' UTR variant rs873601 GG genotypes were significantly associated with favorable outcome for PFS and OS ³⁵³. Furthermore, a significant decreased risk of death was observed among patients carrying the rs2296147 TT genotype compared with the CC genotype ³⁵⁴. In addition, patients with the rs1057768 (C/T) TT genotype had a significantly lower treatment response, short median PFS and OS compared with the rs1057768 CC genotype respectively in NSCLC patient receiving platinum based therapy ³⁵⁴.

The ERCC1 gene is located on chromosome 19 (19q13.32). It is 71,496 bases long gene along with 14 exons and 17 transcripts or splice variants. Increased ERCC1 expression associated with

Platinum drugs resistance in various cancers including ovarian, NSCLC, nasopharyngeal, esophageal, cervical, head and neck squamous carcinoma, liver, osteosarcoma, lung adenocarcinoma, biliary tract adenocarcinoma, mesothelioma, pulmonary adenocarcinoma, bladder cancer, gastric cancer, adrenocortical carcinoma and urothelial carcinoma^{208, 211, 213, 267, 299, 302, 355-373}. EOC patients treated with platinum-based chemotherapy were significantly associated with time to progression (TTP) and OS. OS was significantly longer with low expression group compared with patients showing high expression of ERCC1 protein³⁷⁴. Another study reported that patients with ERCC1-positive tumors had significantly shortened progression-free ($p < 0.00001$) and OS ($p = 0.0006$) compared to patients with ERCC1-negative tumors³⁷⁵⁻³⁷⁸. In advanced cervical cancer patients with negative ERCC1 expression associated with response to Platinum based chemotherapy ($p = 0.021$) as well as disease-free survival for this Patients ($p = 0.046$)³⁷⁹. ERCC1 expression also found to be associated with outcome in gastric cancer patients receiving platinum drugs. Meta-analysis of twenty-one studies involving 1,628 patients of gastric cancer showed high ERCC1 expression was significantly associated with shorter OS and lower response to chemotherapy in advanced GC patients receiving platinum based chemotherapy (HR 1.83; 95 % CI 1.45-2.31; $P < 0.001$; RR 0.49; 95 % CI 0.38-0.62; $P < 0.001$)³⁸⁰⁻³⁸⁸. Many studies have demonstrated a correlation between low levels of ERCC1 and increased survival in NSCLC patients treated with cisplatin³⁸⁹ as well as carboplatin³⁹⁰⁻⁴¹⁶. On the other hand, high levels of expression of ERCC1 mRNA and protein both found to be associated with resistance to cisplatin in NSCLC³⁸⁹. Non-small cell lung cancer (NSCLC) and gastric cancer patients treated with platinum-based chemotherapy showed absence or low expression of ERCC1 was associated with longer disease free survival DFS ($P = 0.037$, $P < 0.0001$, $P = 0.018$) and longer OS ($P = 0.007$)⁴¹⁷. Low ERCC1 expression level was correlated with longer OS (HR (95% CI) of NSCLC patients treated with platinum based therapy. Additionally, NSCLC patients with ERCC1 low expression also had a better PFS ($P < 0.05$) for platinum based chemotherapy. On the other hand, high expression of

ERCC1 was found to be associated with poor prognosis ($P < 0.001$) and decreased survival rates in lung cancer patients treated with platinum based chemotherapy ⁴¹⁷.

A multivariate analysis in platinum-based chemotherapy, low expression of ERCC1 was shown to be an independent predictive factor for response to chemotherapy ($P = 0.018$), time to progression ($P = 0.025$), and OS ($P = 0.038$) ⁴¹⁸. As ERCC1 increased, the probability of response decreased ($P = 0.083$), the risk of progression (ERCC1, HR = 1.59, $P = 0.002$; and death (ERCC1, HR = 1.62, $P = 0.008$) increased ⁴¹⁸. Tumor samples from patients enrolled in the International Adjuvant Lung Cancer Trial concluded that patients with ERCC1-negative tumors appear to benefit from adjuvant cisplatin-based chemotherapy ⁴¹⁹⁻⁴²². Similarly, for patients treated with carboplatin, ERCC1 expression was correlated with OS. The median OS for high expressed ERCC1 patient's vs low expressed group was 9.5 months (95% CI 6.7 to 11.8) vs 15.6 months (95% CI 11.6 to 24.8) ⁴²³. In another study, a NSCLC patient's group treated with carboplatin and paclitaxel combination therapy, the prognosis was significantly better of the ERCC1-negative group compare to ERCC1-positive group ($P = 0.014$) ^{387, 424, 425}. Additionally, meta-analysis evaluated the relationship between ERCC1 expressions and the clinical outcomes including overall response rate (ORR), OS or PFS on 1129 patients with small cell lung cancer (SCLC) receiving platinum-based chemotherapy. Analysis indicated that positive/high ERCC1 expression was associated with unfavorable OS (HR=1.18, 95%CI=1.02-1.37) and PFS (HR=1.46, 95%CI=1.14-1.88) ⁴²⁶⁻⁴²⁸. ERCC1 was also a predictor of shorter survival in Asians and Caucasians for head and neck squamous cell carcinoma (HNSCC) patients treated with platinum-based treatment. High ERCC1 expression was associated with unfavorable OS ($p = 0.009$), PFS ($p = 0.000$) and ORR ($p = 0.044$) ^{429, 430}. Furthermore, ERCC1 was also associated with higher objective response, median PFS, and median OS in patients with advanced bladder cancer treated with platinum-based chemotherapy. Patients with ERCC1 low/negative expression showed significantly prolonged median OS time and the median PFS time compared to ERCC1 high/positive expression (hazard ratio 0.69, 95% CI 0.54-0.89, $P = 0.004$, and

hazard ratio 0.76, 95% CI 0.66-0.89, $P=0.000$, respectively)^{384, 431, 432}. In malignant pleural mesothelioma (MPM) patients receiving platinum based chemotherapy, a significant association was found between ERCC1 protein levels and OS ($P = .032$)^{292, 433}. ERCC1 was also associated with clinical outcomes in patients with pulmonary adenocarcinoma treated with pemetrexed/cisplatin as first-line chemotherapy. Low ERCC1 expression was significantly associated with better relative risk (RR) ($p = 0.015$) and longer PFS ($p = 0.004$). Low ERCC1 expression was also associated with longer OS ($p = 0.003$)⁴³⁴. Nasopharyngeal cancer patients treated with platinum (cisplatin) based chemotherapy also showed that ERCC1-negative tumors had longer disease-free survival ($p = .076$) and OS ($p = .013$) than patients with ERCC1-positive tumors⁴³⁵. Similarly, in biliary tract adenocarcinoma (ABTA) patients treated with platinum-based chemotherapy median PFS was (4.6 vs. 1.9 months, $p = 0.014$) and OS was (9.1 vs. 7.9 months, $p = 0.017$) in ERCC1-negative group than in ERCC1-positive group, respectively. Furthermore, disease control rate (DCR) was also better in patients with ERCC1 negative than in ERCC1 positive patients ($p = 0.048$)⁴¹⁵. ERCC1 was also found to associated with pancreatic cancer patients treated by platinum based chemotherapy Median survival, longer time to progression was significantly higher in group with low ERCC1 levels (Median survival, 11.9 versus 9.9 months, $p \leq 0.05$)⁴³⁶. Similarly, ERCC1 expression also correlated with lack of response ($P=0.006$) and poor disease-specific ($P=0.020$) and OS ($P=0.040$) in gastro-esophageal cancer patients receiving platinum-based chemotherapy⁴³⁷. Furthermore, ERCC1 expression was strongly correlated with poor prognosis and OS in adrenocortical carcinoma (ACC) patients treated with platinum-based chemotherapy. OS was eight months in high ERCC1expressed group whereas, 24 months in low ERCC1 expressed group ($P=0.004$)^{438, 439}.

Figure 1.11 summarizes the structure of ERCC1 gene along with pharmacogenomically relevant polymorphisms that are associated with treatment outcome of platinum based chemotherapy. ERCC1 rs3212986 (C8092A), a missense mutation (Gln506Lys), was found to be associated with

response to platinum based therapy in gastric cancer patients. AA genotype was associated with lower rates of complete remission and partial remission along with higher risk of death in gastric cancer patients ⁴⁴⁰. Similarly, rs3212986, CA and AA genotypes associated with a poorer response to chemotherapy compared to the CC genotype ($P = 0.04$, $P = 0.01$). Patients with the AA genotype associated with longer OS time when compared with the CC genotype (34.91 months vs. 51.19 months, $P = 0.003$) in gastric cancer patients receiving platinum based therapy ^{441, 442}. In gastric cancer patients another SNP rs11615 was also found with clinical outcome. TC+CC genotypes were associated with significantly decreased risk of mortality (hazard ratios = 1.71, 95% CI, 1.06-2.72) ³³⁸. CC genotype of ERCC1 rs11615 was associated with a better response to chemotherapy in gastric cancer patients. CC genotype significantly associated with a longer OS of gastric cancer ^{330, 339}. rs3212986 was also associated with PFS in NSCLC patients. ERCC1 mRNA levels were lower in CC homozygous patients compared with the patients carrying the A allele ($p = .024$)⁴⁴³. NSCLC patients with rs3212986 CA genotype significantly contributed to poorer disease-free survival (DFS) and OS than those with the CC genotype ($p=0.037$ and 0.004) ^{444, 445}. rs3212986 was also found to be associated with a significantly increased risk of grade 3 or 4 gastrointestinal toxicity ($P = 0.03$) in NSCLC patients. Patients with the variant allele of the SNP showed grade 3 or 4 gastrointestinal toxicity (odds ratio, 2.33; 95% confidence interval, 1.07-5.05) ⁴¹⁷. Among 128 advanced NSCLC patients treated with platinum-based chemotherapy, a statistically significant association was observed between the rs3212986 (C8092A) polymorphism and OS ($P = 0.006$, by log-rank test), with median survival times of 22.3 (C/C) and 13.4 (C/A or A/A) months, respectively, suggesting that any copies of the A allele were associated with poor outcome ^{444, 445}. Furthermore, rs3212986 was found to be associated with survival of platinum chemotherapy treated Esophageal squamous cell carcinoma patients ($P=0.01$). Homozygous mutant allele (AA) showing the most significantly reduced survival ($P=0.04$) compared to (CC) ⁴⁴⁶. ERCC1 polymorphism rs3212986 C/A was also associated with PFS and OS of epithelial ovarian cancer (EOC) patients.

The CA or AA genotypes could influence OS (HR = 1.28, 95% CI = 1.05-1.56; and HR = 1.55, 95% CI = 1.17~2.05, respectively) ^{247, 447-449}. On the other hand, rs3212986, C/C wild-type genotype significantly associated with PFS (P = 0.034) in platinum treated malignant mesothelioma (MM) patients ⁴⁵⁰.

In NSCLC, the synonymous coding SNP rs11615 (Asn118Asn), was also found to be associated with response against platinum based chemotherapy. Patients with C/C showed better response compare to C/T + T/T genotype (P = 0.007) ^{243, 257, 315, 451}. Similarly, OS was significantly longer in patient's C/T or T/T allele compare to C/C allele (P=0.014) for rs11615 ^{332, 426, 452}. rs11615 was also found to be associated with ERCC1 expression and platinum drug treated EOC patient's outcome. TT genotype had a significantly poor response (OR = 6.22, 95% CI:1.12-34.42), shorter PFS (HR = 2.19, 95% CI:1.14-4.22, P = 0.009) and OS (HR = 2.22, 95% CI:1.06-4.64, P = 0.021) to platinum-based chemotherapy ^{366, 382, 453, 454}. On the other hand, rs11615 was also associated with the risk of anemia in platinum drug treated ovarian cancer patient. Risk was higher for T allele compare to C allele (T > C; p = 0.031, OR = 1.61, 95% CI = 1.04-2.50) ^{249, 455, 456}. In osteosarcoma patient's rs11615, T/T was strongly associated with a higher event free survival compared to C/C genotype ⁴⁵⁷. Furthermore, ERCC1 C/C significantly associated with better survival and longer survival of colorectal cancer patients treated with platinum based chemotherapy ^{273, 328, 458, 459}. Furthermore, rs11615 polymorphism was associated with therapeutic response in Caucasian patients. Meta-analysis study reported that C allele was found to be associated with better patient response platinum based therapy ⁴⁶⁰. Furthermore, rs11615 TT genotype and T allele and rs3212986 AA and A allele were associated with a poor response and higher risk of death from platinum treated NSCLC patients ^{255, 454, 455, 461-471}. In a study concerning ovarian cancer patients (n=280) median OS was 45, 40 or 30 months for the rs3212986 (C8092A) genotype CC, CA and AA, respectively. Patients with CA+AA versus CC in had increased risk of death (p=0.077) ^{265, 327}. rs3212986, rs11615 were significantly associated with OS in T4 breast cancer patients treated with

chemotherapy containing platinum (p-values = 0.036 and 0.004, respectively). combination of these two genotypes found to be played a role as significant prognostic factor in T4 breast cancer patients receiving platinum-based chemotherapy (p-values = 0.022 and 0.049, respectively) ⁴⁷². rs11615, the T/T genotype was found to be associated with prolonged median cancer-specific survival (p = 0.026) in platinum treated advanced urothelial cancer (UC) ⁴⁷³. Additionally, these two SNPs rs11615 and rs3212986 were associated with higher risk of nephrotoxicity in ovarian cancer and neurotoxicity in advanced gastric cancer patients ⁴⁵⁶. Another 2 SNPs in ERCC1 5'-flanking region, -433T>C and 262G>T was associated with decreased ERCC1 RNA expression. 262G allele is associated with better drug response and longer survival time compared with the 262T allele in SCLC patients ⁴⁷⁴.




Figure 1.11 Genetic variants in ERCC1 associated with survival/outcome and/or toxicity in patients treated with platinum drug-based chemotherapy (Green box denotes coding SNP; Yellow boxes represent non-coding SNPs).

Table 1.5 Summary of SNPs in NER pathway genes associated with Platinum chemotherapy.

SNP ID	Gene symbol	Cancer Type	Associated Phenotype	Associated Genotype	PubMed ID (s)
rs1800975	XPA	NSCLC	PFS, OS	XPA23 (A/G+G/G) were associated with significantly longer PFS (P = 0.001) and overall survival (OS) (P = 0.001)	19430706
			Survival, Progression, Risk	PA23 (A/G+G/G) was associated with poor survival (p = 0.008) and increased risk of death(p = 0.013), progression (p = 0.016) and relative risk (RR). Homozygous G/G genotype showed a higher RR than patients with G/A or A/A genotype (p = 0.013).	22511383, 25069034
rs104894131	XPA	NSCLC	Toxicity	XPA -2166T>C was associated with severe hematological toxicity in 55 NSCLC patients	26193985, 23617284
rs1805160	XPA	Ovarian cancer	Recurrence	HR (AA)= 12.06 (95% CI 1.22–118.76); HR(AG+GG) =9.11 (95% CI 1.12–73.90) in 125 Caucasian patients	17825393
rs2607755	XPC	Head and Neck cancer	OS, DFS	rs2607755 vs OS (HR = 0.62 (0.45, 0.86)); vs DFS (HR = 0.51 (0.30, 0.86)) in African Americans head and neck cancer patients	24487794
rs2228001	XPC	Ovarian cancer	OS	rs2228001 (C/A) vs OS (P=0.048)	23621265
		NSCLC	Response, Toxicity	Heterozygous A/C genotype carriers had a poorer response rate and higher hematological toxicity (p = 0.036.) than the wild A/A genotype	19434073, 26193985
		Osteosarcoma	Toxicity	rs2228001 CC genotype was associated with ototoxicity (P-value= 0.042)	19434073
		Bladder cancer	Toxicity	rs2228001 was significantly associated with grade 3 or 4 neutropenia (p = 0.0026) and grade 3 or 4 hematological toxicity.	21047201
rs1124303	XPC	Bladder cancer	PFS	rs1124303 (GT/GG) genotype (PFS= 22.8 months) vs TT genotype (PFS= 14.9	21751198

				months) was associated with a prolonged PFS(P = .03)	
rs3731108	XPC	Ovarian cancer	PFS	rs3731108 (AG/AA) genotype (PFS=21.3 months) vs GG genotype (PFS=13.4 months) was associated with prolonged progression-free survival (PFS) (P = .03) in patients with stage III and IV papillary serous ovarian cancer.	21751198
rs2228000	XPC	NSCLC	Survival	rs2228000 vs survival (P = 0.002) among 185 stages III-IV NSCLC patients.	21739480
rs3738948	XPB/ERCC3	NSCLC	Response	rs3738948 A/A genotype achieved poor response (68.3%) compare to A/G and G/G genotype (100% and 89.3%) (p = 0.008)	25069034
rs13181	XPD/ERCC2	NSCLC	PFS, RFS, Toxicity	Patients with rs13181 Lys/Gln genotype was 0.400 times chemotherapy sensitivity compare with Lys/Lys genotype (P < 0.05). Lys/Lys vs Lys/Gln had significant differences in PFS and nephrotoxicity (p = 0.017)	26193985, 26124006, 15173214, 16351803, 23977265, 25596702, 24260311
		Colorectal cancer	Response, EFS, OS	Patients cohort with rs13181 Lys/Lys genotype responded well (P = 0.015). Median survival: Lys/Lys =17.4 months; Lys/Gln=12.8 months; Gln/Gln=3.3 (P = 0.002)in 73 metastatic colorectal cancers patient's cohort . rs13181 was significantly associated with EFS and OS. Patients with rs13181 C allele had a shorter median EFS and OS in patients receiving oxaliplatin.	11751380, 26125734, 21278243, 18085999, 18797464, 19432884, 22967467
		Gastric cancer	Respose, PFS, OS	rs13181T>G, the G allele was associated with reduced response and poor PFS and OS in Caucasians gastric and colorectal cancer patients.	21278243
		squamous cell carcinoma,		rs13181 showed a significant longer survival in the squamous cell carcinoma (P=0.034) and ovarian cancer.	21676483, 25881102

		Ovarian cancer			
rs1799793	XPD/ERCC2	Gastric cancer	Response, OS, Risk	rs1799793 was associated with improved response, longer overall survival (OS) and a significantly decreased risk of mortality.	26622786, 25542228
		NSCLC	OS	Significant difference was observed in overall survival (OS) between XPD rs1799793 Asp312Asp and XPD Asp312Asn individuals (P=0.04)	26124006, 15173214, 16351803, 23977265, 24841663, 23534713, 22374424, 24446315
rs13181, rs1799793	XPD/ERCC2	NSCLC, Ovarian cancer and gastric cancer toxicity	Toxicity	rs13181, rs1799793 were associated with Grade 4 neutropenia and nephrotoxicity .	16649224, 19786980, 19332728 , 21047201
		Bladder cancer	Response, Prognosis	rs13181, rs1799793 affected response and prognosis.	16880786
		Osteosarcoma	EFS	rs13181, rs1799793 were associated with event-free survival (EFS) in 130 patients with high-grade osteosarcoma receiving platinum based therapy	21826087, 19434073
rs150461093	XPD/ERCC2	NSCLC	Toxicity	rs150461093 associated with nephrotoxicity.	26193985
rs1052555	XPD/ERCC2	NSCLC	Response, PFS, OS	rs1052555 XPD 711 C/T+T/T was associated with poor responses in 496 and 375 NSCLC patient's cohort. Median PFS and OS of patients of XPD 711 C/T+T/T genotype was significantly lower compared to wild-type homozygous patients.	23534713, 22374424, 24446315
rs238406	XPD/ERCC2	NSCLC	Toxicity	rs238406 vs grade 3 or 4 hematologic toxicity (P = 0.009); vs severe leukopenia (P= 0.005)	22479369, 19458053, 19458053
		Squamous cell cancer	OS, DFS	rs238406 SNPs was associated with poorer DFS and OS.	27246611
rs50871	XPD/ERCC2	Head and Neck cancer	OS, DFS	rs50871 vs OS (HR = 0.80); vs DFS (HR = 0.67).	24487794

rs50872, rs238405 , rs238416	XPD/ERCC2	NSCLC	Survival, Toxicity	Median survival for rs50872 G/G, A/G, and A/A genotypes was 35.75, 36.07 and 16.75 months, respectively (p<0.001); rs50872 was also associated grades 3 and 4 infections; rs238405 and rs238416 were significantly related to OS in NSCLC patients receiving platinum based treatment.	22608006
rs12926685	XPF/ERCC4	Ovarian cancer	PFS	rs12926685 CT/CC genotypes (PFS=16.7) vs TT genotype (PFS= 12.4 months)(P = .03)	21751198
rs3136038, rs3136130	XPF/ERCC4	Head and Neck cancer	Survival	rs3136038 and rs3136130 were associated with survival.	24487794
rs1799801	XPF/ERCC4	NSCLC	Risk	Patients with the rs1799801 T/T genotype showed a significantly lower RR than compare to T/C or C/C genotype (p = 0.021.)	25069034
rs751402	ERCC5/XPG	Squamous cell cancer	Response	rs751402 AA genotype was associated with a better treatment response [AA vs. AG+GG: OR=2.74, 95% CI=1.04-7.26, P=0.036)	23211354
		Pediatric ependymoma	Resistance	rs751402 AA patients were more resistance platinum chemotherapy due to relatively high rates of repair of platinum–DNA adducts within tumors. .	26338418
rs2296147	ERCC5/XPG	NSCLC	PFS, OS	rs2296147 CT+TT genotype had a significantly longer median PFS and OS than CC genotype.	23621222, 24615519, 23621222
rs2094258, rs873601	ERCC5/XPG	Esophageal cancer	OS, DFS	rs2094258 CT+TT vs. CC: for DFS (HR = 1.68 and P = 0.012); for OS (HR = 1.99 and P = 0.0001). rs873601 GA+GG vs. AA, DFS (HR = 1.59 and P = 0.024); OS (HR= 1.91 and P = 0.0005).	27246611
rs2094258	ERCC5/XPG	NSCLC	Progression, Survival	rs2094258 AG+GG genotype had a longer median progression time and OS time than AA genotype.	24353624

rs873601	ERCC5/XPG	NSCLC	Outcome	rs873601 GG genotype was significantly associated with favorable outcome for PFS and OS.	24615519
rs17655	ERCC5/XPG	NSCLC	Infection	rs17655 was related with more infection (P = 0.017).	23118991
		lung cancer patients, Ovarian cancer	PFS, OS	rs17655 was associated with OS (P = 0.004) for lung cancer patients and PFS for ovarian cancer.	17855454, 22158331
		Bladder cancer	Toxicity	rs17655 is associated with grade 3 or 4 neutropenia.	21047201
rs1057768	ERCC5/XPG	NSCLC	Response	rs1057768 TT genotype had a significantly lower treatment response, short median PFS and OS compared with the CC genotype.	23621222
rs11615	ERCC1	Urothelial cancer	Survival	rs11615, the T/T genotype was found to be associated with prolonged median cancer-specific survival (p = 0.026).	23148636
		Pancreatic cancer	Survival	rs11615 was associated with survival.	24403499, 18425336
		Osteosarcoma	EFS	rs11615, T/T was strongly associated with a higher event free survival compared to C/C genotype	23098477
		Colorectal cancer	Survival	rs11615, significantly associated with longer survival	22994779, 22567180, 16144907
rs3212986	ERCC1	Malignant Mesothelioma	PFS	rs3212986, C/C wild-type genotype significantly associated with progression-free survival (PFS).	21765044
		Esophageal cancer	Survival	rs3212986 was found to be associated with survival.	25191856
rs3212986, rs11615	ERCC1	Gastric cancer	Remission, OS	rs3212986 and rs11615, were associated with remission, risk of mortality and OS.	27173253, 26823845, 26722542, 26622786, 25542228, 21278243

		NSCLC	PFS, DFS, OS, Toxicity	rs3212986 and rs11615 were associated with PFS, DFS, OS, death risk, Nephrotoxicity, gastrointestinal toxicity.	25647444, 19361884, 25250341, 24958519, 22532140, 21739480, 22932088, 24859833, 23727606, 22031394, 23226053, 20462983, 20351547, 20143185, 20070981, 19362955, 24370899, 24338713, 24045016, 21676483, 15277258, 15297394, 15746057, 17222938
		Ovarian cancer	PFS, OS, Toxicity	rs3212986 and rs11615 polymorphism in ERCC1 were associated with PFS, OS, anemia and nephrotoxicity in epithelial ovarian cancer (EOC)	24499239, 23632208, 19203783, 18640939, 24444563, 25881102, 22329723, 21496891, 19786980, 19332728, 18024864, 17961161, 23632208
		Brest cancer	OS	rs3212986, rs11615 were significantly associated with OS in T4 breast cancer	25253066
rs2298881	ERCC1	SCLC	Response, Survival	262G allele is associated with better drug response and longer survival time compared with the 262T allele	18451256

1.3.4.2.4 Homologous recombination DNA repair (Double strand break) gene

The XRCC1 gene is located on chromosome 19q13.31 (17 exons, 9 splice variants). XRCC1 expression is a prognostic marker for survival in different cancer patients receiving platinum therapy. Positive XRCC1 was associated with twofold increase risk of death ($p = 0.007$) and progression ($p < 0.0001$) and progression free survival [$p = 0.003$], as well as adverse reactions in ovarian cancer ²¹⁷. XRCC1 expression was associated with disease-specific mortality ($P=0.024$) in bladder cancer patients treated with platinum chemotherapy ³⁵⁹.

The major polymorphisms in this gene associated with platinum based chemotherapy are the missense SNPs rs1799782 (Arg194Trp/C580T) and rs25487 (Arg399Gln/G1196A).

The rs1799782 TT genotype and the CT genotype showed higher efficacy the CC genotype whereas, rs25487 GG genotype was found more sensitive to chemotherapy compared with the AG genotype ^{256, 289, 317, 324, 346, 397, 468, 475-482}.

rs25487 399A/A genotype was significantly associated with longer PFS and OS in 378 lung cancer patients treated with platinum based chemotherapy^{260, 345, 483, 484}. XRCC1 rs25487 is also associated with the clinical outcome of platinum-based chemotherapy in gastric cancer patients. A allele was found to be significantly associated with poor OS ($HR = 1.40$; $95\% CI = 1.04-1.90$) of gastric cancer for platinum-based chemotherapy^{327, 485, 486}. XRCC1 rs25487 was also associated with reduced relative risk (RR) to platinum-based chemotherapy in gastric and colorectal cancer patients. Odds ratio for A/G + A/A vs G/G was 0.73; $95\% CI, 0.55-0.96$ ⁴⁸⁷⁻⁴⁸⁹. Severe neutropenia was associated with rs25487 Arg/Arg genotypes in ovarian cancer patients ²⁶⁵. rs25487 was also shown significantly associated with improved cancer-specific survival in bladder cancer patients receiving platinum drugs ($P=0.009$) ³³¹. Ovarian cancer patients bearing the Trp/Trp genotypes of the missense mutation rs1799782 had longer survival time than Arg/Arg genotype and rs25487, Gln/Gln genotypes had 0.44-fold risk of death than Arg/Arg genotype. The combination of both

194 Trp and rs25487 (Arg399Gln) 399 Gln allele was found to be associated with decrease risk of death in ovarian cancer patients treated with platinum based drugs ^{490, 491}.

Furthermore, rs1799782 was also associated with severe hematological toxicity. The odds ratio (OR) of was significantly lower in patients carries T allele (OR = 0.22, 95 % CI: 0.06-0.82, p = 0.018) in NSCLC patients treated by platinum based drugs ³⁰⁸.

In addition, rs1799782 (Trp vs. Arg, OR=1.342, 95% CI: 1.176) was shown associated with increased risk of cervical cancer and better response to platinum drugs, whereas rs25487 allele A was linked with a poor response ⁴⁹²⁻⁴⁹⁵. rs25487 was also found to be associated with clinical response in platinum drugs treated NSCLC patients. GA or GG genotypes had better response than AA genotype P = 0.028). Additionally, Patients carrying CT or TT genotypes of rs1799782 of XRCC1 showed more sensitive to platinum-based chemotherapy compared to patients with CC genotype (P = 0.002) in NSCLC patients ⁴⁹⁶. A meta-analysis study that included total of 53 studies with 7433 advanced lung cancer patients reported that Trp/Trp and Trp/Arg genotypes of rs1799782 was found to be associated to better response rates to platinum-based chemotherapy compared to Arg/Arg genotype (TrpTrp+TrpArg vs. ArgArg: odds ratio (OR) = 2.02, 95% CI, 1.66-2.45) whereas the Gln/Gln and Gln/Arg genotypes of rs25487 were significantly associated with the poorer response rates compared with the Arg/Arg genotype (GlnGln +GlnArg vs. ArgArg: OR = 0.68, 95% CI, 0.54-0.86)⁴⁹⁷. In another meta-analysis study also concluded that rs1799782 (Arg194Trp) was significantly associated with the efficacy of platinum-based chemotherapy in NSCLC ⁴⁹⁸(Fan X. et al PMID: 26458583). In 122 pancreatic cancer stage-III/IV patients' rs25487 and rs1799782 were associated with clinical response to platinum drug therapy. Arg399Gln + Gln399Gln genotypes had a worse prognosis in pancreatic cancer patients receiving platinum based therapy ⁴⁹⁹.

XRCC2 is a 31,387 bases long gene located at chromosome 7q36.1. It has 3 exons and 2 splice variants. A 5' flanking SNP rs6464268 in XRCC2 was associated with toxicity in esophageal adenocarcinoma patients, who received cisplatin-based preoperative radio-chemotherapy. Patients with AA genotype had grade 3–4 radiation-related toxicity, compared to GG or GA genotypes ($P = 0.005$)²⁷⁵. Further, the XRCC2 41657T allele of the polymorphism C41657T was associated with significantly higher carboplatin and cisplatin sensitivity than lung cancer patients with 41657C/C genotype ($P = 0.046$ and $P = 0.039$, respectively)⁵⁰⁰

XRCC3 variants (Chromosomal location 14q32.33; 10 exons; 18 Splice variants; length=17,897 bases) were found to contribute patient outcome in various types of cancers undergoing platinum chemotherapy. The missense coding SNP rs861539 (C18067T/Thr241Met) was found associated with clinical outcome of 1024 NSCLC patient's cohort receiving platinum based therapy. CC genotype was related to more resistance to platinum-based chemotherapy when compared patients with CT or TT genotypes ($P = 0.009$)⁵⁰¹. Another meta-analysis study also reported that rs861539 was associated with good response to platinum-based chemotherapy in NSCLC patients. Odds ratio (OR) was 1.509 at 95% CI: 1.099-2.072 for ThrMet/MetMet vs. ThrThr.⁵⁰² Similar kind of association was observed in platinum drug treated ovarian cancer patients, where Thr/Thr genotype associated with increased risk of death compare to Met/Met genotype⁴⁹¹. Another study showed that XRCC3 241 Thr/Met or Met/Met allele was associated with longer OS than Thr/Thr allele (19.0 m vs. 12.5 m, $p=0.081$) in NSCLC patients treated with platinum based combination therapy^{322, 503}. The XRCC5 polymorphisms rs1051685 and rs6941 were associated with hematologic toxicity in lung cancer patient treated with platinum based compound⁵⁰¹.

1.3.4.2.5 Translesional Replication DNA repair (SOS) genes

REV1, REV3, and REV7 are involved in translesional DNA repair mechanism in platinum drug metabolism pathway. The G/G genotype of rs240966 (as intronic variant) and A allele of rs456865

(an intronic variant) were associated with higher grade 3 or 4 gastrointestinal toxicity in patients⁵⁰⁴.

NSCLC patients (n=663) treated with platinum based therapy reported that rs240969 and rs3218573 was significantly associated with the treatment response (P=0.0082 and 0.036, respectively). Additionally, patients with G/G genotype of rs240966 or A allele of rs456865 were found to be associated with grade 3 or 4 gastrointestinal toxicity⁵⁰⁴.

POLB is located in chromosome 8 (8p11.21). It has 16 exons with 17 splice variants. High POLB expression had significantly poorer prognosis than patients with low expression ($P < 0.05$) in 97 colorectal cancer patients (CRC) (⁵⁰⁵). The non-synonymous polymorphism in POLB, P242R, was associated with OS ($P = 0.018$) in 456 lung cancer patients treated with platinum-based agents³⁵⁰. is located in chromosome 5 (5q13.3). Finally, rs3213801 within the gene Polymerase (DNA Directed) Kappa (POLK; chromosome 5q13.3) was found to be associated with increased risk of death in NSCLC patients treated with Platinum-Based Chemotherapy. Patients with AA genotype had a shorter survival compared with AG or GG genotype ($P = 0.010$)⁵⁰⁶.

Table 1.6 Summary of SNPs in MMR, NHEJ and SOS pathway genes associated with Platinum drug chemotherapy outcome.

SNP ID	Gene symbol	Gene Function	Cancer Type	Associated Phenotype	Associated Genotype	PubMed ID (s)
rs1412125, rs2249825	HMGB1	pt-DNA adduct recognition	Lung Cancer	Response	rs1412125 and rs2249825 were significantly associated with response in Chinese lung cancer patients.	24684392
rs1805412	PARP1		NSCLC	PFS	rs1805412 (Val762Ala) -762 CC genotype was significantly associated with poor PFS (CC vs. CT/TT: adjusted HR = 1.90, 95 % CI = 1.02-3.52)	23479135
hMSH2 gIVS12-6T/C	MSH2	Mismatch repair pathway	NSCLC	Response	hMSH2 gIVS12-6T/C associated with response to cisplatin- or carboplatin-based chemotherapy in peripheral lymphocytes from advanced NSCLC patients belonging to the Chinese population.	20458443
rs26279, rs1105524	MSH3		NSCLC	PFS	rs26279 (Ala1054Thr) was associated with sensitivity to platinum-based chemotherapy (P = 0.014). rs1105524 G/A and A/A genotypes showed shorter PFS than patients with the G/G genotype (P = 0.04).	25966119
rs1799977	MLH1		B-cell lymphoma	OS, risk	rs1799977 was associated with OS. AG/GG genotype displayed an increased death risk (hazard ratio [HR] = 3.23; P < .001) compared with patients carrying the AA genotype.	21156845
rs5742933	PMS1		NSCLC	OS	rs5742933 was found significantly associated with poorer prognosis on OS in a case-cohort of 568 NSCLC patients	21739480

rs1799782	XRCC1	Double strand break repair pathway	Ovarian cancer	Risk	rs1799782 associated with decrease risk of death	22983827, 22938418
			Mesothelio ma	OS	rs1799782 was associated with overall survival(OS)	22982660
			Bladder cancer	Survival	rs25487 significantly associated with improved cancer-specific survival	24649183
			Ovarian cancer	Toxicity	rs25487polymorphism was associated with neutropenia	19786980
rs25487	XRCC1			OS, risk	rs25487 (Arg399Gln/G1196A) was associated with the OS and risk. A allele was found to be significantly associated with poor OS.	24634242, 24465544, 19332728, 24224851, 17593927, 11712813
			Esophageal cancer	Response , Survival	rs25487 was found to be associated with the response and poor survival	16785472
rs25487 and rs1799782	XRCC1		NSCLC	Response , Toxicity	rs25487 and rs1799782 were associated with clinical response and hematological toxicity.	25784983, 24938464, 24737519, 24729390, 24446315, 23549037, 23479135, 23167352, 22705987, 22551904, 22339849, 20719167, 19362955, 19157633, 15173214, 27248474, 26585370, 26458583, 26193985, 24782167, 21805378, 17504986, 22009704

			Cervical cancer	Risk, Response	rs1799782 was associated increased risk of cervical cancer and better response to platinum drugs. rs25487 A allele was linked with a poor response	23464469, 19563645, 18851872, 16875718
			Pancreatic cancer	Response	rs25487 and rs1799782 was associated with clinical response	22026922
rs6464268	XRCC2		Esophageal cancer	Toxicity	Patients rs6464268 AA genotype had grade 3–4 radiation-related toxicity, compared to GG or GA genotypes	21286719
rs861539	XRCC3		NSCLC	Resistance	rs861539 CC genotype was related to more resistant to platinum-based chemotherapy when compared patients with CT or TT genotypes (P = 0.009)	27248474, 23940523, 22374424, 22152690
			Ovarian cancer	Risk	rs861539 Thr/ Thr genotype associated with increased risk of death compare to Met/Met genotype	22938418
rs861539, rs1799794 and rs861530	XRCC3		Gastric cancer	Survival	rs861539, rs1799794 and rs861530 were associated with survival	21347786
XRCC3 - 316A>G	XRCC3		Mesothelioma	Toxicity	XRCC3 -316A>G was found to associated with treatment-related toxicity.	22982660
rs1051685 and rs6941	XRCC5		Lung cancer	Toxicity	rs1051685 and rs6941 of XRCC5 were associated with hematologic toxicity	26358256
rs240966, rs456865, rs240969, rs3218573	REV1, REV3, and REV7:	Translesional Repair	NSCLC	Response , Toxicity	G/G genotype of rs240966 and rs456865 was associated with higher grade 3 or 4 gastrointestinal toxicity. rs240969 and rs3218573 were significantly associated with the treatment responses.	26278154
rs3792136	REV1		NSCLC	Risk	rs3792136 was associated with increased risk of mortality (P=0.030) .	26611653

rs3087403, rs462779	REV1		Osteosarcoma	EFS, OS	rs3087403 had significantly shorter EFS and OS ($p = 0.004$, $p < 0.001$ respectively). rs462779 was also significantly associated with shorter OS ($p < 0.001$) and shorter EFS ($p = 0.003$) of cisplatin-based chemotherapy.	25748439
POLB P242R	POLB		Lung cancer	OS	POLB P242R was associated with OS ($P = 0.018$) in 456 lung cancer patients treated with platinum-based agents.	17855454
rs3213801	POLK		NSCLC	Survival	Patients with rs3213801 AA genotype had a shorter survival compared with AG or GG genotype ($P = 0.010$).	26611653

1.3.4.3 Additional genetic variants identified by genome-wide analysis studies (GWAS)

Genome--wide gene expression or polymorphism analyses studies (GWAS) are approaches to rapidly scan the complete genome or transcriptome to identify highly significant and actionable genetic variations associated with a target disease or condition ^{507, 508}.

GWAS studies that have so far identified gene expression patterns and SNPs with sensitivity to platinum drugs are listed in **Table 1.7** along with major details ⁵⁰⁹⁻⁵¹³. A GWAS study in 1,244 EOC patients treated with carboplatin and paclitaxel combination therapy identified two SNPs (rs7874043 and rs72700653) in TTC39B which were associated with PFS. On the other hand, genome-wide association scan in 334 NSCLC patient's cohort and 375 patients as validation cohort identified a SNP rs2838566 located at 21q22.3. Minor A allele was found to be significantly associated with increased risk of liver injury. Additionally, rs13014982 at 2q24.3 and rs9909179 at 17p12 were also found to be associated with myelosuppression in NSCLC patients ^{514, 515}. Additionally, another GWAS study explain SNP (rs11138019) association with the expression of gene ABCD2 along with carboplatin and cisplatin response ⁵¹⁶. Associations between NSCLC survival and SNPs were identified by GWAS study in Chinese populations. Three SNPs rs7629386 at 3p22.1, rs969088 at 5p14.1, and rs3850370 at 14q24.3 were associated with worse NSCLC survival while two SNPs rs41997 at 7q31.31 and rs12000445 at 9p21.3 were associated with better NSCLC survival. Furthermore, rs7629386 at 3p22.1 (CTNNB1) and rs3850370 at 14q24.3 (SNW1-ALKBH1-NRXN3) were associated with survival in Caucasian population ⁵¹⁷. Cell-based GWAS approach have been identified single nucleotide polymorphisms associated with platinum susceptibility which were then evaluated in head and neck cancer patients treated with platinum-based therapy. Two SNPs, rs6870861 and rs2551038 were associated significantly with overall response and expression of two important organic cation/anion transporters genes (SLC22A5 and SLCO4C1) which are involved in platinum uptake and clearance ⁵¹⁸. However, these LCLs are derived from

normal healthy individuals and therefore do not represent actual variations of cancer patients. Therefore, in vitro findings using these cell lines cannot be reliably validated using patient data (⁵¹⁸). Similar approach also discovered SNP rs1878022 in the chemokine-like receptor 1 (CMKLR1) which was associated with poor OS in 327 advanced-stage NSCLC patients who received platinum-based chemotherapy⁵¹⁹. GWAS study on SCLC patient cohort identified 3 loci, 20q13.2 (rs4809957G >A), 22q12.2 (rs36600C >T) and 5p15.33 (rs401681C >T) which were significantly associated with the survival time of SCLC patients. Hazard ratio (HR) for patients with the rs4809957 GA or AA genotype was 0.80 (P=0.0187) and 0.73 (P=0.0263) compared with the GG genotype and T allele for rs36600 or rs401681 was 0.78 (P=0.0199) and 1.29 (P=0.0047) respectively compared with the CC genotype ⁵²⁰. Another study in SCLC identified association between rs1820453 T>G within the promoter region of YAP1 on chromosome 11q220 and rs716274 A>G in the region of downstream of DYNC2H1 (11q22.3) with small-cell lung cancer survival. Functional analysis showed that the rs1820453 T>G change creates a transcriptional factor binding site and downregulates YAP1 expression ⁵²¹. GWAS in LCLs for cisplatin cytotoxicity identified rs11169748 and rs2440915 SNPs in DAPK3 and METTL6 which were also associated with OS in lung cancer patients ^{522, 523}. GWAS approach on brain tumor had identified genetic variations in ACYP2 associated with cisplatin-related ototoxicity (rs1872328: P = 3.9×10^{-8} ; hazard ratio = 4.5) in brain tumors in children population ⁵²⁴.

Epstein-Barr Virus (EBV)-transformed human lymphoblastoid cell lines (LCLs) are used as model systems to identify germ-line genetic variations and gene expression associated with anticancer drug response. Using Epstein-Barr Virus (EBV)-transformed human lymphoblastoid cell lines (LCLs) as model system, SNPs in a number of genes have been identified as candidates for platinum-drug and paclitaxel response: The International HapMap Project is an organization that developed a panel of EBV-transformed LCLs (Hapmap panel)

from populations with African, Asian and European ancestry with a goal to create a publicly available catalog of common patterns of human genome variations including SNPs and gene expression ⁵²⁵. Genome-wide association analysis with carboplatin and cisplatin cytotoxicity (IC₅₀) in HapMap LCLs with European (CEU), African (YRI) and Asian ancestries also been reported ⁵²⁶. Association of SNPs and gene expression levels with carboplatin-induced cytotoxicity among 89 HapMap LCLs belonging to African ancestry (YRI) has been studied and results was validated by using 377 ovarian cancer patients receiving at carboplatin-based chemotherapy. SNP rs1649942 could be replicated in an independent LCL set and was also found significantly associated with decreased PFS in ovarian cancer patients ⁵²⁷. Another Genome-wide meta-analysis of SNPs was associated with carboplatin- and cisplatin-induced cytotoxicity among 608 LCLs representing world population panels of diverse ancestry. Results showed a rs7572081 located in intronic region of the gene NBAS (neuroblastoma amplified sequence) and was most significantly associated with carboplatin cytotoxicity ($P=5.1 \times 10^{-7}$). Additionally, rs7210837 was significantly associated with cisplatin IC₅₀ and rs244903 (located in the first exon of RARS) associated with carboplatin IC₅₀. rs10138824 (located in an intron of MPP5-membrane protein, palmitoylated 5), rs8008724 (located in an intron of EIF2S1- eukaryotic translation initiation factor 2, subunit 1 alpha) and rs10431718 (located in an intron of FAM71D-family with sequence similarity 71, member D) associated with IC₅₀ of both drugs and the expression of BCL2, GSTM1 (glutathione S-transferase mu 1) respectively. Furthermore, GSTM1, GSTT1, ERCC2 and ERCC6 were found associated with platinum drug response. ⁵²⁸.

Table 1.7 SNPs found significantly associated with Platinum drug chemotherapy outcome in GWAS studies

SNP ID (Gene symbol)	Cancer Type	Associated Platinum-drug Phenotype	PubMed ID (s)
rs7874043, rs72700653 (TTC39B)	Ovarian cancer	PFS	26840454
rs201023017 (SLC9A9), rs66696671 (TIAL1), rs10674174 (PCDH20), rs1525599 (LRP1B), rs2748151 (CDH4), rs12025262 (ZNF731P), rs150303591 (FRAS1)	Ovarian cancer LCLs	Associated with cytotoxicity in LCLs developed form EOC patients receiving carboplatin combination therapy	27047539
rs1649942	Ovarian cancer	PFS	21705454
rs2838566, rs13014982, rs9909179	NSCLC	rs2838566 A allele was significantly associated with increased risk of liver injury	26100964, 25823687
rs11138019 (ABCD2)		Gene expression of ABCD2; carboplatin and cisplatin response.	24739237
rs7629386 (CTNNB1), rs969088, rs3850370, rs41997, rs12000445	NSCLC	Worse NSCLC survival	22872573
rs6870861 (SLC22A5), rs2551038 (SLCO4C1)	Head and neck cancer	rs6870861 was associated significantly with overall response and expression of SLC22A5, which are involved in platinum uptake and clearance.	21497773
rs1878022 (CMKLR1)	NSCLC	Associated with poor OS in advanced-stage NSCLC patients who received platinum-based chemotherapy	21483023
rs4809957, rs36600, rs401681	SCLC	rs4809957 GA or (HR=0.80; P=0.0187) and AA genotype (HR=0.73; P=0.0263) compared with the GG genotype. Patients with rs36600 T allele (HR=0.78; P=0.0199) compared with the CC genotype. Patients with rs401681 T allele of rs401681 (HR=1.29; P=0.0047) compared to CC genotype.	25415319
rs1820453 (YAP1), rs716274 (DYNC2H1)	SCLC	rs1820453 T>G within the promoter region of YAP1 associated with small-cell lung cancer survival.	21118971
rs11169748 (DAPK3), rs2440915 (METTL6)	Lung cancer	Associated with OS in lung cancer patients	21775533
rs7572081 (NBAS), rs7210837, rs10138824 (MPP5), rs8008724 (EIF2S1), rs10431718 (FAM71D), rs244903 (RARS)	LCLs	Associated with carboplatin cytotoxicity in LCLs	21844884
rs1872328 (ACYP2)	Brain tumor	rs1872328 associated with cisplatin-related ototoxicity (P = 3.9 × 10 ⁽⁻⁸⁾) in brain tumors in children population.	25665007

1.4 Paclitaxel chemotherapy

1.4.1 Introduction

Taxol is a microtubule binding, widely used chemotherapeutic agent ⁴⁶. It was first isolated from the bark of the yew tree (*Taxus brevifolia*) in 1967 by Monroe Wall and Mansukh Wani⁵²⁹. In 1979, Peter Schiff and Susan Horwitz discovered that paclitaxel stimulated microtubule polymerization ⁵³⁰. Major members include paclitaxel and docetaxel. Basic structure of taxanes. **Figure 1.12** shows the basic structure of Taxanes along with differences in paclitaxel and docetaxel based on the moieties attached to the sites marked R1 and R2 in the basic structure ⁵³¹. paclitaxel (Empirical formula: C₄₇H₅₁NO₁₄; Molecular weight: 853.9 g/mol; Average mass: 853.9 Da; Trade names: Abraxane, Taxol, Onxol, Nov-Onxol) was approved for clinical use in 1995 and it is now widely used to treat various cancers including breast and ovarian cancer, non-small-cell lung cancer and Kaposi's sarcoma ⁵³². paclitaxel is a mitotic inhibitor that stabilizes microtubules. The stabilized microtubules interfere with the normal breakdown of microtubules during cell division leading to induction of apoptosis ⁵³²⁻⁵³⁸.

However, the response rate for paclitaxel based therapy is 65%, however tumor recurrence occurs in almost all patients approximately after 15 months ⁴⁶. Drug resistance is the principal cause for 90% of the deaths among EOC patients ⁵³⁹. Furthermore, side effects such as severe hypersensitivity reactions (sHSR), hematological toxicity mainly severe neutropenia and peripheral neuropathy (PNP) are the major challenges for this drugs based chemotherapy ^{531, 540}.

Lack of reliable biomarkers that would enable prediction of response or association with drug related toxicity is another challenge for developing paclitaxel individualized therapy. Identification of genes and polymorphisms associated with paclitaxel treatment outcome is the major goal for developing new therapeutic strategies with maximum drug efficacy along with decreasing drug related toxicity. Here we explore how differential expression and/or activity of genes are involved

in paclitaxel pathway and presence of SNPs can have an impact on treatment outcome and toxicity in cancer patients being treated with paclitaxel.

Understanding the pharmacogenomics of paclitaxel will provide an opportunity to identify likely responders along with poor responders, non-responders and patients with adverse effects which will help to provide better treatment.

1.4.2 Mechanism of action of paclitaxel

Paclitaxel is administered (175mg/m²) in combination with carboplatin (AUC 5–7.5) every 3 weeks for 6 cycles intravenously for EOC and NSCLC patients ^{43, 45, 58, 541-545}. Paclitaxel diffuses through the small openings on the microtubule surface or microtubule lattice fluctuations and binds to β -tubulin on the inside surface of the microtubule resulting in a conformational change in tubulin form (**Figure 1.13**) ⁵⁴⁶. This alteration enhances microtubule assembly, inhibits depolymerisation and stability followed by mitotic arrest resulting from lack of microtubule bundling which eventually leads to apoptotic cell death (**Figure 1.13**) ⁵³²⁻⁵³⁸. In addition, after entering the body, paclitaxel is metabolized by cytochrome enzymes CYP3A and CYP2C ⁵⁴⁷⁻⁵⁵⁰. The major metabolites of paclitaxel are 6- α -hydroxypaclitaxel and p-3'-hydroxypaclitaxel which are formed by CYP2C8 and CYP3A4/5 metabolism, respectively (**Figure 1.12**) ^{551, 552}. Taxol metabolism was studied using membrane fractions from Hep G2 cells contain several human cytochrome P450 enzymes. Only P450 2C8 formed 6 alpha-hydroxytaxol and 3A3 and 3A4 was catalyzed Metabolite B formation. The correlation between hepatic 2C8 protein content and 6 alpha-hydroxytaxol formation was found to be high ($r^2 = 0.82$) (Rahman et al. 1994). The role of individual modifications was then investigated by high-pressure liquid chromatography/mass spectrometry using human liver microsomes and recombinant P450 expressed in Ad293 cells ^{547-549, 549, 550}. Notably, these metabolites do not have appreciable antitumor activity ⁵⁵³.

Paclitaxel is excreted (70–80%) in bile as metabolites or as the parent drug by adenosine triphosphate- (ATP) binding cassette multidrug transporters including P-glycoprotein (P-gp or

ABCC1) and multidrug resistance protein 2 (MRP-2 or ABCC2). Renal clearance occurs only about 5–10% of drugs ⁵⁴⁷⁻⁵⁵⁰.




Figure 1.12 a) Basic structure of taxanes showing differences in paclitaxel and docetaxel (R¹ and R²)⁵³¹ b) Major metabolites of paclitaxel with the enzymes involved⁵⁵³.




Figure 1.13 Mechanism of action of paclitaxel.

Figure shows a) binding to tubulin and b) overall schema of events leading up to cellular sensitivity to paclitaxel ⁵³²

1.4.3 Metabolic (PK/PD) pathway of paclitaxel

Transporters are involved in the regulation of intracellular drug concentration for effective cytotoxic activity of paclitaxel ⁵⁵⁴. paclitaxel is a substrate for several cell membrane transporting enzymes. paclitaxel enters into the cells through OATP1B3 (SLCO1B3) influx transporter. Polymorphism in this gene were found to be associated with paclitaxel pharmacokinetics ^{555, 556}. Additionally, OAT2 (SLC22A7) is highly expressed in human liver and was found to transport paclitaxel in vitro ⁵⁵⁷. On the other hand, ATP-binding cassette (ABC) transporters, which include 48 proteins in humans, is involved in paclitaxel transport. Studies has been shown that inhibition of ATP-binding cassette sub-family B member 1 (ABCB1) or P-gp is associated with increased level of paclitaxel bioavailability ⁵⁵⁸⁻⁵⁶⁰. Furthermore, it has been reported in several studies that ABCB1 was overexpressed in platinum resistant ovarian cancer cells, human lung carcinoma clonal cell line and also in other paclitaxel-resistant cell lines ^{539, 561}. Non-small cell lung carcinoma (NSCLC) patients receiving paclitaxel therapy with no or low expression of MDR-1/P-gp/ABCB1 showed better response compared to patients with increased MDR-1/P-gp/ABCB1 expression ($p<0.05$) ⁵⁶². In ovarian cancer, MDR-1/P-gp overexpression was also found to be significantly associated with poorer paclitaxel response and survival ($p=0.004$) ^{563, 564}. The multi-resistance-associated protein (MRP) family of transport proteins or ABCC (adenosine triphosphate-binding cassette C group) transporters is another class of drug efflux pumps which include ABCC1, ABCG2 and ABCC2 that are involved in paclitaxel transport as well as development of resistance against taxanes ^{168, 550, 565-569}. paclitaxel related neurotoxicity was associated with ABCB1, CYP2C8 and CYP3A5 polymorphisms ⁵⁷⁰⁻⁵⁷³.

CYP2C8 and CYP3A4 are key enzymes involved in the metabolism of paclitaxel. CYP2C8 metabolizes paclitaxel to the primary metabolite 6 α -hydroxypaclitaxel whereas, two minor metabolites 3'-p-hydroxypaclitaxel and 6 α , 3'-pdihydroxypaclitaxel are produced by CYP3A4. Extensive inter-individual variations in CYP genes influence paclitaxel treatment response ^{531, 574-}

⁵⁷⁶. Furthermore, polymorphisms and gene expression changes in pregnane X receptor (PXR), a nuclear receptor involved in CYP enzymes induction and molecular regulation of CYP3A, may also contribute to drug resistance to chemotherapeutic agents in cancer patients ⁵⁷⁷.

Tubulin and its binding capacity with paclitaxel is a major player in paclitaxel drug resistance ⁵⁷⁸. Altered expression levels of microtubule binding proteins (MBPs) are associated with paclitaxel drug response ⁵⁷⁹⁻⁵⁸². For example, microtubule-associated protein (MAP) 7 domain-containing protein 3 plays an integral role in breast cancer growth and metastasis ⁵⁸³⁻⁵⁸⁵. Sub-categories MBPs include canonical MBPs, microtubule-destabilizing proteins, microtubule plus end-tracking proteins, and noncanonical MBPs. Canonical MBPs are structural MAPs and bind to the side of microtubules to stabilize it by promoting the polymerization and formation of microtubule bundles through MAP phosphorylation. MAP2 and tau are the member of this family ⁵⁸⁶. MAP2 promotes paclitaxel sensitivity in breast cancer cell lines ⁵⁸⁷. Knockdown of tau in breast cancer cells increases cancer cell sensitivity to paclitaxel ⁵⁸⁸. Furthermore, studies reported that low concentration of tau is associated with increased paclitaxel binding to microtubules compare to the presence of higher levels of tau. So, low tau expression was found to be associated with the greater response to treatment whereas, high levels of tau was associated with residual tumor and resistance to treatment ⁵⁸⁹⁻⁵⁹². Additionally, overexpression of MAP4 increased microtubule polymerization and increased taxane binding with microtubule which resulted in increased sensitivity to the drugs ^{593, 594}. Microtubule-destabilizing proteins are important regulators of microtubule dynamic instability by restoring free α/β -tubulin heterodimers into a ternary complex, resulting in microtubule de-polymerization which eventually promotes the formation of spindle microtubules ^{595, 596}. Stathmin is the most important member of this family. Elevated expression of stathmin plays a role in cancer susceptibility to paclitaxel treatment ^{597, 598}. Overexpression of stathmin in breast cancer cells significantly decreases the binding of paclitaxel to microtubules and causes paclitaxel resistance, whereas, functional knockdown of stathmin using siRNA results in increased sensitivity

to paclitaxel ^{599-602, 602}. p53 regulates the expression of the MAP4 and statin ^{593, 594, 603-605}. Microtubule plus end-tracking proteins (+TIPs) accumulate at the plus ends of growing microtubules and modulate microtubule dynamics and functions. Members of the end-binding protein (EB) family are master regulators of the dynamic +TIPs network. EB1 regulates paclitaxel-induced mitotic arrest and apoptosis. Expression of EB1 is found to be associated with response to paclitaxel-based chemotherapy in breast cancer patients ^{606, 607}. CLIP-170 is another member of the +TIPs family that is found to be associated with breast cancer cell sensitivity to paclitaxel. Another member, Mitotic centromere-associated kinesin (MCAK), shows association with paclitaxel resistance through microtubule detachment from centrosomes ^{608, 609}.

Non canonical MBPs are involved in the regulation of cell proliferation and survival and are also associated with paclitaxel sensitivity. Parkin is an important member of this sub-category. Higher expression of Parkin is associated with a better response to paclitaxel-containing chemotherapy in breast cancer ⁶¹⁰. Survivin, is another member that locates to the mitotic spindle during mitosis. Upregulation of survivin is associated with paclitaxel resistance in head and neck cancer breast cancer cells ^{611, 612}. Lower expression of Spindle genes MAD2 and BUBR1 in breast cancer cell lines shows resistance to paclitaxel ⁶¹³.

Paclitaxel also has important effects on immune cells including macrophages, dendritic cells (DCs), natural killer (NK) cells, effector T-cells, regulatory T-cells (Tregs), and B-cells which eventually trigger antitumor immune response. paclitaxel stimulates macrophages, which cause cytotoxicity against tumor cells and start anti-tumor immune response ⁶¹⁴. In addition, paclitaxel also downregulates the expression of the anti-apoptotic molecule Bcl-2 in and upregulates expression of the pro-apoptotic member Bax, which leads cells towards apoptosis ⁶¹⁵. Furthermore, multiple translation factors along with gene-related oxidative stress (UGT1A6, MAOA, and CYBA), glycolysis (ADH1A, HK1, and ENO3) and glutathione metabolism are mainly responsible for paclitaxel resistance ⁶¹⁶⁻⁶¹⁸. **Figure 1.14** is a modified version of PharmGKB pathway that

represents genes involved in metabolism and transport of paclitaxel and docetaxel, and the downstream effects of the drugs ⁶¹⁹.




Figure 1.14 Metabolic pathway of taxanes.
The pathway was derived from the PharmGKB database ⁶¹⁹

1.4.4 Pharmacogenomics of paclitaxel (PK/PD) pathway genes

Differential gene expression and polymorphisms in genes involved in paclitaxel metabolism including influx and efflux transporters, pharmacodynamics and therapeutic targets may be associated to the inter-individual variability in response and toxicity. Here we report the relevant genetic variations in the paclitaxel-drug pathway and evaluate their association with chemotherapy. We performed a thorough and comprehensive study of the literature on the genes involved on taxane or paclitaxel-drug based chemotherapy. Our goal was to evaluate all the studies investigating association between gene expression changes and genetic polymorphisms in taxane or paclitaxel pathway genes and clinical response to paclitaxel therapy. One of our criteria for identifying the genes to be selected for the study was based on the pathway information available from the resources providing information on key pathways, e.g. KEGG (Kyoto Encyclopedia of Genes and Genomes), GO (NCBI's Gene Ontology database), PharmGkb (The Pharmacogenomics Knowledgebase) and PGRN's (Pharmacogenomics Research Network) drug-specific pathway information.. Care was taken to ensure uniformity with respect to gene names and SNP IDs based on scientifically accepted notations and nomenclatures.

The following segment presents the salient genes and genetic variations that have been demonstrated to be associated with response to paclitaxel chemotherapy, as well as toxicity.

1.4.4.1 Pharmacogenomics of efflux-influx transporter and drug metabolizing (PK) genes

The organic anion transporting polypeptide (OATP)1B3 or SLCO1B3 is a key influx transporter for paclitaxel⁵⁵⁶. Polymorphisms within this gene have been reported with functional consequences⁵⁵⁵. Two polymorphisms in SLCO1B3, rs4149117 (c.334 T>G, Ser112Ala) and rs7311358 (c.699 G>A, Met233Ile) were associated with grade 3/4 anemia (P=0.002) in NSCLC patients. As shown in **Table 1.8**, rs7311358 (c.699 G>A) was also associated with overexpression of SLCO1B3 resulting significantly decreased uptake of paclitaxel compared with the wild-type⁶²⁰.

The OS time and disease-free survival (DFS) were higher in patients with lower expression of ATP-binding cassette subfamily B member 1 (ABCB1)⁶²¹⁻⁶²³. ABCB1 polymorphism rs2032582 (c.2677 T>G/A) was found to be associated with inferior PFS (hazard ratio=1.49, P=0.017) in NSCLC patients^{620, 624}. Other studies also reported that ABCB1 polymorphisms rs1128503 (1236C>T), rs2032582 (2677G>T) and rs1045642 (3435C>T) have been found to be associated with paclitaxel response in various cancers including ovarian, lung prostate and breast cancer^{247, 570, 622, 625-631, 631-634, 634, 635}. ABCB1 polymorphism rs2032582 (c.2677 T>G/A) was also found associated with grade 3 or 4 hematological and gastrointestinal toxicities in ovarian cancer patients²⁴⁷. Additionally, the 2677 GG genotype showed a significant association with paclitaxel resistance (P = 0.04)⁶³⁶. The ABCB1 polymorphism 3435C>T was associated with increasing risk to develop neuropathy (p=0.09)⁵⁷¹. Another study in metastatic breast cancer patients treated with paclitaxel monotherapy showed that ABCB1 3435 C>T was associated with lower disease (p = 0.025) as well as shorter OS (P = 0.026). A study reported that the synonymous coding SNP rs1128503 (ABCB1; Gly412Gly; C1236T) was found associated with increased risk of anemia (p = 0.023)²⁴⁹. Another SNP p.Ser894Ala/Thr in ABCB1 was found associated with the risk of disease progression (p = 0,010) in ovarian cancer patients receiving paclitaxel/cisplatin chemotherapy⁶³⁷. The SNP rs2229109 (G1199T/A) in the ABCB1 gene was also shown to be associated with paclitaxel resistance in ovarian cancer patients treated with paclitaxel and carboplatin⁶³⁸. The ABCA1 SNP rs4149313 showed significant (p=0.03) association with thrombocytopenia⁶³⁹.

Among SNPs in the ABCC2 gene, rs12762549 (ABCC2, C > G) was associated with decrease risk of anemia (p = 0.004) in ovarian cancer patients. rs2073337 (ABCC2, A > G) and rs1695 (ABCC1, A > G) was associated with colony stimulating factors (CSF) p = 0.039; p = 0.017, respectively), while rs2074087 (ABCC1, G > C) was associated with (p = 0.011) use of erythropoiesis stimulating agents (ESAs)²⁴⁹. rs2725264 in ABCG2 was found to be associated with OS in NSCLC receiving taxane-based chemotherapy (P=0.041)²³⁸, while the non-synonymous variant in ABCG2,

rs2231142 (C421A encoding Q141K, Gln141Lys), was associated with longer median PFS ($p=0.041$) and a reduced risk of disease progression ($p=0.022$) in NSCLC patients ²³⁶.

CYP2C8 polymorphisms were shown associated with reduced metabolizing activity. Two Polymorphisms CYP2C8*3 (R139K; K399R) and CYP2C8*4 (I264M) are common polymorphisms in Caucasian population and associated with paclitaxel clinical response ⁶⁴⁰. Patients with heterozygous for CYP2C8*3 showed lower clearance of paclitaxel ^{570, 641, 642}. Another study reported that two polymorphisms CYP2C8*2 and CYP2C8*3 associated with two-fold higher K_m due to lower activity and lower intrinsic clearance of paclitaxel ⁵⁷⁴. One study found the genetic variants CYP2C8*3, and CYP3A5*3 to be associated with paclitaxel-induced toxicity. CYP2C8-HapC showed association with lower level values of both leukocytes and neutrophils ($p < 0.05$) than patients with the wild-type genotype. Combining the two genotypes CYP3A5*3/*1 and CYP2C8-HapC (rs1113129 G/C) showed significant correlation with both leukopenia and neutropenia ($p = 0.01$) ⁶⁴³. Furthermore, in a study in Chinese epithelial ovarian cancer patients treated with paclitaxel/carboplatin therapy, CYP3A5*3 (rs776746 G/C) was found to be associated with paclitaxel toxicities such as myelo-suppression where CYP3A5*3/*1 genotype showed higher risk of developing leukopenia ($p < .001$) and neutropenia ($p < .001$) ⁶⁴⁴. One study reported that CYP3A4*1B was found to be associated with paclitaxel pharmacokinetics in ovarian cancer patients ⁵⁷⁰. Another CYP3A4 polymorphism rs4986910 (CYP3A4, T > C) was found to be associated with increased risk of thrombocytopenia ($p = 0.025$) ²⁴⁹. Two other CYP3A4 variants, rs2740574 (CYP3A4*1B) and rs35599367 (CYP3A4*22), were found associated with a higher CYP3A4 activity and clinical response of ovarian cancer patients treated with first-line paclitaxel and cisplatin or carboplatin chemotherapy. Patients carrying the CYP3A4*1BG allele showed decreased mean survival rate and lower OS when compared to the AA genotype patients ($p = 0.010$) ⁶⁴⁵. Another study reported that polymorphism CYP3A4*16B was associated with both reduced 3'-p-hydroxylation of paclitaxel and increased levels of 6alpha-hydroxypaclitaxel. CYP3A4*16B

showed lower 3'-p-hydroxypaclitaxel metabolites production ($P = .04$) and a 2.4-fold higher production of 6alpha-hydroxypaclitaxel to paclitaxel ($P < .001$) compared with wild-type (1/1) patients⁶⁴⁶.

Table 1.8 Summary of SNPs in efflux-influx transporter and drug metabolizing (PK) genes associated with Paclitaxel chemotherapy.

SNP ID	Gene symbol	Gene function	Cancer Type	Associated Phenotype	Associated Genotype	PubMed ID (s)
rs2032582	ABCB1	Drug transporter	Ovarian cancer	Risk	p.Ser894Ala/Thr associated with on the risk of disease progression (p = 0.010).	25591549
rs4149313	ABCA1	Drug transporter	Lung cancer, ovarian cancer	Toxicity	rs4149313 was significantly (p=0.03) association with thrombocytopenia.	22759513
rs1045642	ABCB1	Drug transporter	Advanced solid tumours	Toxicity	rs1045642 was associated with increasing risk to develop neuropathy (P=0.09).	16950614
	ABCB1	Drug transporter	Breast cancer	OS	rs1045642 C>T was associated with lower disease (p = 0.025) as well as shorter OS (P = 0.026).	18836089
rs1128503	ABCB1	Drug transporter	Ovarian cancer	Toxicity	rs1128503 was associated with increased risk of anemia (p = 0.023).	25881102
rs2032582	ABCB1	Drug transporter	Ovarian cancer	Toxicity	rs2032582 was associated with grade 3 or 4 hematological and gastrointestinal toxicities.	19203783
rs2032582, rs1128503 and rs1045642	ABCB1	Drug transporter	Ovarian, lung prostate and breast cancer	Response	rs2032582 was found associated with inferior PFS (P=0.017) and GG genotype associated with paclitaxel resistance (P = 0.04) in NSCLC patients. Also associated with paclitaxel response in various cancers including ovarian, lung prostate and breast cancer.	23917080, 19203783, 18765553, 24810093, 21955855, 21883677, 21687948, 20944127, 19401306, 16803472, 16467099, 15901749 and 17062699

rs2229109	ABCB1	Drug transporter	Ovarian cancer	Response	rs2229109 associated with paclitaxel resistance.	17828752
rs12762549	ABCC2	Drug transporter	Ovarian cancer	Toxicity	rs2725264 was associated with decrease risk of anemia (p = 0.004).	25881102
rs2231142	ABCG2	Drug transporter	NSCLC	PSF,Risk	Associated with longer median PFS (p=0.041) and a reduced risk of disease progression (p=0.022).	22112610
rs2725264	ABCG2	Drug transporter	NSCLC	OS	Associated with OS (P=0.041).	23689644
rs4149117, rs7311358	SLCO1B3	Drug transporter	NSCLC	Toxicity	c.334 T>G (Ser112Ala) was associated decreased uptake of paclitaxel and grade 3/4 anemia (P=0.002). c.699 G>A (Met233Ile) was associated with grade 3/4 anemia (P=0.002).	26641474
CYP3A4*16B	CYP3A4	Drug metabolizing enzyme	NSCLC, SCLC, breast cancer, other cancers	PK	Associated with both reduced 3'-p-hydroxylation of paclitaxel and increased levels of 6-alpha-hydroxypaclitaxel	16890579
rs2740574 (CYP3A4*1B)	CYP3A4	Drug metabolizing enzyme	Ovarian cancer	PK, Response	Associated with paclitaxel pharmacokinetics and clinical response of ovarian cancer patients. CYP3A4*1BG allele showed decreased mean survival rate and lower OS when compared to the AA genotype patients (p = 0.010)	23936594, 19143748
rs35599367 (CYP3A4*22)	CYP3A4	Drug metabolizing enzyme	Ovarian cancer	Activity, Response	Associated with a higher CYP3A4 activity and clinical response of ovarian cancer patients	23936594
rs4986910	CYP3A4	Drug metabolizing enzyme	Ovarian cancer	Toxicity	Associated with increased risk of thrombocytopenia (p = 0.025)	25881102

CYP3A5*3	CYP3A5	Drug metabolizing enzyme	Epithelial ovarian cancer	Toxicity	CYP3A5 *3/*1 genotype showed higher risk of developing leukopenia (p < .001) and neutropenia (p < .001). Associated with paclitaxel-induced toxicity such as myelo-suppression.	21702053, 26179145
CYP2C8*3	CYP2C8	Drug metabolizing enzyme	Ovarian cancer	Clearance	Associated with lower clearance of paclitaxel	20368717, 19143748, 17923851, 11668219
rs1113129	CYP2C8	Drug metabolizing enzyme	Ovarian cancer	Toxicity	rs1113129 G/C association with lower level values of both leukocytes and neutrophils (p < 0.05).	21702053

1.4.4.2 Pharmacogenomics of Microtubule binding protein, apoptosis and immune response (PD) genes

Among the genes involved in paclitaxel pharmacodynamics, low/negative class III β -tubulin expression was associated with significantly higher response rate for paclitaxel based chemotherapy in NSCLC patients ($P < 0.00001$). Median survival time was longer for patients with low/negative expression of class III β -tubulin compared with patients with high/positive expression ($P < 0.00001$)⁶⁴⁷. In addition, significantly increased MAP2 Expression in breast cancer patients showed complete response to neoadjuvant paclitaxel⁵⁸⁷. TUBB is one of the primary genes involved in the encoding of tubulin β subunits. Polymorphism of TUBB gene was found to be associated with both poor response and shortened OS to paclitaxel-containing chemotherapy in NSCLC patients⁶⁴⁸. Promoter polymorphisms -101, -112, and -157 in TUBB2A associated with 63-fold variation in β -tubulin IIa gene (TUBB2A) mRNA expression. These polymorphisms were beneficial, and protected patients from paclitaxel-induced peripheral neuropathy (HR, 0.62; 95% confidence interval (CI), 0.42–0.93; $P = 0.021$). Additionally, these polymorphisms were inversely correlated with paclitaxel-induced apoptosis level ($P = 0.001$) in lymphoblastoid cell lines confirming that higher TUBB2A gene expression conferred lower paclitaxel sensitivity⁶⁴⁹.

Most important among the microtubule-destabilizing enzymes, high stathmin (STMN1) expression have been shown to be associated with poor prognosis in ovarian cancer patients treated with paclitaxel and platinum⁶⁰². The promoter polymorphism rs182455 in STMN1 gene is associated with taxane outcome (**Table 1.9**). Patients with TT genotype showed longer PFS and the lower risk of early disease progression ($p = 0.0154$)⁶⁵⁰. Mitotic arrest deficient 1 (MAD1) is a mitotic spindle assembly checkpoint (SAC) protein plays an important role in metaphase arrest. The polymorphism rs1801368 (MAD1 1673 G>A) was associated with the therapeutic response of patients with ovarian cancer with paclitaxel based therapy. The GG patients showed a higher rate of response and cells showed higher percentage of mitotic arrest ($p < 0.05$)⁶⁵¹.

Studies concerning TP53 gene reported that the Pro/Pro genotype of rs1042522 (p.Arg72Pro TP53) in p53 was associated with better progression free survival (PFS), higher response rate, and OS ($p = 0,008$) among ovarian cancer patients receiving paclitaxel/cisplatin chemotherapy^{637, 652-656}. Polymorphisms in apoptotic genes were also found to be associated with paclitaxel response. These include as rs1061624 (TNFRSF1B), rs2279115 (BCL2), rs9904341 (BIRC5), and rs3769818 (CASP8) were significantly associated with OS in NSCLC patients receiving paclitaxel-cisplatin chemotherapy⁶⁵⁷.

A study investigating the role of immune checkpoints genes polymorphisms with paclitaxel-cisplatin chemotherapy in NSCLC patients identified rs2297136T > C and rs4143815C > G in PD-L1 was significantly associated with better chemotherapy response, whereas rs2297136T > C was also associated with better OS^{658, 659}.

Table 1.9 Summary of SNPs in Microtubule binding, apoptosis and immune response (PD) genes associated with Paclitaxel chemotherapy.

SNP ID	Gene symbol	Gene function	Cancer Type	Associated Phenotype	Associated Genotype	PubMed ID (s)
rs1801368	MAD1	Microtubule assembly	Ovarian cancer	Response	rs1801368 was associated with therapeutic response of paclitaxel based therapy. The GG patients showed a higher rate of response and cells showed higher percentage of mitotic arrest (P<0.05).	23407047
rs182455	STMN1	Microtubule assembly	NSCLC	PFS, Risk	Patients with rs182455 TT genotype showed longer PFS and the lower risk of early disease progression (p = 0.0154)	26148901
rs2279115 (BCL2), rs9904341 (BIRC5), rs3769818 (CASP8), rs1061624 (TNFRSF1B)		Apoptosis	NSCLC	OS	Significantly associated with OS.	23973201
rs1042522	TP53	Apoptosis	Ovarian cancer	PFS, OS	rs1042522 was associated with better PFS, higher response rate, and OS (p = 0.008).	25591549, 23574945, 22331725, 16739339, 16364249, 15958617
rs2297136, rs4143815	PD-L1	Immune checkpoint genes	NSCLC	OS, Response	rs2297136T > C was associated with better OS; rs4143815C > G in PD-L1 was significantly associated with better chemotherapy response.	27181838, 27198292

1.4.4.3 Genome-wide association study (GWAS) genes

Genome-wide association study (GWAS) in 1,244 EOC patients treated with the carboplatin and paclitaxel showed two SNPs (rs7874043 and rs72700653) in TTC39B associated with PFS by interacting with the promoters of PSIP1, CCDC171 and an alternative promoter of TTC39B ⁵²³. Another GWAS study in EOC patients treated with paclitaxel/carboplatin standard doses identified a number of SNPs rs4910232 (11p15.3), rs2549714 (16q23), and rs6674079 (1q22) and rs6674079 were significantly associated ($P \leq 1.0 \times 10^{-5}$) with poorer outcomes ⁶³². Furthermore, a GWAS study on advanced NSCLC patients with squamous cell histology showed that tag SNPs rs4151510, rs4151465, rs9568036 in RB1 gene was associated with clinical response of first-line platinum-taxane. The patients with G/G genotype of rs4151510 had longer OS ($p=0.018$) ⁶⁶⁰. An extensive GWAS study performed in 276 population panel LCLs and validated in 76 small and 798 non-small cell lung cancer (SCLC, NSCLC) patients treated with paclitaxel found a SNP rs1106697 on chromosome 7 to be associated with paclitaxel response in both LCLs and lung cancer patients (NSCLC and SCLC) ($p = 0.016$ and $p = 0.007$ respectively). The three SNPs rs1778335, rs2662411 and rs7519667 were associated with SCLC OS ($p = < 0.05$) ⁶⁶¹. Among other GWAS studies, one analysis showed association of SNPs in AIPL1 and BCR with sensory neuropathy in cancer patients ⁶⁶²; rs1656402 in the EIF4E2 gene, rs1209950 in the ETS2 gene and rs9981861 in the DSCAM gene was found associated with poor OS in NSCLC patients ⁶⁶³. A study performing analysis of single nucleotide polymorphisms (SNPs) located in the miRNA target sites and their role in paclitaxel-cisplatin chemotherapy in advanced NSCLC showed ETS2 rs461155A>G was significantly associated with decreased ETS2 mRNA expression⁶⁶⁴. **Table 1.10** summarizes the SNPs that were found significantly associated with paclitaxel chemotherapy in GWAS studies conducted so far.

Table 1.10 SNPs found significantly associated with paclitaxel drug chemotherapy outcome in GWAS studies.

SNP ID (Gene symbol)	Cancer/Cell Type	Associated Phenotype	Associated Genotype	PubMed ID (s)
rs7874043, rs72700653 (TTC39B)	EOC	PFS	rs7874043 and rs72700653 were associated with PFS	26840454
rs4910232, rs2549714, rs6674079, rs6674079	EOC	Response	significantly associated ($P \leq 1.0 \times 10^{-5}$) with poorer outcomes	26152742
rs4151510, rs4151465, rs9568036 (RB1)	NSCLC	Response	associated with clinical response	25684524
rs1106697, rs1778335, rs2662411 and rs7519667	NSCLC, SCLC	Response, OS	rs1106697 was associated with paclitaxel response in both LCLs and lung cancer patients (NSCLC and SCLC). rs1778335, rs2662411 and rs7519667 were associated with SCLC OS ($p = < 0.05$)	23006423
rs1656402 (EIF4E2), rs1209950 (ETS2), rs9981861 (DSCAM)	NSCLC	OS	Associated with poor OS	21079520
rs461155 (ETS2)	LCLs	mRNA expression	rs461155A>G was significantly associated with decreased ETS2 mRNA expression	26893365

1.5 Research Study Objectives

1.5.1 Gaps in literature

Understanding the pharmacogenomics of response to carboplatin and paclitaxel chemotherapy is a pre-requisite to identify reliable genetic signatures of drug response and toxicity. Several studies, including GWAS, have investigated the association of differential expression and/or activity of genes involved in the PK/PD pathway due to SNPs with platinum drug-based treatment outcome and/or toxicity in various cancers and have identified several key pathway genes polymorphisms^{227, 229, 665, 666}. However, a comprehensive pharmacogenomics-based evaluation of these genes to identify genomic markers predictive of efficacy/sensitivity is still lacking. However, most SNPs identified in such studies have not been validated using independent investigation.

One of the strategies to identify biomarkers associated with cellular sensitivity to carboplatin and/or paclitaxel is use of Epstein-Barr Virus (EBV)-transformed human lymphoblastoid cell lines (LCLs) as model systems. SNPs associated with sensitivity have been identified by genome wide analysis using LCLs that are part of International HapMap project^{189, 191, 192, 667}. However, one of the biggest disadvantages of using these LCLs is that they were derived from normal healthy individuals and therefore do not represent variations in cancer patients. Secondly, these cell lines might not reflect gene expression profiles of the target patient subject. Furthermore, large-scale association studies conducted so far have not provided reliable and clinically applicable biomarkers of carboplatin/paclitaxel response in ovarian/lung cancer due to low sample sizes or lack of reproducibility^{138, 186, 288, 668, 669}. Notably, no study so far has focused on the assessment of carboplatin + paclitaxel combination treatment response using *in vitro* model systems. Therefore, a systematic and comprehensive genotype-phenotype association study is required to understand the association between genetic variants and carboplatin + paclitaxel drug response in cancer treatment.

1.5.2 Objectives and Specific Aims

We hypothesize that we will identify a robust set of genetic variants and gene expression signatures significantly associated with drug response and treatment outcomes within *in vitro* cancer cell lines model systems as well as cancer patients undergoing carboplatin and paclitaxel-based chemotherapy. Thus, the objectives of this study were to: 1) Use LCLs derived from ovarian cancer patients to identify predictive markers of chemo-sensitivity; 2) Comprehensively and systematically evaluate association of gene expression and SNPs with treatment outcome in *in vitro* cell line-based models and clinical trials of ovarian and lung cancer patients.

Our objectives were achieved through the following specific aims:

Specific Aim 1. To create an *in vitro* cytotoxicity profile of EBV-transformed LCLs from epithelial ovarian cancer patients in response to carboplatin and paclitaxel as single agents and in combination.

Hypothesis: LCLs from ovarian cancer patients demonstrate inter-patient variation in *in vitro* cellular sensitivity to carboplatin and paclitaxel.

Specific Aim 2. To evaluate gene expression levels and SNPs in key PK and PD pathway genes in epithelial ovarian cancer patients and correlate these with cytotoxicity data obtained from Aim1.

Hypothesis: Variation in gene expression levels and the presence of SNPs in PK/PD pathway genes influence treatment outcomes in ovarian cancer patients.

Specific Aim 3. To determine correlation of cytotoxicity, gene expression and SNPs identified using Genome-wide association study with clinical response (progression free and OS) and toxicity (myelosuppression, gastrointestinal toxicity) in the ovarian cancer patients from which the LCLs are derived.

Hypothesis: *In vitro* cytotoxicity profiles, gene expression and SNPs are correlated with clinical outcome and toxicity in cancer patients treated with carboplatin/paclitaxel-containing chemotherapy.

Specific Aim 4: To identify genetic variations in platinum drugs and taxane pathway genes as predictors of outcome and toxicity in advanced non-small-cell lung cancer (NSCLC)

Hypothesis: SNPs within the genes involved in the metabolic pathways of taxanes and platinating agents are associated with clinical outcome and toxicity in NSCLC patients treated with carboplatin/paclitaxel-containing chemotherapy.

Long term goal: Successful completion of the project will generate a robust panel of genetic biomarkers of carboplatin and paclitaxel single-agent and combination treatment response that can serve as a first step towards the development of a precision medicine approach to develop pharmacogenomics-guided chemotherapy for maximum efficacy and minimum toxicity.

CHAPTER 2 PHARMACOGENOMIC MARKERS INFLUENCE THE
VARIATION OF IN VITRO CHEMOSENSITIVITY TO CARBOPLATIN AND
PACLITAXEL AS SINGLE AGENTS AND AS COMBINATION: A PATHWAY
BASED APPROACH

2.1 Introduction

Ovarian cancer is the most lethal gynecologic malignancy in the United States (NCI SEER cancer statistics) ¹. An estimated 220,000 new ovarian cancer (OC) cases are diagnosed every year worldwide while around 140,000 women die from this disease ². Epithelial ovarian cancer (EOC) is the one of the most common gynecological malignancies and the fifth leading cause of cancer death among women in the United States ³. EOC occurs primarily in middle aged or older women, rarely before puberty ⁴.

The standard treatment for 75% of EOC patients with advanced disease is initial debulking surgery followed by carboplatin-paclitaxel combination chemotherapy⁵. carboplatin [cis-diammine (1,1-cyclobutanedicarboxylato) platinum(II)] or Paraplatin was approved by FDA in 1989 for the treatment of advanced ovarian cancer ⁶. carboplatin forms DNA inter-strand, intra-strand, and DNA-protein crosslinks which is attributed to the formation of DNA helix-distorting adducts that result in strand breakage followed by activation of DNA repair mechanisms that interfere with cell division, which ultimately results in apoptosis ⁷. Some of the major genes involved in platinum pathway are influx and efflux transporters *SLC31A1*, *ABCC2*, *ATP7*; DNA repair genes *MSH6*, *MLH1*, *XRCC1*, *ERCC1*, *ERCC2*, *XPA*; and xenobiotic metabolism genes *MPO*, *SOD1*, *GSTM1*, *NQO1*, *GSTP1* ⁸⁻¹⁰. paclitaxel is another chemotherapeutic agent, used generally in combination with platinum drugs for various cancers including lung and ovarian cancers ¹¹. Taxanes like paclitaxel block cell division by binding to α -tubulin that stabilizes the microtubules resulting in cell death ^{11,12}, a process regulated by microtubule- associated proteins (*MAPs*) *MAPT*, *MAP2* and *MAP4* ¹³. paclitaxel is transported into the cell through influx transporter *OATP1B3* (*SLCO1B3*) and metabolized by *CYP3A4* and *CYP2C8* ^{14, 15} while the drug is effluxed out of the cell by ATP binding cassette (*ABC*) multidrug transporters *ABCB1*, *ABCG2*, *ABCC1* and *ABCC2*.

However, despite currently available therapeutic strategies, 5-year overall survival (OS) in EOC remains around 44.2% along with wide inter-patient variation in response ¹⁶. Response rate is 10–

15% and overall survival is ~12 months in resistant patients ¹⁷. 90% of the patients relapse after 18 months of treatment ¹⁸.

Though patients commonly develop resistance to these chemotherapeutic agents, the underlying mechanisms still remain unclear. Alterations in expression of key candidate genes involved in drug metabolism pathways and/or the presence of candidate SNPs may have a profound impact on treatment outcome/resistance and toxicity in ovarian cancer patients. However, there are not many reliable prognostic biomarkers for either carboplatin or platinum to help improve clinical response.

Epstein-Barr virus (EBV)-transformed Lymphoblastoid cell lines (LCLs) are immortalized cell lines generated from human normal lymphocytes through the infection of Epstein-Barr virus (EBV) ¹⁹. EBV-transformed LCLs comprising of multi-ethnic panels representing multiple world subpopulations, including the International HapMap project ²⁰⁻²² and 1000 genomes project ²³, have been extensively used to identify genetic biomarkers for various chemotherapeutic agents using both Genome-wide association and candidate gene approaches ^{20, 24-26}. However, since these LCLs are derived from 'normal' healthy individuals, they do not truly represent the genetic architecture of EOC patients.

In the current study, we created EBV-transformed LCLs from cells derived from EOC patients, performed *in vitro* chemo-sensitivity assays to derive cytotoxicity profiles for carboplatin, paclitaxel and combination treatments, and conducted genetic association studies between *in vitro* drug response phenotypes, gene expression levels and genetic variations of drug pharmacokinetic (PK) and pharmacodynamics (PD) pathway genes to identify reliable genomic signatures of treatment outcomes.

2.2 Materials and Methods

2.2.1 Creation of EBV-transformed Ovarian cancer LCLs

Prior to initiation of chemotherapy, blood was collected following informed consent from ovarian cancer patients, diagnosed at the Mayo Clinic between 2000 and 2003. Epstein Barr Virus (EBV)-transformed LCLs were generated from patients and subjected to *in vitro* drug testing. All patients provided informed consent, including for passive and active follow-up, using protocols approved by the appropriate Institutional Review Board at the Mayo Clinic in Rochester, MN.

2.2.2 Cell Culture

All the cell lines were cultured in RPMI1640 media (Gibco) supplemented with 2 mM L-glutamine (Gibco) and 15% fetal bovine serum (HyClone) and were maintained in a humidified incubator at 37°C with 5% CO₂. The cells were passaged every 2 to 3 days in order to maintain them in logarithmic growth phase. To confirm healthy growth, cell viability was measured using Countess Automated Cell Counter (Life Technologies) and cell line with >80% viability was used for further experiments.

2.2.3 Drugs

Carboplatin and paclitaxel were procured from Sigma Aldrich (St. Louis, MO). carboplatin was freshly prepared in PBS while paclitaxel was prepared in DMSO (Dimethyl sulfoxide) and stored in -20°C in aliquots for 8 weeks.

2.2.4 In vitro cytotoxicity assays

MTT (5-diphenyltetrazolium bromide) reagent (Sigma Aldrich, St. Louis, MO) was used to perform *in vitro* cytotoxicity assays on the cell lines treated with single-agent or combination regimens. Briefly, LCLs were plated in 96-well plate at seeding density of 2.5×10^5 cells/ml and incubated at 37°C overnight. Following 24 hour incubation, LCLs were treated with increasing concentrations of carboplatin and/or paclitaxel (in duplicate). Concentrations were 0, 5, 10, 20, 40, 80 and 128 uM for carboplatin and 0, 4.5, 7.5, 10, 20, 40 and 80 nM for paclitaxel treated as single agents. In combination treatment, individual drug doses were half (0.5) of the doses for each drug

used as single agent in increasing concentrations. Cells were incubated at 37C for 48hrs following treatment. Cell viability was determined 48hrs post-treatment by incubating the cells with MTT reagent followed by measurement of absorbance at 570 nm the using Synergy plate reader (BioTek, USA). The percent cell survival at each concentration and IC₅₀ values and area under survival curve (AUC) were calculated as described in the statistical analysis sub-section.

2.2.5 Caspase-Glo 3/7 activity Assay

Apoptotic activity within the LCLs was determined 48 hours following treatment with carboplatin, paclitaxel, or combination, the using Caspase-Glo® 3/7 Assay (Promega, USA). Luminescence Intensity were determined as measures of caspase activity using Synergy plate reader (BioTek, USA). The caspase activity at each concentration was normalized to control.

2.2.6 Cell cycle analysis

Prior to plating for cytotoxicity and apoptosis experiments, an aliquot from each fresh cell line was collected to measure %S phase. Briefly, cells were stained with Propidium Iodide (PI) followed by assessment of cell cycle phases using BD FACSCalibur Flow Cytometer (BD Biosciences). FlowJo software was used to analyze the % of cells in different cell cycle phases (synthesis phase, growth phases).

2.2.7 SNP Genotyping

Patient germline DNA was isolated and genotyped on the Illumina Infinium 610K array, as previously described ²⁷. Genotypes were predicted by STRUCTURE ²⁸ analysis to have greater than 80% European ancestry. The 1000 Genomes Project ²³ was used as reference, imputation was completed with *mach* and *minimac* in a two-step process ²⁹. SNPs within carboplatin and paclitaxel drug pathway genes were extracted by using the web-based SNP Nexus software.

2.2.8 RNA Isolation and Gene expression assays

Total RNA was isolated from LCLs before drug treatment using RNeasy Plus Mini Kit (QIAGEN, USA) according to manufacturer's protocol. RNA quantity and concentration were measured using

NanoDrop 2000 UV-Vis spectrophotometer (ThermoScientific, USA). The absorbance ratio between absorbances at 260 nm and 280 nm (A_{260}/A_{280}) was computed for assessing RNA sample purity. A_{260}/A_{280} ratio between 1.8-2.1 was considered as highly pure RNA. Further, RNA Quality check was performed using Agilent Bioanalyzer (Agilent RNA 6000 Pico kit) and estimated by calculating RNA Integrity Number (RIN) with 2100 expert software version B.02.02. RIN greater than 7 was considered indicating good quality RNA. The total RNA was reverse transcribed to cDNA using High Capacity cDNA Reverse Transcription Kit (Applied Biosystems, USA) according to manufacturer's protocol.

Expression of key pharmacokinetic and pharmacodynamics pathway genes of carboplatin and paclitaxel was determined using the TaqMan® Low Density Array (TLDA) cards (Applied Biosystems, USA). Briefly, each TLDA card was custom designed with pre-loaded assays for measuring the mRNA expression of selected genes that includes influx and efflux transporters, DNA repair gene, detoxifying gene for carboplatin and genes for microtubule associated protein, transporters for paclitaxel, along with the housekeeping genes GAPDH and Actin β (**Table 2.1**). Each TLDA card consists of eight separate loading ports which can fill into 48 separate wells for each sample. 24 different genes can be assayed in duplicates in a total of 384 wells per card. Therefore, each card could analyze the expression of 8 different samples. High-quality cDNA and equal volume of 2X TaqMan Universal PCR Master Mix was loaded in to the TLDA card along with 100 μ l of the sample-specific PCR mix. To distribute reaction mix to the reaction wells properly, the TLDA cards were then centrifuged twice for 1 minute at 1200 rpm using a Sorvall Legend T Centrifuge that has Sorvall Custom Buckets with TLDA card holder. The cards were then sealed by the TaqMan Array Micro Fluidic Card Sealer and real-time quantitative polymerase chain reaction (qRT-PCR) was performed on microfluidic card thermal cycling block of Applied Biosystems 7900HT Fast Real-time PCR System (Applied Biosystems). Thermal cycling conditions were: initial denaturation of 10 minutes at 94.5°C followed by 35 cycles of: annealing

and extension of 2 minutes at 50°C, and denaturation at 30 seconds at 97°C and finally 1 minute at 59.7°C for 40 cycles. For every run, Ct (cycle threshold) values or the cycle number required for the fluorescent signal to cross the background level were determined for each gene and used for further analysis. Average Ct values representing the mRNA expression levels were normalized to the housekeeping/control gene (GAPDH).

Table 2.1 Genes included in TaqMan Low-Density array (TLDA) card.

Gene	Gene Name	Pathway (s)	Role
ABCB1	ATP-binding cassette, sub-family B (MDR/TAP), member 1	Paclitaxel	Transporter
ABCC1	ATP-binding cassette, sub-family C (CFTR/MRP), member 1	Carboplatin, paclitaxel	Transporter
ABCC2	ATP-binding cassette, sub-family C (CFTR/MRP), member 2	Carboplatin, paclitaxel	Transporter
ABCG2	ATP-binding cassette, sub-family G (WHITE), member 2	Carboplatin, paclitaxel	Transporter
ACTB	actin, beta	Control	House-keeping gene
ATP7A	ATPase, Cu ⁺⁺ transporting, alpha polypeptide	Carboplatin	Transporter
ATP7B	ATPase, Cu ⁺⁺ transporting, beta polypeptide	Carboplatin	Transporter
BIRC5	baculoviral IAP repeat-containing 5	Paclitaxel	Anti-apoptotic
BIVM/ERCC5	basic, immunoglobulin-like variable motif containing	Carboplatin	DNA repair
BRCA1	breast cancer 1, early onset	Carboplatin, paclitaxel	Genomic stability/Tumor suppressor
CCNB1	cyclin B1	Paclitaxel	Cell cycle regulator
EGFR	epidermal growth factor receptor (erythroblastic leukemia viral (v-erb-b) oncogene homolog, avian)	Paclitaxel	Cell proliferation
ERCC1	excision repair cross-complementing rodent repair deficiency, complementation group 1 (includes overlapping antisense sequence)	Carboplatin	DNA repair
ERCC2	excision repair cross-complementing rodent repair deficiency, complementation group 2	Carboplatin	DNA repair
ERCC6	excision repair cross-complementing rodent repair deficiency, complementation group 6	Carboplatin	DNA repair
FOXC2	forkhead box C2 (MFH-1, mesenchyme forkhead 1)	Carboplatin, paclitaxel	Differentiation
FOXL1	forkhead box L1	Carboplatin, paclitaxel	Differentiation
GAPDH	glyceraldehyde-3-phosphate dehydrogenase-like 6; hypothetical protein LOC100133042; glyceraldehyde-3-phosphate dehydrogenase	Control	House-keeping gene
GSTM1	glutathione S-transferase mu 1	Carboplatin	Xenobiotic metabolism
GSTM3	glutathione S-transferase mu 3 (brain)	Carboplatin	Xenobiotic metabolism
GSTP1	glutathione S-transferase pi 1	Carboplatin	Xenobiotic metabolism
GSTT1	glutathione S-transferase theta 1	Carboplatin	Xenobiotic metabolism
HMGB1	high-mobility group box 1; high-mobility group box 1-like 10	Carboplatin	Pt-DNA adduct repair
KISS1	KiSS-1 metastasis-suppressor	Carboplatin	Metastasis suppressor <i>gene</i>
MAD2L1	MAD2 mitotic arrest deficient-like 1 (yeast)	Paclitaxel	Microtubule- associated proteins
MAP2	microtubule-associated protein 2	Paclitaxel	Microtubule- associated proteins
MAP4	microtubule-associated protein 4	Paclitaxel	Microtubule- associated proteins

MAPT	microtubule-associated protein tau	Paclitaxel	Microtubule- associated proteins
MLH1	mutL homolog 1, colon cancer, nonpolyposis type 2 (E. coli)	Carboplatin	DNA repair
MPO	myeloperoxidase	Carboplatin	Xenobiotic metabolism
MSH2	mutS homolog 2, colon cancer, nonpolyposis type 1 (E. coli)	Carboplatin	DNA repair
MSH6	mutS homolog 6 (E. coli)	Carboplatin	DNA repair
NQO1	NAD(P)H dehydrogenase, quinone 1	Carboplatin	Xenobiotic metabolism
NR1I2	nuclear receptor subfamily 1, group I, member 2	Paclitaxel	Xenobiotic metabolism
PMS2	PMS2 postmeiotic segregation increased 2 (S. cerevisiae)	Carboplatin	DNA repair
POLB	polymerase (DNA directed), beta	Carboplatin	DNA repair
POLH	polymerase (DNA directed), eta	Carboplatin	DNA repair
REV3L	REV3-like, catalytic subunit of DNA polymerase zeta (yeast)	Carboplatin	DNA repair
SLC22A7	solute carrier family 22 (organic anion transporter), member 7	Paclitaxel	Transporter
SLC31A1	solute carrier family 31 (copper transporters), member 1	Carboplatin	Transporter
SOD1	superoxide dismutase 1, soluble	Carboplatin	Xenobiotic metabolism
TP53	tumor protein p53	Carboplatin, paclitaxel	Tumor suppressor
XPA	xeroderma pigmentosum, complementation group A	Carboplatin	DNA repair
XRCC1	X-ray repair complementing defective repair in Chinese hamster cells 1	Carboplatin	DNA repair
XRCC2	X-ray repair complementing defective repair in Chinese hamster cells 2	Carboplatin	DNA repair
XRCC3	X-ray repair complementing defective repair in Chinese hamster cells 3	Carboplatin	DNA repair
XRCC5	X-ray repair complementing defective repair in Chinese hamster cells 5 (double-strand-break rejoining)	Carboplatin	DNA repair

2.2.9 Data analysis

All statistical analyses were performed using GraphPad Prism software version 7.0 and R statistical analysis software version 3.3.1, a free software environment as well as a programming language for statistical computing and graphics. Drug IC_{50} (effective dose that inhibits 50% of the cells), area under the survival curve (AUC), EC_{50} (concentration required to induce caspase 3/7 activity by 50%) and the area under the relative caspase activity curve (AUC) were also calculated using four parameter logistic dose response curves fitting to the *in vitro* drug response measurements (cell survival and caspase3/7 activity assays) ²⁷.

To cluster cell lines based on cytotoxicity parameters, unsupervised K-means clustering was performed using the algorithm of Hartigan and Wong (1979) to identify the most-resistant and relatively sensitive subgroups of cell lines ³⁰.

We used CalcuSyn software version 2.0 (Biosoft) based on Chou-Talalay method for the analysis of drug combination treatment ³¹. This quantitative analysis of combination drug effects is based on the median-effect equation, derived from the mass-action law principle, that estimates the following: 1) combination index (CI) as the measure of the effect of each individual drug in combination treatment and 2) dose reduction index (DRI) or the reduction in dose requirement of a drug in combination treatment compare to single drug treatment and also 3) the required dose to achieve IC_{50} in combination (carboplatin + paclitaxel) treatment for each cell line ³¹. Combination index (CI) is the quantitative measure of interactions between multiple drug effects interactions derived from the median-effect equation or Unified theory of the law of Mass-Action, where $CI < 1$ is defined as synergistic, $CI > 1$ as antagonistic and $CI = 1$ as additive ³¹. Additionally, DRI or dose reduction index (DRI) of an individual drug in combination treatment is a measure of estimated reduction of its dose at a given effect level compared to when the drug is administered as a single agent. Therefore, higher DRI corresponds to lower dose requirement which implies reduced toxicity at the increased effect in synergistic combination ³¹.

The association of SNPs with *in vitro* drug response phenotypes IC₅₀ or EC₅₀ was evaluated with linear models. For annotation of results across gene regions, SNPs were mapped to genes within 2KB using Biofilter (assembly CRCh37.p10, genome build 104.0) and SNP Nexus, a web-based application for the selection of functionally relevant Single Nucleotide Polymorphisms (SNP) for large-scale genotyping studies ²⁷.

Gene expression data generated using TaqMan Low Density arrays (TLDA) was analyzed using Partek Genomics Suite software v6.6 to perform differential expression analysis and to find correlation between drug pathway gene expression signatures with drug chemo-sensitivity parameters using Analysis of variance (ANOVA) model. Heatmaps were generated for each set of pathway genes (Carboplatin and paclitaxel) using unsupervised hierarchical clustering (HC) analysis based on the genes expressed differentially. Gene expression data was standardized: shifted to mean=0 and scaled to standard deviation=1 prior to Hierarchical clustering analysis.

2.3 Results

2.3.1 *In vitro* profiling of carboplatin and paclitaxel chemo-sensitivity

A total of 112 EBV-transformed ovarian cancer cell lines were created. Among these, 81 cell lines were successfully treated with carboplatin and paclitaxel as single agents and as combination.

Figure 2.1 demonstrates the quality control (QC) steps undertaken to check for variability or batch effects in the data owing to variations of experimental conditions. Briefly, a few cell lines were randomly selected and cytotoxicity assays were performed at multiple time points to check for consistency. Our results showed there were no batch effects observed for our drug cytotoxicity or caspase activity data. Furthermore, we divided the dataset into two subsets and performed statistical verification of any batch effect, as previously reported ²⁷.




Figure 2.1 Workflow representing Data QC (quality check) of chemo-sensitivity assay results

Table 2.2 provides a summary of all the *in vitro* chemo-sensitivity measures assessed for single agent and combination treatments, including IC₅₀, AUC, Caspase AUC and Caspase EC₅₀. Results from our *in vitro* chemo-sensitivity analysis showed wide variability in response to carboplatin and paclitaxel treatment as single agent among LCLs from ovarian cancer patients. For carboplatin cytotoxicity, IC₅₀ and AUC showed fold changes of 8.34 and 3.84 between the lowest (most-sensitive) and highest (most-resistant) values, respectively whereas the fold changes were 7.72 and 5.65, respectively for paclitaxel.

Boxplots in **Figure 2.2** represent the distribution of log transformed drug-sensitivity parameters across the LCL panel. **Figure 2.3** depicts a scatterplot matrix showing the correlation between the

observed drug chemo-sensitivity parameters in EOC-LCLs. We observed a significantly high ($r>0.5$; $p<0.0001$) correlation between the following: caspase AUC values for combination treatment with caspase AUC for carboplatin ($r= 0.86$) and paclitaxel ($r= 0.84$) as single agent treatment; paclitaxel IC₅₀ vs paclitaxel AUC ($r=0.71$); paclitaxel IC₅₀ vs Combination AUC ($r=0.59$); Combination AUC vs paclitaxel AUC ($r=0.7$); Combination AUC vs carboplatin AUC ($r=0.59$); and carboplatin caspase vs paclitaxel caspase ($r=0.74$) (**Figure 2.3**). As expected, IC₅₀ and AUC values were negatively correlated with corresponding caspase AUC values for a treatment regimen indicating increase in caspase 3/7 activity and apoptosis in the dying cells.

Our results demonstrate a wide range of diversity in response to treatment with carboplatin and paclitaxel single agent treatment as well as strong correlation with combination treatment. Our caspase 3/7 activity data corroborated with cytotoxicity data (data not shown).

We also performed flow cytometry analysis of the cell lines prior to performing chemosensitivity assays to assess cell cycle phases based on quantification of DNA content. Our results showed no correlation between cell cycle phases and drug cytotoxicity revealing drug response was not influenced by either cell viability or cell cycle phases (**Table 2.3**).

Table 2.2 Summary of *in vitro* phenotype measures in LCLs from EOC patients.

Drug	Phenotype	N	Mean	Minimum	Maximum	Fold Change (Max/Min)
Carboplatin	IC ₅₀ (μM)	77	52.56	13.63	113.24	8.31
Carboplatin	AUC	81	6165.32	2759	10598	3.84
Carboplatin	Caspase_AUC	79	288.23	100	640	6.4
Carboplatin	Caspase_EC ₅₀ (μM)	79	27.12	6	96	16
Paclitaxel	IC ₅₀ (nM)	76	19.33	5.84	45.12	7.72
Paclitaxel	AUC	81	3159.52	1165	6584	5.65
Paclitaxel	Caspase_AUC	79	173.49	80	330	4.13
Paclitaxel	Caspase_EC ₅₀ (μM)	79	10.75	2.9	56	19.31
Combination	AUC	81	2917.99	1262	6568	5.2
Combination	Caspase_AUC	79	139.35	62	270	4.35
Combination	Caspase_EC ₅₀	79	10.73	3.7	41	11.08




Figure 2.2 Boxplots representing distribution of a) drug cytotoxicity and b) caspase activity following treatment.

Cytotoxicity values were log-transformed (represented in the y-axes) and boxplots were generated using the R statistical packages. (Carb = Carboplatin; Pacl = Paclitaxel; Comb = Combination; Casp = Caspase)




Figure 2.3 Scatterplot matrix representing pairwise correlations between IC₅₀ and AUC estimates of paclitaxel and carboplatin single-agent and combination treatment.

Cytotoxicity estimates were log-transformed and a scatterplot matrix was generated using the R graphing package. Scatterplots, r values and corresponding significance (p-values) for each pairwise correlation are included. Marginal plots on the diagonal axis represent histograms of the individual cytotoxicity parameters. Significance levels are associated with the following symbols and represent p-values (0.001, 0.01, 0.05, 0.1, 1) <=> symbols ("***", "**", "*", ".", " ")


Table 2.3 Summary of correlation between cell viability & cell cycle phases and drug IC₅₀

Phenotype	Minimum	Mean	Maximum	sd	Carboplatin. IC ₅₀	Paclitaxel. IC ₅₀	Combination. IC ₅₀
Cell viability %	80	91.33	97	3.81	r= 0.2156	r = 0.1135	r= 0.1375
S-Phase %	12.3	24.843	38.36	4.997	r=0.042	r=-0.1049	r=-0.099
G1- Phase %	41.62	63.072	76.35	6.906	r=-0.126	r=-0.142	r=-0.165
G2- Phase %	1.01	11.007	34.95	5.654	r=0.051	r=0.254	r=0.218

2.3.2 K-Means clustering identified resistant subgroups

K-Means clustering analysis was performed to identify clusters or subgroups based on our based on our *in vitro* cytotoxicity parameters observed in the EOC-LCLs ³⁰. Three (3) distinctly different drug response subgroups were identified (**Figure 2.4**). We then focused on identifying differences in response to carboplatin, paclitaxel and combination treatments between the 3 subgroups generated using K-Means clustering. **Table 2.4** provides numerical summary of the drug response parameters for each K-means cluster. Our results showed that cytotoxicity AUC and IC₅₀ for carboplatin, paclitaxel and combination were significantly higher ($p < 0.0001$) in Cluster 2 (resistant subgroup) compared to other cell lines. 7 cell lines (Cluster 2) were identified as most resistant to the drug treatments. Cluster1 (n= 26) was identified as the most sensitive while cluster 3 (n=84) is the intermediate subgroup (**Figure 2.4**). Pvalues of Kruskal-Wallis test by ranks are also provided for each drug response phenotype (**Figure 2.5**).

a)



b)




Figure 2.4 Results from Kmeans clustering of *in vitro* cytotoxicity data

a) Heatmap representing Kmeans clusters generated using *in vitro* cytotoxicity assay data on the LCL panel. K Means clustering (k=3) identified 7 cell lines (middle panel) as most resistant to the drug treatments; b) Bar plots showing subgroups of cell lines in the LCL panel clustered using Kmeans clustering analysis (k=3)




Figure 2.5 Plots showing differences in response to carboplatin, paclitaxel and combination treatments between the clusters generated using *in vitro* cytotoxicity assay data on the LCL panel.

K-Means clustering analysis was performed to identify clusters or subgroups based on our based on our *in vitro* cytotoxicity profiling data in the cell lines.

Table 2.4 Summary of drug response phenotypes in subgroups identified by Kmeans clustering.

Drug	Phenotype	Cluster 1 (n=26)	Cluster 2 (n=7)	Cluster 3 (n=48)	P (Kruskal-Wallis)
Carboplatin	IC ₅₀ (μM)	34.99	236.43	99.69	< 0.0001
Paclitaxel	IC ₅₀ (nM)	8.98	125.57	16.12	< 0.0001
Carboplatin	AUC	4907.73	8580.43	6494.31	< 0.0001
Combination	AUC	2120.5	4706	3089.21	< 0.0001
Paclitaxel	AUC	2255.88	5191.43	3352.67	< 0.0001

2.3.3 Analysis of carboplatin-paclitaxel combination treatment using CI theorem

Furthermore, we investigated the *in vitro* interactions between the two drugs, carboplatin and paclitaxel, in combination compared to use as single agent in our cell line panel of 81 EOC-LCLs using Chou-Talalay's Combination Index-Isobologram Theorem to evaluate synergism, antagonism or additive effects ³¹. We studied the effect of combination (carboplatin + paclitaxel) treatment in our cell line panel of 81 EOC-LCLs. Results showed, for either of the drugs, reduced dose was required to achieve a similar effect (% kill) in combination treatment compared to single agent treatment for majority of the cell lines. We also observed large variability in the IC₅₀ of single-agent treatment compared to its IC₅₀ when used in drug combination (**Figure 2.6**). Our observed CI values showed wide variation (0.14-5.51) within the cell line panel. As mentioned in the Methods section, where CI<1 is defined as synergistic, CI>1 as antagonistic and CI=1 as additive ³¹. We also observed that dose reduction (DRI) for a drug is low in cell lines where the combination results in predicted antagonistic drug-drug interaction (high CI value). High DRI corresponds to lower dose requirement, as described earlier, which suggests reduced toxicity corresponding to increasing synergism in combination treatments ³¹. We also observed high negative correlation between DRI and CI for carboplatin IC₅₀ (r=- 0.732), as well as for paclitaxel IC₅₀ (r=-0.641). **Figure 2.7** summarizes the CI and DRI values in the diagonal box plots and provides pair-wise correlations between CI and DRI values at predicted IC₂₅, IC₅₀ and IC₇₅. Thus, our DRI calculations revealed lower dose requirements to achieve similar anti-cancer effects when

the drugs were used in combination, compared to single-agent regimens, based on our observations from their reduced predicted IC_{50} and IC_{75} values in combination treatments (**Table 2.5**).




Figure 2.6 Representation of variation of the relative IC₅₀s of treatment with single agents and drug combination that indicates variation across LCL panel. a) carboplatin; b) paclitaxel.

Table 2.5 CI and DRI as measures of Synergism and Antagonism of drug combination treatment in LCLs from EOC patients.

	Drug Phenotype	Minimum	Mean	Maximum
CI	IC ₅₀	0.14	1.03	5.51
DRI	Carboplatin IC ₅₀	0.57	2.84	14.04
DRI	Carboplatin IC ₇₅	0.00	2.88	11.23
DRI	Paclitaxel IC ₅₀	0.26	2.74	15.26
DRI	Paclitaxel IC ₇₅	0.00	2.14	7.79

Combination Index (CI) and Dose-Reduction Index (DRI) values were calculated using Compusyn software program that applies Median-Effect Principle (by Chou) and the Combination Index-Isobologram Theorem (by Chou-Talalay). CI < 1, Synergism; CI = 1, Additive; CI > 1, Antagonism.




Figure 2.7 Scatterplots showing correlation between combination index (CI) and dose reduction index (DRI) values at IC₅₀.
Diagonal axis shows boxplots of the distribution of each parameter.

2.3.4 SNPs in pathway genes associated with chemo-sensitivity

Genotype call rates observed for all samples were >95%, and minor allele frequency (MAF) ≥ 0.01 . We performed a genotype-phenotype association analysis between SNPs in carboplatin and platinum pharmacokinetic and pharmacodynamics pathway genes extracted from our GWAS study and our observed *in vitro* chemo-sensitivity measures. **Table 2.6** provides details of the top 15 pathway SNPs that were found significantly associated with *in vitro* cytotoxicity phenotypes and their predicted functional effects. We identified significant association between pathway genes of drugs with response to carboplatin, paclitaxel as single agent and as combination therapy. At $p < 0.01$, 85 carboplatin pathway gene mutations were significantly associated with carboplatin IC_{50} , 50 paclitaxel pathway genes were associated with paclitaxel IC_{50} while a total of 67 genetic variations were significantly associated with combination treatment chemosensitivity.

Analysis of association between genetic variations in drug PK/PD pathway genes and *in vitro* caspase activity revealed 46, 23, 63 mutations associated with caspase activity in carboplatin single agent, paclitaxel single agent and combination treatment, respectively. **Table 2.7** lists the top 15 Pathway SNPs that were found most significantly associated with caspase activity phenotypes and their predicted functional effects.

2.3.4.1 SNPs in genes associated with drug pharmacokinetics

Top among the SNPs in drug disposition (PK) genes significantly ($p < 0.05$) associated with carboplatin cytotoxicity were the SNPs rs186868889 ($p = 0.001$) and rs72659636 ($p = 0.001$) in the gene *ABCG2* and rs182123265 (*ABCC2*; $p = 0.002$). Most significant SNP versus paclitaxel single-agent cytotoxicity association was found among rs1153867 (*ABCC1*) and rs9472030 ($p = 0.002$). On the other hand, cytotoxicity of carboplatin-paclitaxel combination therapy was highly significantly associated with the following SNPs: rs183572778, rs17224787, rs62140866, rs17224528 (*MSH2*); rs3887412 ($p = 0.0034$), rs35598 ($p = 0.0035$), rs35599 ($p = 0.0035$) (*ABCC1*). The top PK pathway gene mutations associated with caspase activity following carboplatin

treatment include the transporter genes *ATP7B* and the xenobiotic metabolism gene *GSTM1*. Mutations in the genes *MAP4*, *MAPT*, *MAP2*, were among the top caspase activity-associated PK gene variations following paclitaxel single-agent treatment. Finally, genetic variations in *ABCC1*, *BIRC5*, *MAP4*, and *GSTP1* were highly associated with caspase activity following treatment with carboplatin and paclitaxel as combination.

2.3.4.2 SNPs in genes related to pharmacodynamics response

The following SNPs in pharmacodynamics (PD) genes were most significantly associated with carboplatin cytotoxicity: rs1425118 (p=0.003), rs55911615 (p=0.003), rs6749415 (p=0.003) in *XRCC5*, and rs75475960 (*MSH2*; p=0.003). On the other hand, significantly high SNP vs paclitaxel single-agent cytotoxicity association was found among *TP53* SNPs rs17887200, rs17881556 and rs17886760 (p<0.001 for all three SNPs), and the *EGFR* SNPs rs7779645, rs7804688, rs6947594, rs6593209 (p<or =0.001 for all three SNPs). The top PD pathway gene mutations associated with caspase activity following carboplatin treatment include DNA repair genes *XRCC5*, *XRCC1*, *ERCC2*, and *ERCC1*, as well as *REV3L*, *MSH6* and *KISS1*. Top PD pathway gene mutations associated with caspase activity following carboplatin treatment include: mutations in the genes *BIVM/ERCC6*, *EGFR* and *BRCA1* were among the top caspase activity-associated PD variations following paclitaxel single-agent treatment. Whereas, genetic variations in *BIRC5*, *KISS1*, and *EGFR* were highly associated with caspase activity following treatment with carboplatin and paclitaxel as combination.

Table 2.6 Top pathway SNPs found most significantly associated with *in vitro* cytotoxicity parameters.

	Ch r	chromPositio n	dbSNP ID	Gene/Gene	Allele 1	Allele 2	Directio n	P-value (SNP vs IC50)	Predicted function
Carboplatin IC50									
	2	217037273		<i>XRCC5</i>	t	c	-1	0.001	
	4	88994224	rs18686888 9	<i>ABCG2</i>	t	g	-1	0.001	
	4	88999267	rs72659636	<i>ABCG2/PKD2</i>	c	g	-1	0.001	3downstrea m
	7	6061988		<i>PMS2/EIF2AK1</i>	a	g	1	0.001	3utr
	2	217028174		<i>XRCC5</i>	a	g	-1	0.002	
	10	101573872	rs18212326 5	<i>ABCC2</i>	a	g	1	0.002	
	2	216958899	rs1425118	<i>XRCC5</i>	c	g	1	0.003	
	2	217028646		<i>XRCC5</i>	t	g	-1	0.003	
	2	216958412		<i>XRCC5</i>	a	g	-1	0.003	
	2	216958402	rs6749415	<i>XRCC5</i>	a	g	1	0.003	
	2	216968340	rs55911615	<i>XRCC5/TMEM16 9</i>	a	g	1	0.003	3downstrea m
	2	47616643	rs75475960	<i>MSH2</i>	c	g	1	0.003	
	2	216970326		<i>XRCC5</i>	t	c	-1	0.003	
	2	216972934		<i>XRCC5/XRCC5</i>	a	g	1	0.003	5upstream
Paclitaxel IC₅₀									
	17	7571071	rs17887200	<i>TP53</i>	t	c	-1	<0.001	3downstream
	17	7567681		<i>TP53</i>	t	c	1	<0.001	
	17	7568628		<i>TP53</i>	a	g	1	<0.001	
	7	55185302	rs7779645	<i>EGFR</i>	a	t	-1	<0.001	
	7	55190098	rs7804688	<i>EGFR</i>	a	t	-1	<0.001	

	7	55203297	rs6947594	EGFR	a	c	-1	<0.001	
	7	55203838		EGFR	a	g	1	<0.001	
	17	7570869	rs17881556	TP53	t	c	-1	0.001	3downstream
	17	7570978	rs17886760	TP53	t	c	-1	0.001	3downstream
	17	7570823		TP53	c	g	1	0.001	3downstream
	17	7570956		TP53	a	g	1	0.001	3downstream
	16	16203202	rs1153867	ABCC1	t	g	1	0.001	
	7	55200625	rs6593209	EGFR	a	g	-1	0.001	
	6	43271237	rs9472030	SLC22A7/CRIP3	t	g	-1	0.002	3downstream
	7	55208284		EGFR	a	c	1	0.002	
Combination									
	2	48010163		MSH6	t	c	1	0.0007	5upstream
	2	47636927	rs183572778	MSH2	a	g	-1	0.0011	
	16	16173221		ABCC1	t	c	-1	0.0012	S667S
	2	47701362	rs17224787	MSH2	a	t	1	0.0019	
	2	47682634	rs62140866	MSH2	a	c	1	0.0021	
	2	47684043	rs17224528	MSH2	a	g	1	0.0022	
	16	16173516		ABCC1	t	c	-1	0.0034	
	16	16175030	rs3887412	ABCC1	a	t	1	0.0034	
	16	16182528		ABCC1	t	c	-1	0.0034	
	16	16183045		ABCC1	a	g	-1	0.0034	
	16	16159023	rs35598	ABCC1	a	g	1	0.0035	
	16	16159061	rs35599	ABCC1	t	c	1	0.0035	
	16	16168608		ABCC1	t	c	-1	0.0039	
	16	16169465		ABCC1	t	g	-1	0.0039	
	16	16170771		ABCC1	a	g	-1	0.0039	

Table 2.7 Top pathway SNPs found significantly associated with *in vitro* combination treatment caspase activity.

	Chr	chromPosition	dbSNP ID	Gene	Allele1	Allele2	Direction	P-value (SNP vs Caspase)	Predicted function
Carboplatin									
	1	110211681	rs428135	<i>GSTM1</i>	C	g	1	0.001	
	2	217003152	rs2303400	<i>XRCC5</i>	T	c	1	0.002	
	6	111801193		<i>REV3L</i>	A	g	1	0.003	
	13	52497540		<i>ATP7B</i>	A	g	-1	0.003	
	19	44062781		<i>XRCC1</i>	D	r	-1	0.003	
	19	45837542		<i>ERCC2</i>	C	g	1	0.003	
	19	45841318		<i>ERCC2</i>	A	g	1	0.003	
	2	217012092	rs2160981	<i>XRCC5</i>	A	g	1	0.004	
	19	44041363		<i>XRCC1</i>	T	c	-1	0.004	3' downstream
	19	45931387	rs28586606	<i>ERCC1</i>	A	g	1	0.004	
	19	45936292		<i>ERCC1</i>	T	c	-1	0.004	
	1	110291406		<i>GSTM3</i>	T	c	-1	0.005	3' downstream
	1	204181317	rs72749759	<i>KISS1</i>	A	c	-1	0.005	
	2	48012375		<i>MSH6</i>	C	g	-1	0.005	
	2	48022688	rs111861797	<i>MSH6</i>	T	c	1	0.005	
Paclitaxel									
	10	96845842		<i>CYP2C8</i>	D	r	-1	0.003	
	3	47977447		<i>MAP4</i>	D	r	1	0.005	
	17	44072651		<i>MAPT</i>	A	g	1	0.005	
	17	44090778		<i>MAPT</i>	A	t	1	0.007	
	17	44090779		<i>MAPT</i>	A	c	1	0.007	
	2	210512893		<i>MAP2</i>	A	g	-1	0.008	
	2	210574383		<i>MAP2</i>	A	g	1	0.008	
	2	210596950	rs114531804	<i>MAP2</i>	A	g	1	0.008	3' UTR
	7	55197234	rs79884022	<i>EGFR</i>	T	c	-1	0.008	

	7	55206022		<i>EGFR</i>	A	g	1	0.008	
	17	41250221		<i>BRCA1</i>	D	r	-1	0.008	
	17	41260985		<i>BRCA1</i>	C	g	-1	0.008	
	2	210534297	rs141068360	<i>MAP2</i>	T	g	-1	0.009	
	3	47896738		<i>MAP4</i>	A	c	1	0.009	Coding (K1087N)
	3	47918295		<i>MAP4</i>	A	g	1	0.009	
Combination									
	17	76238765		<i>BIRC5</i>	T	c	-1	<0.001	3' downstream
	17	76239023	rs7223704	<i>BIRC5</i>	T	c	1	<0.001	3' downstream
	17	76231842		<i>BIRC5</i>	T	c	-1	0.002	
	3	48067426	rs184824121	<i>MAP4</i>	T	c	-1	0.003	
	1	204150976		<i>KISS1</i>	T	c	1	0.005	
	7	55124701	rs6593202	<i>EGFR</i>	T	c	-1	0.005	
	11	67362385	rs7108038	<i>GSTP1</i>	A	g	-1	0.005	
	16	16195257		<i>ABCC1</i>	A	g	1	0.005	
	16	16206501	rs45499397	<i>ABCC1</i>	C	g	-1	0.005	
	17	76230981	rs8064778	<i>BIRC5</i>	C	g	1	0.005	
	17	76231434		<i>BIRC5</i>	A	t	-1	0.005	
	11	67358076		<i>GSTP1</i>	T	g	1	0.006	
	11	67359090	rs7939505	<i>GSTP1</i>	C	g	-1	0.006	
	11	67359718	rs6591259	<i>GSTP1</i>	A	g	-1	0.006	
	11	67359947	rs7937159	<i>GSTP1</i>	T	g	-1	0.006	

2.3.5 Differential GEP signatures of drug response

By comparing gene expression versus response phenotype using the Spearman rank-order correlation analysis, several key pathway genes were found significantly associated with drug chemosensitivity phenotype measure (details provided in **Table 2.8**).

Pre-treatment gene expression levels of candidate genes involved in carboplatin and paclitaxel PK/PD pathway were assayed using TaqMan Low Density Arrays. Average Ct values for each gene were then normalized using average Ct value of the housekeeping/control gene (GAPDH). Data was log-transformed and used for analysis of association with phenotype measures. Genes showing no expression in >5% cell lines were filtered out prior to analysis. Differential gene expression analysis was performed using the remaining genes to identify gene expression changes that were highly correlated with drug chemo-sensitivity. Clustering analysis using the most differentially expressed genes revealed distinct clusters/subgroups within the LCL panel. **Figure 2.8 and 2.9** show heatmaps that cluster the EOC LCL panel based on most differentially expressed pathway genes for carboplatin and paclitaxel, respectively.

Expression of the following genes were highly correlated with carboplatin cytotoxicity: *ERCC2* (p=0.015); *GSTM1* (p=0.0097); *GSTT1* (p=0.0086); *NQO1* (p=0.0497), while expression levels of the following genes were correlated with response to paclitaxel single agent treatment: *BIRC5* (p=0.003); *FOXL1* (p=0.0148); *FOXC2* (p=0.0034). When compared to the phenotype measures of carboplatin-Paclitaxel combination treatment, CI values at IC₅₀ was correlated with the carboplatin pathway genes *ATP7A* and *ATP7B*. In addition, DRI values at IC₂₅ were correlated with the following carboplatin pathway genes: *POLH* (p=0.022); *ATP7A* (p=0.042) and *FOXC2* (p=0.032), as well as the paclitaxel pathway genes *FOXC2* (p=0.0114), *BIRC5* (p=0.0237) and *EGFR* (p=0.0458).


Spearman correlation analysis between caspase activity and pathway gene expression revealed trends towards significance for expression of the genes *GSTP1* (p=0.08) and *NQO1* (p=0.096) for

carboplatin treatment and *FOLXL1* (p=0.07), *BIRC5* (p=0.07) and *EGFR* (p=0.08) for caspase activity induced by paclitaxel single agent treatment (**Table 2.8**).

Table 2.8 Summary of results from Spearman correlation analysis between TLDA gene expression and drug chemo-sensitivity outcomes (p<0.1).

Drug	Phenotype	Column ID	p-value (correlation)
Carboplatin	IC50	<i>ERCC2</i>	0.035
		<i>XPA</i>	0.087
	MTT.AUC	<i>GSTM1</i>	0.010
		<i>ERCC2</i>	0.002
		<i>MSH2</i>	0.040
		<i>TP53</i>	0.083
		<i>GSTP1</i>	0.023
		<i>NQO1</i>	0.032
		<i>MSH6</i>	0.035
		<i>XRCC2</i>	0.037
		<i>SOD1</i>	0.039
		<i>ATP7A</i>	0.039
		<i>POLB</i>	0.043
		<i>FOXL1</i>	0.045
Paclitaxel	Caspase.AUC	<i>GSTP1</i>	0.085
	Caspase.EC50	<i>NQO1</i>	0.096
	IC50	<i>BIRC5</i>	0.094
		<i>BIRC5</i>	0.002
	MTT.AUC	<i>FOXC2</i>	0.003
		<i>FOXL1</i>	0.011
		<i>MAD2L1</i>	0.093
	Caspase.EC50	<i>FOXL1</i>	0.070
		<i>BIRC5</i>	0.074
Combination	MTT.AUC	<i>GSTM1</i>	0.017
		<i>ERCC6</i>	0.053
		<i>BIRC5</i>	0.039
		<i>FOXC2</i>	0.064
		<i>NQO1</i>	0.085
		<i>FOXL1</i>	0.090
	Casp.AUC	<i>GSTP1</i>	0.022
	Casp.EC50	<i>XPA</i>	0.017
		<i>HMGB1</i>	0.029
		<i>BIRC5</i>	0.050
		<i>ERCC1</i>	0.061
		<i>NQO1</i>	0.064
		<i>FOXC2</i>	0.078

a)



b)





Figure 2.8 Heatmap showing hierarchical clustering of the LCL panel based on differentially expressed carboplatin pathway genes. a) All pathway genes; b) Pathway genes most significantly associated with cytotoxicity ($p < 0.1$).
Columns represent cell lines; Rows represent carboplatin pathway gene analyzed using TLDA arrays.

a)



b)




Figure 2.9 Heatmap showing hierarchical clustering of the LCL panel based on differentially expressed paclitaxel pathway genes. a) All pathway genes; b) Pathway genes most significantly associated with cytotoxicity ($p < 0.1$).
Columns represent cell lines; Rows represent paclitaxel pathway gene analyzed using TLDA arrays.

2.4 Discussion and Conclusions

EOC accounts for 6% of all cancer deaths among women ³. Despite the routine use of carboplatin-paclitaxel combination treatment, one of the biggest challenges faced by clinicians is wide inter-patient variation in clinical response, ^{7, 32} as well as toxicities including myelosuppression and gastrointestinal toxicities ⁷ and drug resistance ³². Approximately 80% ovarian epithelial cancer patients relapse after first line based chemotherapy.

We hypothesized that modelling drug resistance using *in vitro* model systems will enable identification of genetic variants and gene expression signatures significantly associated with drug response. These drug response markers derived from *in vitro* models may then be validated using patient data to identify actionable genetic changes that influence drug resistance and treatment outcomes. In the current study, we used a panel of 81 LCLs derived from human ovarian cancer patients to identify key mutations and gene expression changes associated with carboplatin and paclitaxel response, both as single agents and when used as combination.

Our *in vitro* chemo-sensitivity assays showed wide variability in response to carboplatin and paclitaxel single agent and combination treatment, that mirrors the wide variability in clinical response among ovarian cancer patient samples. We also performed caspase activity assays to demonstrate that drug cytotoxicity reflected alterations in caspase-3 and caspase-7 activity, key effectors involved in cellular apoptosis and programmed cell death ³³.

We then used Chou-Talalay's Combination index-Isobologram Theorem to understand the interactive effects of carboplatin and paclitaxel when used in combination treatment ³¹. Evaluation of combination index, which is a direct measure of synergism, antagonism or additive interaction between the drugs revealed more than 25-fold variation across the cell line panel. Similar variation was also observed for DRI values, indicating that drug combinations are also variable based on the degree of refractoriness of a tumor. To understand the genetic determinants underlying this variability, we performed association analysis between the chemo-sensitivity parameters as

phenotype measures and mutations in genes involved in the PK/PD pathway of carboplatin and paclitaxel.

We identified significant association between carboplatin single-agent IC_{50} and genetic variations in drug transporter genes *ABCG2* and *ABCC2*, several mutations in the DNA repair gene *XRCC5* as well as mutations in *PMS2* and *MSH2*. Mutations in the following paclitaxel pathway genes were highly associated with paclitaxel IC_{50} : *TP53*, *EGFR*, and the transporter genes *ABCC1* and *SLC22A7*. On the other hand, several of the top genetic mutations associated with combination treatment belonged to the ATP-binding cassette transporter gene *ABCC1*, and the genes *MSH2* and *MSH6*. Notable among the associations we found between coding mutations and drug cytotoxicity were F150F in *TP53* gene and the S667S in *ABCC1* that were associated with paclitaxel and combination treatment chemosensitivity, respectively. In addition, we also found association between caspase activity and the coding variants R704R (*ERCC6*), R338R and P428P (*MSH6*) and carboplatin single-agent treatment. The genes *ABCC2* and *ABCG2* are involved in carboplatin efflux and multiple studies have shown association of these transporters with survival in ovarian cancer patients ^{9, 34}. Additionally, localization of *ABCC2* in the nuclear membrane was shown associated with cisplatin resistance and clinical outcome in ovarian carcinoma while *ABCC2* overexpression was associated with resistance to cisplatin in melanoma cells ³⁵. Similarly, *ABCG2* overexpression was found associated with poor survival for platinum-containing cancer chemotherapy ³⁶. In contrast, high *ABCC1* expression was associated with favorable survival measures³⁷. Polymorphisms in *ABCC2* and *ABCG2* have been shown associated with response to platinating agents have shown correlated with treatment response and survival in ovarian cancer patients ³⁸. Polymorphisms in DNA repair genes may modulate not only DNA repair capacity, also clinical outcome of DNA damage-inducing anticancer drugs. For example, polymorphisms in *XRCC5*, a gene involved in non-homologous end joining repair process ³⁹, have been shown associated with hematologic toxicity in cancer patients treated with platinum based compound ⁴⁰.

PMS2, *MSH2* and *MSH6* are genes involved in DNA mismatch repair pathway that recognize and repair single stranded DNA damage due to Pt-DNA adducts ⁴¹. We also found a non-synonymous coding variant K1087N in the gene *MAP4* significantly associated with caspase activity in paclitaxel single-agent treatment. Overexpression of *MAP4* have earlier been shown to increase microtubule polymerization and taxane binding with microtubule resulting in increased chemosensitivity ^{42, 43}. Overall, polymorphisms in these genes representing key nodes in the drug metabolism pathway that may have considerable impact on gene function and/or activity thereby affecting drug response.

We also performed differential gene expression of carboplatin and paclitaxel pathway genes to identify changes in gene expression levels associated with drug response. Significant among the differentially expressed carboplatin pathway genes were *ERCC2* (p=0.035 for IC₅₀ and p=0.002 for AUC), *GSTM1* (p=0.010) and *NQO1* (p=0.032). Notably, earlier studies have shown deregulation of *GSTM1* enzyme activity influenced treatment response, 5-year survival and time to progression in various cancers including NSCLC and ovarian cancer patients ⁴⁴. Furthermore, differential expression of *GSTM1* also influences cisplatin sensitivity in cancer patients treated with platinum compound ⁴⁵. Upregulation of the nucleoside excision repair (NER) pathway genes *ERCC1* and *XPB* expression was found significantly associated with higher resistance in platinum-based chemotherapy in advanced EOC ^{46, 47}. NER genes play significant role in the identification and repair of platinum-DNA adducts (1, 2-intrastrand DNA cross-links) ⁴⁸.

Expression of the paclitaxel pathway genes that were significantly associated with cytotoxicity included *FOXO2* (p=0.003), *FOXO1* (p=0.011) and *BIRC5* (p=0.002). Alterations in the apoptotic gene *BIRC5* have been earlier shown associated with overall survival in patients receiving paclitaxel-cisplatin chemotherapy where OS was observed to decrease with increase in number of risk alleles ('bad genotypes') within DNA repair genes including *BRCA1*, *XRCC1* and *BIRC5* ⁴⁹.

Although earlier studies have reported platinum and taxane sensitivity-related changes in DNA and gene expression using cell-based models, the modeling in these studies largely involved datasets that comprised of publicly available EBV-transformed LCL panels from healthy individuals including the Hapmap project ²⁰⁻²² . Therefore, our study is the largest study so far conducted on LCLs modeled using cells derived directly from ovarian cancer patients.

Use of immortalized LCLs directly derived from patient subjects has several advantages over commercially available LCLs since it empowers us to develop *in vitro* drug response models that can be directly correlated with patient clinical responses. Here we successfully demonstrate that the use of candidate-gene based approaches in immortalized LCLs from patient samples can provide reliable genetic markers of drug responses/treatment outcomes that may eventually benefit clinical decision-making.

**CHAPTER 3 RESPONSE TO PLATINUM, TAXANE, AND COMBINATION
THERAPY IN OVARIAN CANCER: IN VITRO PHENOTYPES, INHERITED
VARIATION, AND DISEASE RECURRENCE**

(Published in *Frontiers in Genetics* (2016), 22; 7:37)

3.1. Introduction

Epithelial ovarian cancer (EOC) is the fifth leading cause of cancer death among women in the United States (6% of cancer deaths); in 2015, it is estimated that 14,180 women will die from the disease ¹. The standard treatment for patients with advanced disease is initial debulking surgery followed by carboplatin-paclitaxel combination chemotherapy ². Five-year overall survival remains around 45% ², yet there is a wide inter-patient variation in response. Currently there are few reliable prognostic biomarkers for the classification of patients and treatment response.

Platinating agents, such as carboplatin, interfere with DNA via inter-strand, intra-strand, and DNA-protein crosslinks, thereby causing DNA damage and prevention of cell division and growth, resulting in cell-cycle arrest and apoptosis ³. Although platinum-based drugs are widely used in cancer treatment, many tumors are completely resistant to these drugs, and no clinical response is attained. Major molecular mechanisms underlying this resistance might involve alteration in platinum inactivation or reduced intracellular accumulation by uptake/efflux transporters, increased repair of adducts, increased adduct tolerance or failure of apoptotic pathway. Taxane agents, such as paclitaxel, are commonly used chemotherapeutic drugs often in combination with platinating agents. Taxanes block cell division by binding to α -tubulin, stabilizing the microtubules, thus resulting in cell death ^{4,5}. Although development of taxane resistance is common, the mechanism underlying this resistance is unclear. Molecules implicated in taxane metabolism and disposition include cytochrome P450s and drug transporters (e.g., ABCB1, ABCG2, ABCC1, ABCC2, and SLC01B3) ^{6,7}. Inherited variation in many of the genes encoding these molecules has been assessed for association with clinical outcome with inconsistent results ⁸⁻¹⁰; genome-wide searches to date have also failed to identify variants associated with outcome after correction for genome-wide testing ($p < 5 \times 10^{-8}$).

Patient-derived cell line based model systems represent a novel way to identify genomic predictors of drug response. Although lymphoblastoid cell lines (LCLs) derived from participants in the

international HapMap project have been used to identify genomic predictors of cytotoxic effects of various chemotherapeutic agents ¹¹⁻¹⁶, they are limited as they are not derived from the EOC population but from healthy individuals. Here, we generated LCLs derived from Mayo Clinic EOC patients, conducted *in vitro* cytotoxic studies, and associated *in vitro* drug response phenotypes with germline genotype. Utilizing patient-derived LCLs, as opposed to commercially available LCLs, allows us to screen and directly correlate *in vitro* phenotypes and clinical responses. These genome-wide association scans (GWAS) should contribute to the identification of predictive markers of treatment responses and ultimately improve clinicians' ability to tailor therapy decisions for EOC patients.

3.2. Materials and Methods

3.2.1. Patients, Lymphoblastoid Cell Lines and Cytotoxicity Assays

Prior to initiation of chemotherapy, ovarian cancer patients diagnosed at the Mayo Clinic between 2000 and 2003 provided blood for immediate germline DNA extraction and for the creation of Epstein Barr Virus (EBV)-transformed LCLs. Samples from 74 patients were successfully transformed and subjected to *in vitro* drug testing. All patients provided informed consent, including for passive and active follow-up, using protocols approved by the appropriate Institutional Review Board.

In vitro cellular chemo-sensitivity studies of LCLs were performed in two batches (N=33, N=41) using identical procedures and assays. Cells were maintained in RPMI1640 media supplemented with 2 mM L-glutamine, and 15% fetal bovine serum at 37°C under 5% CO₂. Following 24 hour incubation, LCLs were treated with increasing concentrations of carboplatin and/or paclitaxel (in duplicate). The concentrations of carboplatin were 0, 5, 10, 20, 40, 80 and 128 uM, while for paclitaxel were 0, 4.5, 7.5, 10, 20, 40 and 80 nM when used as single agents. In drug combination experiments, we used half of the doses for each drug in increasing doses (i.e. 2.5 uM carboplatin + 2.25 nM paclitaxel for “dose level 1” and so on). Cell viability 48 hours post-treatment was determined using standard MTT assay^{13, 17, 18}. Caspase3/7 (Promega) apoptosis assays were performed at the same time in parallel plates^{18,19}. A Synergy 3 plate reader (BioTek Instruments) was used to read absorbance (cell viability using MTT) or fluorescence (for caspase3/7 activity) intensities.

Four parameter logistic dose response curves were fit to the *in vitro* drug response measurements (cell survival and caspase3/7 activity assays) for each LCL and treatment (paclitaxel, carboplatin, and combination):

$$Y_i = \alpha + (\beta - \alpha) / (1 + [D_i / \theta]^\phi)$$

where Y_i is the measurement at dose i (D_i), α is the estimated bottom of the curve (i.e., measurement as $D_i \rightarrow \infty$), β is the estimated top of the curve (i.e., measurement as $D_i \rightarrow 0$), ϕ is the slope of the curve, and θ is the inflection point of the curve (i.e., concentration giving a response half way between the top and bottom of the curve). The estimated inflection point of the curve was used as the drug response phenotype. That is, using these dose-response curves, we estimated the relative IC_{50} (effective dose that kills 50% of the cells) for MTT cell viability and the relative EC_{50} (concentration required to induce caspase 3/7 activity by 50%) for caspase3/7 activity²⁰. For simplicity of presentation, we will refer to these quantitative values as the IC_{50} for the MTT cytotoxicity assays and EC_{50} for the caspase 3/7 assays. We then applied rank-based inverse Gaussian transformation (i.e., Van de Waerden rank transformation).

Summaries of the IC_{50} and EC_{50} values for both experimental batches are presented in **Table 3.1**. We observed a high degree of correlation between many of the drug response phenotypes, as expected (**Figure 3.1**).

Table 3.1 Mean (SD) *in vitro* phenotype measures collected on LCLs from 74 EOC patients and comparison across experimental batch.

Phenotype	Treatment	Batch 1 (N=33)	Batch 2 (N=41)	P*
MTT IC_{50}	Paclitaxel	14.22 (8.4)	27.66 (55.1)	0.131
	Carboplatin	55.16 (73.9)	105.52 (138.0)	0.049
	Combination	31.10 (37.6)	47.16 (84.8)	0.281
Caspase 3/7 EC_{50}	Paclitaxel	11.34 (6.7)	9.39 (6.2)	0.865
	Carboplatin	26.00 (13.9)	26.58 (15.3)	0.203
	Combination	10.36 (4.39)	10.28 (5.9)	0.953

*P is result from testing difference in two groups based on a two-sample t-test assuming unequal variances (Welch Test)




Figure 3.1 Scatterplot matrix of drug response phenotypes by experimental batch.

Spearman correlations are presented on the upper panel of each plot (log scale). (A) Batch 1 (N = 33), the highest correlation being between paclitaxel IC₅₀ and combination IC₅₀ (r = 0.81) and caspase EC₅₀ measurements for paclitaxel and combination treatments (r = 0.78). (B) Batch 2 (N = 41), the highest correlation was between paclitaxel and combination IC₅₀ measurements (r = 0.39).

3.2.2. Genotyping and Statistical Methods

Germline DNA was genotyped on the Illumina Infinium 610K array, as previously described^{19, 21, 22}. All samples had genotype call rate >95% and were predicted by STRUCTURE²³ analysis to have greater than 80% European ancestry. SNPs were excluded with call rate < 95%, Hardy-Weinberg Equilibrium $p < 10^{-4}$, or no variation in this set. Using the 1000 Genomes Project²⁴ as reference, imputation was completed with *mach* and *minimac* in a two-step process²⁵, resulting in data on more than 30 million SNPs. Assessment of imputation quality was completed and high quality imputed markers ($r^2 \geq 0.30$ and $MAF \geq 0.01$) were retained (6,243,550 SNPs).

The association of each SNP with *in vitro* drug response phenotypes IC₅₀ or EC₅₀ was evaluated with linear models²⁶ using the expected genotype or “dosage” (i.e., additive or dose-response/trend model). Thus, a negative effect estimate indicates that the carriers of the minor/rare allele have lower IC₅₀ (EC₅₀) values (i.e., were more “sensitive” to treatment). The two *in vitro* experimental batches were analyzed separately followed by meta-analysis was conducted using metal²⁷, with

weights applied for the number of samples in each group. We completed the GWAS analyses for each individual *in vitro* drug response phenotype, as opposed to a combined analysis with all phenotypes model together, due to the difference in mechanism of action between the drugs (i.e., not in the same drug class)²⁸. For annotation of results across gene regions, SNPs were mapped to genes within 2KB using Biofilter (assembly CRCh37.p10, genome build 104.0)²⁹. Pathway analysis used Ingenuity Pathway Analysis (IPA) (Ingenuity® Systems, www.ingenuity.com).

3.3. Results

We examined the relationships of *in vitro* phenotypes with time to recurrence of EOC (40 of 74 patients had recurred or died). Of the 74 patients with LCLs included in this study, 51 had available information on the first two treatments used: 43 patients were treated with paclitaxel / carboplatin, 2 treated with paclitaxel / cisplatin, 3 treated with carboplatin / topotecan, and 3 treated with carboplatin / other less common agent. paclitaxel *in vitro* drug response phenotypes were moderately associated with time to EOC recurrence (HR = 1.90 per unit increase in MTT IC₅₀, p = 0.008; HR = 1.84 per unit increase in caspase 3/7 EC₅₀, p = 0.058) (**Figure 3.2**) (**Table 3.2**).

This suggests that patients whose LCLs demonstrated greater sensitivity to the chemotherapeutics tested had improved outcome; as **Figure 3.2** illustrates, LCLs that were sensitive to paclitaxel (as reflected by having low IC₅₀ and low EC₅₀ values) were from patients with longer time to progression as compared to patients with LCLs with high values. Although based on a small sample size, this provides, for the first time, a link between *in vitro* chemo-sensitivity testing and clinical outcome in EOC.

Results of genome-wide association analyses for each drug response phenotype are presented in **Figure 3.3**. Regions with p < 10⁻⁶ are highlighted and are further displayed in **Figure 3.3**. **Table 3.3** presents the SNPs associated with the drug response phenotype with p < 10⁻⁶. Overall, we found a greater proportion of significant results (e.g., at p < 10⁻⁶) for the combination therapy as compared to the single agent therapies. In particular, we found strong SNP associations with combination therapy in the following gene regions: SLC9A9 (MAF = 0.41, p = 6x10⁻⁷), TIAL1 (MAF = 0.23, p = 7.3x10⁻⁷), ZNF731P (MAF = 0.39, p = 6.6x10⁻⁷), and PCDH20 (MAF = 0.42, p = 8.2x10⁻⁷). None of these regions were found to be moderately associated with single agent carboplatin or paclitaxel IC₅₀ in other pharmacogenomic studies involving commercially available LCLs, such as those of the International HapMap Consortium (personal communications for carboplatin study and paclitaxel study published by Niu et al ³⁰).

Table 3.2 Hazard ratios (p-value) associating *in vitro* drug phenotypes with time to disease recurrence.

	Caspase 3/7 EC₅₀	MTT IC₅₀
Paclitaxel	1.84 (0.058)	1.90 (0.008)
Carboplatin	1.00 (0.991)	1.00 (1.000)
Combination	1.42 (0.376)	1.01 (0.930)

* P-value determined from a likelihood ratio test; Hazard Ratio > 1 indicates worse outcome for subjects with high IC₅₀ ("resistant"); 40 events; bold, p<0.10




Figure 3.2 Kaplan-Meier curves for time to progression and paclitaxel *in vitro* phenotypes. The groupings were defined by 3 quartiles representing low, medium or high values for the phenotype based (A) MTT assay IC₅₀ or (B) Caspase 3/7 EC₅₀ assay. The p-value presented is from a Cox proportional hazards model with the *in vitro* phenotype modeled as a continuous measurement on the log-scale.

Table 3.3 SNPs with $p < 10^{-6}$ association with a drug response phenotype.

Drug	Phenotype	Nearest Gene	SNP	Chr	Position	MAF	Meta-Analysis P	Group 1 (N=33)		Group 2 (N=41)	
								Estimate	P	Estimate	P
Paclitaxel	MTT IC ₅₀	<i>BOD1L1</i>	rs185229225	4	13609129	0.02	2.2E-07	-2.53	8.0E-03	-11.64	4.7E-06
	Caspase 3/7 EC ₅₀	<i>MGC32805</i> / <i>SNCAIP</i> *	rs3842595	5	121778606	0.14	2.6E-07	-1.37	1.7E-03	-1.34	4.1E-05
Carboplatin	MTT IC ₅₀	<i>FRAS1</i>	rs150303591	4	79009309	0.29	5.9E-07	0.86	2.5E-03	1.02	6.3E-05
Combination	MTT IC ₅₀	<i>SLC9A9</i> *	rs201023017	3	143103669	0.41	6.0E-07	0.84	5.2E-04	0.70	3.3E-04
		<i>TIAL1</i>	rs66696671	10	121366953	0.23	7.3E-07	-0.89	2.9E-04	-0.76	6.6E-04
	Caspase 3/7 EC ₅₀	<i>ZNF731P</i>	rs12025262	1	247356732	0.39	6.6E-07	-0.84	2.5E-04	-0.71	6.9E-04
		<i>PCDH20</i>	rs10674174	13	61892075	0.42	8.2E-07	-0.73	5.3E-03	-0.86	3.7E-05

*Nearest gene within 2000 gene pairs.

For regions with multiple SNPs with $p < 10^{-6}$, the most significant SNP is presented. A negative estimate indicates that carriers of the minor allele had, on average, lower IC₅₀ or EC₅₀ ("sensitive") while a positive estimate indicates that carriers of the minor allele had, on average, higher IC₅₀ or EC₅₀ ("resistant").




Figure 3.3 Manhattan plots of the single SNP meta-analysis genome-wide association results based the six *in vitro* drug response measures among ovarian cancer patient LCLs.

(A) carboplatin MTT; (B) paclitaxel MTT; (C) Combination of carboplatin and paclitaxel MTT; (D) carboplatin Caspase3/7; (E) paclitaxel Caspase3/7; and (F) Combination of carboplatin and paclitaxel Caspase3/7. Blue line indicates $p\text{-value} = 0.00001$; Red line indicates $p\text{-value}=5 \times 10^{-8}$. Highlighted regions (circled) are displayed in **Figure 3**.




Figure 3.4 Locus Zoom plots for regions with $p < 10^{-6}$.

Note: BOD1L1 region not presented as only one SNP in region with $p < 0.001$.

We evaluated potential overlap of loci associated with both phenotypes for a given drug. Only one SNP was found to be associated with a $p < 10^{-4}$ with same direction of the effect for MTT IC_{50} and caspase3/7 EC_{50} values for paclitaxel, carboplatin, or combination treatment. An intronic SNP rs35067965 in COLEC12 (chromosome 18, bp 455396) was associated with response to paclitaxel (MTT IC_{50} $p = 2.2 \times 10^{-5}$, caspase 3/7 EC_{50} $p = 3.8 \times 10^{-5}$) (**Table 3.4**).

We also examined overlap of associations at the level of genes, considering SNPs within 20kb. This showed consistency of IC₅₀ and EC₅₀ results for paclitaxel response and COLEC12 and revealed similar IC₅₀ and EC₅₀ associations for carboplatin response in the gene regions of CTIF and CDH4. As presented in **Table 3.5**, additional regions showed joint associations with response to multiple drugs, including variants in protein coding regions of BRE, EML6, CTNNA2, LRP1B, EYS, NKAIN2, ANTXRL, COL13A1, and MTCL1 (SNPs in gene regions with $p < 0.0001$).

Because of the suggested association between *in vitro* paclitaxel MTT IC₅₀ response and time to EOC recurrence (**Table 3.2**), we also examined recurrence association with SNPs rs185229225 (intronic BOD1L1) rs35067965 (intronic COLEC12) and rs1525599 (intronic LRP1B) which were associated with paclitaxel MTT IC₅₀ (**Table 3.3.**, **Table 3.4.** and **Table 3.5.**, respectively). However, none of these SNPs were associated with time to recurrence with a nominal p-value < 0.05 (data not shown).

Table 3.4 Gene regions with SNPs associated with both phenotypes for a given drug.

Drug	Gene	Phenotype	SNP*	Chr	Position	P	Direction [†]
Paclitaxel	COLEC12	MTT IC ₅₀	rs35067965	18	455396	2.2E-05	--
		Caspase 3/7 EC ₅₀	rs35067965	18	455396	3.8E-05	--
Carboplatin	CTIF	MTT IC ₅₀	rs8091660	18	46087936	8.9E-06	--
		Caspase 3/7 EC ₅₀	rs113867814	18	46259604	1.2E-05	--
	CDH4	MTT IC ₅₀	rs2748151	20	60133486	4.7E-05	++
		Caspase 3/7 EC ₅₀	rs113594423	20	60379048	2.4E-05	++

*Presenting most significant SNP in the region for the giving drug/phenotype. SNP within ± 20 KB of the listed gene. †A positive estimate indicates that carriers of the minor/variant allele had, on average, higher IC₅₀ or EC₅₀ ("resistant").

Table 3.5 Gene regions with SNPs associated with multiple drugs for any phenotype (p < 0.0001).

Gene	Drug	Phenotype	SNP*	Chr	Position	P	Direction [†]
<i>BRE</i>	Carboplatin	Caspase 3/7 EC ₅₀	rs5830067	2	28537890	1.70E-05	++
	Combination	Caspase 3/7 EC ₅₀	rs7572644	2	28320033	5.80E-06	--
<i>EML6</i>	Paclitaxel	Caspase 3/7 EC ₅₀	rs75314082	2	55087315	7.90E-05	--
	Combination	MTT IC ₅₀	rs17046344	2	55023600	4.90E-05	++
<i>LINC01122</i>	Paclitaxel	Caspase 3/7 EC ₅₀	rs72817940	2	58998563	6.40E-05	++
	Carboplatin	Caspase 3/7 EC ₅₀	rs4233974	2	59295043	2.60E-05	--
<i>CTNNA2</i>	Carboplatin	MTT IC ₅₀	rs17261321	2	80197843	3.60E-05	++
	Combination	MTT IC ₅₀	rs6719499	2	80193386	6.00E-05	--
<i>LRP1B</i>	Paclitaxel	MTT IC ₅₀	rs1525599	2	141778702	8.60E-05	++
	Combination	Caspase 3/7 EC ₅₀	rs13020675	2	142212928	6.20E-05	--
<i>EYS</i>	Paclitaxel	Caspase 3/7 EC ₅₀	rs201083182	6	65736914	2.30E-06	--
	Combination	Caspase 3/7 EC ₅₀	rs2064701	6	65676556	3.60E-05	++
<i>NKAIN2</i>	Paclitaxel	Caspase 3/7 EC ₅₀	rs550987	6	124905510	4.10E-05	--
	Combination	Caspase 3/7 EC ₅₀	rs670616	6	124885773	7.80E-05	++
<i>C7orf65</i>	Carboplatin	Caspase 3/7 EC ₅₀	rs10230114	7	47705506	2.40E-05	++
	Combination	Caspase 3/7 EC ₅₀	rs11771997	7	47712495	2.40E-05	++
<i>ANTXRL</i>	Paclitaxel	Caspase 3/7 EC ₅₀	rs12572446	10	47665906	4.30E-05	++
	Combination	Caspase 3/7 EC ₅₀	rs10906942	10	47670851	4.90E-05	++
<i>COL13A1</i>	Carboplatin	Caspase 3/7 EC ₅₀	rs10999018	10	71654602	2.40E-05	++
	Combination	Caspase 3/7 EC ₅₀	rs77535242	10	71652985	3.50E-05	++
<i>TMEM132D</i>	Paclitaxel	Caspase 3/7 EC ₅₀	rs77438645	12	130304313	7.60E-05	--
	Carboplatin	Caspase 3/7 EC ₅₀	rs1451904	12	130166947	6.50E-05	++
<i>MTCL1</i>	Carboplatin	Caspase 3/7 EC ₅₀	rs690089	18	8845223	7.80E-05	--
	Combination	Caspase 3/7 EC ₅₀	rs35765215	18	8839469	6.00E-05	--

*Presenting most significant SNP in the region for the giving drug/phenotype. SNP within ±20 KB of the listed gene. †A negative estimate indicates that carriers of the minor/variant allele had, on average, lower IC₅₀ or EC₅₀ ("sensitive") while a positive estimate indicates that carriers of the minor/variant allele had, on average, higher IC₅₀ or EC₅₀ ("resistant").

3.4. Discussion and Conclusions

In this proof of concept study, we explored use of LCLs derived from EOC patients followed for clinical response as a model for discovery of pharmacogenomics markers. LCLs were treated with varying concentration of the chemotherapeutics agents (carboplatin and paclitaxel and, uniquely, their combination) that were used for the treatment and cellular chemo-sensitivity was determined by measuring cell viability and activation of caspase activity (as a marker of apoptosis) post drug treatment. Genome-wide association studies were performed to identify inherited markers associated with these measures of *in vitro* chemo-sensitivity (i.e., MTT IC₅₀ and caspase 3/7 EC₅₀ values) and the relationships between *in vitro* measures and clinical outcome was explored.

Although the sample size was small limiting the power of the study, some of the implicated biologically interesting genes are worthy of discussion. Pathway analysis of genes with SNPs showing association with one of the drug response phenotypes (at $p < 10^{-6}$), both phenotypes for a given drug (at $p < 10^{-4}$), or multiple drugs for any phenotype (at $p < 10^{-4}$) found enrichment in genes related to malignant solid tumor and epithelial cancers (**Figure 3.5A**). Among the top canonical pathways represented by these genes were “Epithelial Adherens Junction Signaling”, “Sertoli Cell Junction Signaling”, and “Endometrial Cancer Signaling” (**Figure 3.5B**). In addition, genes such as CTNNA2 and CDH4, both tumor suppressor genes with role in cell adhesion were found to be associated with chemo-sensitivity in carboplatin alone or combination treatments. SNPs in CTNNA2- catenin (cadherin-associated protein) alpha2, a structural constituent of cytoskeleton and cadherin binding was associated with *in vitro* cytotoxicity to carboplatin alone as well as in combination with paclitaxel. CTNNA2 has been shown to be frequently mutated in laryngeal carcinomas with mutations predictive of poor prognosis ³¹. Additionally, SNPs within CTNNA2 have recently been implicated in breast cancer ³² and its role in tumor progression and metastasis has been suggested for multiple cancers ³³. Variants in CTNNA2 have also been implicated in schizophrenia ³⁴ and alcohol addiction ³⁵. CTNNA2 SNPs associated with carboplatin and

paclitaxel MTT IC₅₀ were both intronic and present functional relevance of these is not known. CDH4, codes for cadherin, and has been implicated in nasopharyngeal carcinoma ³⁶ and aberrant methylation of CDH4 promoter has been colorectal and gastric cancer ³⁷. Our results identified two intronic SNPs (rs2748151 and rs113594423) that were associated with carboplatin resistant as measured by cell death (IC₅₀) and apoptosis (caspase 3/7 EC₅₀). Variants in PCDH20, another member of cadherin family, were also found to be associated *in vitro* drug response. PCDH20 codes for protocadherin20 and functions as a tumor suppressor by interacting with Wnt/b-catenin signaling^{38, 39}.

Another gene with role in cell adhesion identified in our study was FRAS1, which encodes for an extracellular matrix protein and is involved in the regulation of epidermal-basement membrane adhesion and organogenesis during development. Inherited mutations in FRAS1, and FREM2, have been associated with development of Fraser syndrome. FRAS1 has also been implicated in ERK signaling and influence migration and invasion of lung cancer cell line by influencing FAK signaling ⁴⁰, suggesting its role in tumorigenesis and metastasis of lung cancer. Although the genes described above are involved cell adhesion/cell migration, the functional significance of the intronic SNPs identified in this study is not known and would require further investigation.

Two intronic variants within BRE were found to be associated with caspase 3/7 levels for carboplatin and combination therapy (indel rs5830067 and rs7572664). BRE encodes for Brain and reproductive Organ-Expressed (TNFRSF1A modulator) and is a component of BRCA1-A DNA damage repair complex that recognizes Lys 62linked ubiquitinated H2A and H2Ax at DNA lesions, resulting in recruitment of BRCA1-BARD1 to double strand DNA breaks ⁴¹. BRE expression has been shown to be predictive of disease free survival in non-familial breast cancer patients ⁴² and recent studies show its involvement in both intrinsic and extrinsic apoptotic pathways by influencing XIAP ⁴³. Variation within EML6, which is involved assembly dynamics of microtubules, was found to be associated with platinum-sensitivity which was of interest since

paclitaxel's mechanism of action involved disruption of microtubules; however no evidence exists in the literature on functional relevance of these particular SNPs within EML6. No other genes involved in microtubule protein were identified with respect to paclitaxel chemo-sensitivity.

Lastly, two intronic variants within LRP1B (low density lipoprotein related protein 1B) were associated with paclitaxel and combination therapy drug response phenotypes. LRP1B is a tumor suppressor with decreased expression in several primary cancers and is among ten most significantly deleted genes across 3312 cancer samples ⁴⁴⁻⁴⁷. In renal cell cancer, down-regulation of LRP1B has been shown to regulate cell motility and actin cytoskeleton reorganization ⁴⁸. Germline SNPs/ haplotype in LRP1B have been associated with aging without cognitive decline ⁴⁹; however, associations of germline SNPs with incidence/progression of cancer and pharmacogenomics have yet to be reported.




Figure 3.5 Pathway analysis of genes with SNPs that demonstrated significant association with LCL chemosensitivity with carboplatin or paclitaxel alone or in combination.

A) Significant number of genes mapped to malignant solid tumor or epithelial cancer; B) Top canonical pathways and interaction network among the pathways represented by significantly associated genes.

In summary, using a patient-derived cell-based model system to generate several *in vitro* drug response phenotypes on a clinically followed set of EOC cases we have identified genetic loci associated with response to platinum-taxane therapies. Overall our results identified germ-line SNPs in multiple cell adhesion molecules and several tumor suppressor genes (PCDH20, LRP1B, CDH4, and CTNNA2). However, none of the most associated SNPs were reported by Huang et al¹⁵ or associated with mRNA gene expression in lymphoblastoid cell lines (<http://www.ncbi.nlm.nih.gov/projects/gap/eql/index.cgi>). Further studies are needed to determine if these SNPs are truly associated with drug response or if they represent false-positive

findings. Similar to other studies comparing *in vitro* chemo-sensitivity with clinical outcomes^{15, 50}, our findings suggest that *in vitro* response to paclitaxel correlates with time to disease recurrence indicating that this model may have utility in several types of future studies. On possible explanation for the observation that paclitaxel correlates with recurrence and not carboplatin may be the fact that the majority of EOC patients eventually develop platinum resistant tumors and the main factor related to future response maybe attributed to response to taxane therapy. Further research is needed to understand the mechanism by which genomic loci impact clinical response in ovarian cancer patients to the most common regimen used in the treatment of ovarian cancer following surgery.

**CHAPTER 4 GENETIC VARIATION IN PLATINATING AGENT AND
TAXANE PATHWAY GENES AS PREDICTORS OF OUTCOME AND TOXICITY
IN ADVANCED NON-SMALL-CELL LUNG CANCER**

(Published in *Pharmacogenomics* (2014), 15(12):1565-74)

4.1 Introduction

Lung carcinoma is the most common malignancy worldwide, and the leading cause of cancer deaths. According to the most recent estimates, the global incidence of lung cancer is more than 1.6 million cases/year, resulting in more than 1.3 million deaths/year (>18% of all cancer deaths)¹. In the USA, the incidence of lung cancer was estimated to be 228,000 cases in 2013, leading to nearly 160,000 deaths (27.5% of all cancer deaths)². Histologically, the majority (approximately 80%) of lung cancers are non-small-cell carcinomas (NSCLCs). The 5-year relative survival of patients with advanced NSCLC remains dismal at approximately 4%². Indeed, the median survival of unselected patients diagnosed with metastatic NSCLC is only 4 months³. Platinating agent-based combination chemotherapy is the standard treatment for patients with advanced NSCLC who have adequate organ function and performance status, and whose tumor does not have a driver mutation amenable to treatment with a specific inhibitor. However, combination chemotherapy achieves objective response in only approximately 26% of patients, improves survival modestly, and is associated with diverse side effects⁴. Interpatient variation in response and toxicity, which cannot be predicted for individual patients, precludes the selection and tailoring of chemotherapy that might improve outcomes and minimize adverse events. Although clinical factors including age, performance status and stage influence the likelihood of benefit from and tolerability of chemotherapy, the genetic profile of individual patients may contribute significantly to the marked variation in response and toxicity.

Adverse effects associated with platinating agent-based combination chemotherapy include gastrointestinal toxicity, myelosuppression, nephrotoxicity, peripheral neuropathy and ototoxicity⁵. Platinating drugs such as cisplatin and carboplatin act by forming platinum–DNA adducts, which lead to cell cycle arrest and apoptosis. Several genes are involved in the carboplatin drug pathway (platinating agents pathway PharmGKB: **(Figure 4.1)** Intracellular levels of carboplatin are regulated by drug transporters SLC31A1 (CTR1), ABCC2 (MRP2), ATP7A and ATP7B⁶, as well

as drug-metabolizing enzymes including MPO, SOD1, GSTM1, NQO1, GSTP1 and MT, which are implicated in the development of cellular resistance to these drugs ⁷⁻¹⁰. Genes of pharmacodynamic significance include HMGB1, which is involved in recognition and cellular response to platinum–DNA, adducts and DNA repair genes including mismatch repair genes MSH6 and MLH1, and nucleotide excision repair genes XRCC1, ERCC1, ERCC2 and XPA ^{11, 12}. Genetic variation in these genes of importance to pharmacokinetic and pharmacodynamics pathways of platinating agents may thus contribute to interpatient variation in response and tolerability ¹³.




Figure 4.1 Platinum pharmacokinetics/pharmacodynamics pathway

Representation of the candidate genes involved in the metabolism of platinum containing drugs ¹⁴.

Taxanes are routinely given in combination with platinating agents, genes involved in efflux (ABCC1, ABCC2, ABCG2 and ABCB1) and metabolism of taxane (CYP3A4 and CYP2C8) are critical for its therapeutic efficacy (taxane pathway: PharmGKB) (**Figure 4.2**). Although the role of genetic variation in taxane response is unclear at this time, some studies have found no significant association between SNPs and treatment outcome ¹⁵, whereas others have found

significant associations between ABCB1 SNPs and response to paclitaxel ¹⁶, and gastro intestinal toxicity in patients treated with taxane and platinum combination therapy ¹⁷. In the present study, we evaluated SNPs in genes of relevance to the pharmacokinetic/pharmacodynamics pathways of platinating agents and taxanes in patients diagnosed with advanced NSCLC and treated primarily with carboplatin-based doublet chemotherapy, and determined the association of individual SNPs with outcomes and toxicity. We identified specific SNPs that were predictive of progression-free survival (PFS) and multiple adverse effects, after adjusting for known clinical prognostic factors in multivariate models. Results of our proof-of-concept study provides evidence that in real world clinical settings the association of genetics with clinical outcome is evaluable and although validation in larger cohorts is required, genetic information can be utilized to develop to more effective treatment strategies.




Figure 4.2 Taxane pharmacokinetics pathway

Representation of the genes involved in the metabolism and transport of paclitaxel and docetaxel, and the downstream effects of the drugs ¹⁸.

4.2 Materials and methods

4.2.1 Patients cohort

A total of 635 patients diagnosed with stage IIIB or IV NSCLC between December 1998 and December 2008 were identified by the institutional tumor registry (**Figure 4.3**). Of these, 90 patients met the following eligibility criteria: histological or cytological confirmation of NSCLC, presence of measurable disease, performance status Eastern Cooperative Oncology Group (ECOG) 0–2, no neoadjuvant or concurrent radiation therapy or surgery, no second malignancies, availability of adequate diagnostic tumor tissue, treatment with first-line platinating-agent based doublet chemotherapy and complete follow-up at the Minneapolis VA Health Care System (MN, USA). The study was approved by the institutional human subjects committee.




Figure 4.3 Patient selection and Study design.

NSCLC: Non-Small-Cell lung carcinoma, MAF: Minor allele frequency, XRT: Radiotherapy

4.2.2 Treatment schedules, dose & toxicity assessments

Of the 90 patients identified, 87 received carboplatin, two received cisplatin, and two received cisplatin and carboplatin as first-line chemotherapy. Patients were evaluated every 2–3 cycles using the Response Evaluation Criteria in Solid Tumors (RECIST) 1.1. We assessed the following hematologic toxicities: anemia, neutropenia, febrile neutropenia and thrombo-cytopenia. Nonhematologic toxicities assessed included nausea, vomiting, diarrhea, mucositis, sensory and motor neuropathy, renal dysfunction (increase in creatinine), and liver dysfunction (abnormal liver function tests). Grading of toxicities was performed using the National Cancer Institute Common Toxicity Criteria Version 4.0. Clinical evaluation and blood tests were performed prior to the first cycle of chemotherapy and before each subsequent cycle. Baseline CT scans were obtained before the start of treatment, and repeated every 2–3 cycles to evaluate treatment response. Patients had open access to the oncology clinic for reporting any side effects or concerns during treatment.

4.2.3 SNP analysis:

4.2.4 DNA isolation

Genomic DNA was isolated from formalin-fixed paraffin-embedded specimens (n=86 NSCLC patients) using Qiagen DNA isolation from formalin-fixed paraffin-embedded tissues kit (Qiagen, Valencia, CA, USA) as per manufacturer's instructions.

4.2.5 Genotyping

Genes of importance to platinating agents and paclitaxel as well as potentially significant SNPs within them were selected from PharmGKB¹⁹ and literature search from PubMed. Overall, 63 SNPs within 29 genes in 86 subjects with detailed clinical information were genotyped using the Sequenome platform at the University of Minnesota Genomics Center (MN, USA). Only include SNPs with MAF>0.1 (**Figure: 4.4, Table 4.1**).

Six samples were removed from further statistical analyses owing to low call rates (<90%). There was no significant difference in age, performance status and survival between patients excluded versus included in study ($p>0.05$). Quality control of the SNP data resulted in one SNP being removed with a call rate <90%, with no SNPs removed owing to deviations from Hardy–Weinberg equilibrium.




Figure 4.4 SNP selection process.

Table 4.1 Summary of SNPs genotyped in the NSCLC patients.

SNP	Chr	Position	Gene	Major Allele	Minor Allele	MAF	Call rate	HWE
rs1695	11	67352689	GSTP1/missense	A	G	0.336	0.95	0.8
rs7483	1	110279701	GSTM3	G	A	0.342	1	0.802
rs11615	19	45923653	ERCC1/cds-synon	T	C	0.386	0.9875	1
rs13181	19	45854919	KLC3/nearGene-3	T	G	0.431	1	0.821
rs25487	19	44055726	XRCC1/missense	G	A	0.27	0.95	0.151
rs25489	19	44056412	XRCC1/missense	G	A	0.05	1	0.168
rs35592	16	16141823	ABCC1/intron	T	C	0.201	0.9625	0.722
rs119774	16	16086833	ABCC1/intron	G	A	0.069	1	1
rs246240	16	16119024	ABCC1/intron	A	G	0.131	1	0.614
rs316019	6	160670282	SLC22A2/missense	G	T	0.125	1	0.094
rs447978	3	120491086	GTF2E1/intron	A	G	0.312	1	0.117
rs520227	1	190569312		C	G	0.307	0.8875	0.001
rs744154	16	14015081	ERCC4/intron	C	G	0.312	1	1
rs851974	6	151971687	ESR (7 kb 5')	T	C	0.436	1	0.357
rs1042522	17	7579472	TP53/nearGene-5,TP53/missense	G	C	0.291	0.9875	0.584
rs1045385	6	10398116	TFAP2A/UTR-3	A	NA	NA	0.9875	NA
rs1045642	7	87138645	ABCB1	T	C	0.456	1	0.366
rs1047768	13	103504517	ERCC5/cds-synon	C	T	0.342	0.9875	0.624
rs1058930	10	96818119	CYP2C8/missense	C	G	0.069	1	0.308
rs1128503	7	87179601	ABCB1	C	T	0.456	1	0.652
rs1138272	11	67353579	GSTP1/missense	C	T	0.094	1	0.515
rs1347851	12	90566978		A	G	0.244	1	0.542
rs1625895	17	7578115	TP53/intron	G	A	0.144	1	0.193
rs1799735	1	110280254	GSTM3	-	TCC		1	NA
rs1799793	19	45867259	ERCC2/missense	G	A	0.383	0.9625	0.81
rs1800566	16	69745145	NQO1/missense	C	T	0.222	0.9875	0.512
rs1801244	13	52544805	ATP7B/missense	C	G	0.481	0.975	0.82
rs1801247	13	52520471	ATP7B/cds-synon	G	A	0.051	0.9875	1
rs1944118	11	111352032	BTG4/intron	G	A	0.35	1	0.466
rs2031920	10	135339845	CYP2E1/nearGene-5	C	T	0.013	0.95	1
rs2032582*	7	87160618	ABCB1				0.9875	NA
rs2228001	3	14187449	XPC/missense	A	C	0.344	1	0.806
rs2229109	7	87179809	ABCB1	G	A	0.038	1	1
rs2230671**	16	16228242	ABCC1/cds-synon	A	G	NA	0.9875	NA
rs2231142	4	89052323	ABCG2/missense	C	A	0.081	1	0.411

rs2235015	7	87199564	ABCB1	G	T	0.131	1	0.614
rs2238476	16	16213872	ABCC1/intron	C	T	0.062	1	0.259
rs2273697	10	101563815	ABCC2/missense	G	A	0.2	0.9375	1
rs2798389	14	83051150		A	G	0.262	1	0.775
rs3212948	19	45924362	ERCC1/intron	C	G	0.369	1	0.811
rs3212986	19	45912736	CD3EAP/missense,ERCC1/UTR-3	G	T	0.269	0.975	1
rs3740066	10	101604207	ABCC2/cds-synon	G	A	0.308	0.975	1
rs3957357	6	52668687	GSTA1/nearGene-5	G	A	0.394	1	0.35
rs4128317	2	29659854	ALK/intron	A	C	0.419	1	0.818
rs4148396	10	101591944	ABCC2/intron	C	T	0.342	0.95	0.8
rs6863960	5	114995781		G	A	0.469	0.9375	1
rs7255865	19	21804968		A	G	0.412	1	1
rs8187710	10	101611294	ABCC2/missense	G	A	0.056	0.9875	1
rs9282564	7	87229440	ABCB1	A	G	0.088	0.975	0.463
rs9312960	5	204082	CCDC127/5'UTR	C	A	0.294	1	0.282
rs10158985	1	226050609	TMEM63A/intron	C	A	0.247	0.9625	1
rs11572080	10	96827030	CYP2C8/missense	G	A	0.106	1	0.589
rs12188653	5	101383803		G	C	0.257	0.95	0.005
rs13120400	4	89033527	ABCG2/intron	T	C	0.288	1	0.272
rs17098912	14	100104329		G	A	0.206	1	1
rs17222723	10	101595996	ABCC2/missense	T	A	0.051	0.9875	1
rs17718902	11	17784707	KCNC1/intron	A	G	0.388	1	0.642
rs17731538	4	89055379	ABCG2/intron	G	A	0.2	1	0.725
rs28364006	16	16228249	ABCC1/missense	A	G	0.012	1	0.006
rs28730837	17	56355397	MPO/missense	C	T	0.025	1	1
rs45511401	16	16173232	ABCC1/missense	G	T	0.014	1	1
rs72552784	7	87145914	ABCB1	C	NA	NA	1	NA

(MAF= minor allele frequency; HWE= Hardy Weinberg equilibrium)

4.2.6 Statistical methods

All analyses were restricted to subjects with self-reported race of white or 'unknown'. Association of each clinical feature with clinical outcomes of PFS and overall survival (OS) was assessed using Cox-proportion hazards models. Associations of each SNP genotype with clinical outcomes were examined using Cox-proportion hazards models, where the SNP genotyped was coded under a dominant, codominant or additive genetic model. Models were fit with and without adjustment for clinical features associated with clinical outcomes. For analysis of PFS, we included covariates of

stage and performance status. For OS, we included performance status as a covariate. For analyses with clinical features, stage was treated as a categorical variable and performance status was treated as a continuous variable. Age at diagnosis was categorized into four levels based on the 25, 50 and 75% percentiles of the distribution. All statistical analyses were completed using R (version 2.15.3). Adjustment for multiple testing was carried out using Bonferroni correction.

4.3 Results

4.3.1 Demographic features and clinical characteristics

Demographic and clinical features of the 90 evaluable patients at diagnosis are shown in **Table 4.2**.

The median age was 66 years (range: 45–81). All the patients were males, a reflection of the patient population at a VA Medical Center. The majority (93%) of patients was white; 81% of patients had stage IV disease. Adeno-carcinoma was the most common histological subtype (42% of patients) of NSCLC, followed by squamous cell (20%) and large cell (16%) carcinoma. The median performance status (ECOG scale) was 1 (range: 0–2).

Table 4.2 Patient demographics and clinical characteristics (n = 90).

	n(%)	p-value (PFS)	p-value (OS)
Age at diagnosis (years)			
Median	66	0.95	0.25
Range	45-81		
Race			
White	84 (93.3)		
African-American	1 (1.1)		
American Indian/Alaska Native	2 (2.2)		
Unknown	3 (3.3)		
Stage			
IIIB	17 (19)	0.02	0.48
IV	73 (81)		
Histology			
Adenocarcinoma	40 (44.4)		
Adenocarcinoma, mucinous	2 (2.2)		
Squamous cell	18 (20)		
Large cell	12 (13.3)		
Large cell neuroendocrine	2 (2.2)		
Non-small-cell carcinoma, NOS	16 (17.8)		
Brain metastasis	12 (13.3)		
Performance status (ECOG)			
Median	1	0.03	0.02
Range	0-2		

ECOG: Eastern Cooperative Oncology Group; NOS: Not otherwise specified; OS: Overall survival; PFS: Progression-free survival.

4.3.2 Treatment, response and outcomes

Chemotherapy regimens administered, including the number of cycles and average dose intensities of each agent, are shown in **Table 4.3**. The overall response rate was 19%, with a median duration of response of 182 days, median PFS of 148 days and median OS of 288 days. No patient achieved a complete response.

Hematological and non-hematological toxicities selected for further analyses are shown in **Table 4.4**. No patients developed treatment-related abnormalities in liver function tests, or experienced CTCAE grade 5 (fatal) toxicity.

Table 4.3 Chemotherapy regimens, dose intensity, response and survival.

Regimen	Number of patients	Number of cycles, mean (range)	Dose intensity (mg/m ² /week) [†] , mean ± SE		Response rate, n (%)	Median duration of response (days)	Median PFS (days)	Median OS (days)
			Platinum [‡]	Second drug				
C + T	77	3.3 (1–6)	1.8 ± 0.3	60.0 ± 10.0	14 (18)	163	148	292
C + G	9	3.2 (1–5)	1.4 ± 0.4	455.1 ± 169.1	3 (33)	185	211	465
C + E	2	3 (1–3)	1.9 ± 0.5	84.9 ± 19.0	0 (0)	0	NE	NE
Cis + E	2	1.5 (1–2)	21.8 ± 4.2	75.0 ± 0.0	0 (0)	0	NE	NE
Total	90	3.2 (1–6)	–	–	17 (19)	182	148	288

None of the patients had a complete response.

The median PFS and OS were calculated from the Kaplan–Meier estimates.

[†]The target dose intensities for each agent were: C AUC 2/week; Cis, 27 mg/m²/week; T, 67 mg/m²/week; G, 667 mg/m²/week and E, 100 mg/m²/week (with C) and 80 mg/m²/week (with Cis).

[‡]Dose of C is expressed as AUC.

AUC: Area under the concentration–time curve; C: carboplatin; Cis: Cisplatin; E: Etoposide; G: Gemcitabine; NE: Not evaluable; OS: Overall survival; PFS: Progression-free survival; SE: Standard error; T: paclitaxel.

Table 4.4 Hematological and non-hematological toxicities.

Toxicity and grade†	Chemotherapy regimen			
	C + T, n (%)	C + G, n (%)	C + E, n (%)	Cis + E, n (%)
Neutropenia				
1–2	10 (12)	5 (56)	0 (0)	0 (0)
3–4	19 (25)	2 (22)	1 (50)	0 (0)
Febrile neutropenia				
1–2	0 (0)	0 (0)	0 (0)	0 (0)
3–4	3 (4)	0 (0)	0 (0)	1 (50)
Anemia				
1–2	72 (94)	7 (78)	2 (100)	0 (0)
3–4	2 (3)	2 (22)	0 (0)	1 (50)
Thrombocytopenia				
1–2	18 (23)	4 (44)	0 (0)	1 (50)
3–4	3 (4)	2 (22)	0 (0)	1 (50)
Nausea				
1–2	28 (36)	3 (33)	0 (0)	2 (100)
3–4	5 (7)	0 (0)	0 (0)	0 (0)
Vomiting				
1–2	4 (5)	1 (11)	0 (0)	0 (0)
3–4	4 (5)	0 (0)	0 (0)	0 (0)
Diarrhea				
1–2	15 (20)	2 (22)	0 (0)	0 (0)
3–4	7 (10)	0 (0)	1 (50)	1 (50)
Mucositis				
1–2	1 (1)	1 (11)	0 (0)	0 (0)
3–4	0 (0)	0 (0)	0 (0)	0 (0)
Sensory neuropathy				
1–2	24 (31)	3 (33)	0 (0)	0 (0)
3–4	5 (7)	0 (0)	0 (0)	0 (0)
Motor neuropathy				
1–2	0 (0)	0 (0)	0 (0)	0 (0)
3–4	2 (3)	0 (0)	0 (0)	0 (0)
Increased creatinine				
1–2	2 (3)	1 (11)	0 (0)	0 (0)

†Toxicities were graded according to Common Terminology Criteria for Adverse Events (CTCAE) Version 4.0. None of the patients developed abnormal liver function tests or experienced grade 5 toxicity. The number of patients who received each chemotherapy regimen was: C + T = 77, C + G = 9, C + E = 2 and Cis + E = 2. C: carboplatin; Cis: Cisplatin; E: Etoposide; G: Gemcitabine; T: paclitaxel.

4.3.3 Pharmacogenetics

We first examined the association of known clinical prognostic factors (covariates) with clinical outcomes. Stage ($p=0.02$) and performance status ($p=0.03$) were predictive of PFS, and performance status ($p=0.02$) was associated with OS. Therefore, we adjusted for these covariates when analyzing the association between SNPs and outcomes. In the analysis of toxicity, we adjusted for the total number of cycles received, since the cumulative dose of chemotherapeutic agents received would be expected to increase the risk of toxicities such as peripheral neuropathy. For genetic association testing, 86 patients had both clinical and genotype data. Of these 86 patients we included patients with white ancestry ($n=80$) for further analysis. Of 58 SNPs with a minimum allele frequency of >0.05 , one SNP was excluded due to call rate of <0.90 . Although number of chemotherapy cycles were associated with OS, we decided not to adjust for them as genetic factors might be influencing number of cycle and if we adjust for cycles, we might mask some of the effects due to patient genetics (as it might be ‘surrogate’ for length of survival[depends on time; changes over time]). PFS was associated with SNPs in drug efflux transporters ABCC1 (intronic SNPs: rs246240 and rs2238476) and ABCB1 (coding synonymous SNP, rs1128503); a Quinone reductase NQO1 (missense

SNP, rs1800566); TMEM63A (rs10158985); and other genes such as KCNC1 (rs17718902) and CCDC127 (rs9312960; summarized in **Table 4.5**). SNPs within ABCB1 (rs1045642; $p=0.04$), ABCC1 (rs2238476; $p=0.03$), ABCG2 (rs17731538; $p=0.04$) and TMEM63A (rs10158985; $p=0.03$) were also associated with OS.

We also evaluated association of SNPs with major hematological and non-hematological toxicities. Toxicities were not associated with the initial performances status ($p>0.05$) but were associated with age and number of treatment cycles. After adjusting for age and number of treatment cycles, we found associations of ABCB1SNP with thrombocytopenia (rs2235015; $p=0.04$), and ABCG2

(rs2231142; $p=0.045$) and ATP7B (rs1801244; $p=0.027$) SNPs with nausea (**Table 4.6**), and ABCG2SNP (rs13120400, $p=0.027$) with sensory neuropathy, respectively. Interestingly, SNPs in the DNA repair pathway genes ERCC4 (rs744154; $p=0.04$) and XPC (rs2228001; $p=0.045$) demonstrated associations with neutropenia and sensory neuropathy, respectively (**Table 4.6**). Additionally, we observed SNPs in TP53 (rs165895; $p=0.02$), CYP2C8 (rs11572080; $p=0.017$) and CCDC127 (rs9312960; $p=0.019$) to be associated with diarrhea. rs447978, an intronic SNP in transcription factor GTF2E1, was associated with both neutropenia ($p=0.024$) and nausea ($p=0.027$). After adjustment for multiple testing (Bonferroni) the above indicated associations were not significant and the results need to be validated in a larger study.

Table 4.5 Association of SNPs with progression-free survival after adjusting for covariates (performance status and stage).

SNP	Gene	Drug	Chr	Major Allele	Minor Allele	MAF	P value PFS Additive Model (Dominant Model)	Hazard Ratio PFS Additive Model (Dominant Model)	95% CI, additive model (dominant model)
rs1800566 (missense)	NQO	platinating agents	16	C	T	0.222	0.02 (0.037)	1.615 (1.689)	0.089 to 0.870 (0.040 to 1.009)
rs10158985 (intron)	TMEM63A [#]	platinating agents	1	C	A	0.247	0.032 (0.025)	0.655 (0.592)	-0.821 to -0.025
rs246240 (intron)	ABCC1	Taxane	16	A	G	0.131	0.037 (0.072)	0.640 (0.617)	0.409 to 1.001
rs2238476 (intron)	ABCC1	Taxane	16	C	T	0.062	0.045 (0.037)	1.972 (2.32)	-0.894 to 0.001
rs1128503 (synonymous)	ABCB1	Taxane	7	C	T	0.456	0.066 (0.019)	0.730 (0.541)	-0.652 to 0.022
rs17718902 (intron)	KCNC1 [#]	platinating agents	11	A	G	0.388	0.069 (0.035)	1.353 (1.717)	-0.017 to 0.623
rs9312960	CCDC1278 [#]	platinating agents	5				0.059 (0.042)	1.378 (1.620)	-0.003 to 0.645

Data are presented as: values from additive model, three group analysis (values from dominant model, variant carrier vs noncarrier).
MAF: Minimum allele frequency.

Table 4.6 Association of SNPs with toxicity after adjusting for covariates (age as categorical variable and total number of cycles).

Toxicity	Gene	SNP	P value, log Additive Model (Dominant Model)	Odds Ratio Additive Model (Dominant Model)	95% CI, log-additive model (dominant model)
Nausea	ABCG2	rs2231142_CA (missense)	0.045(0.045)	3.938 (4.052)	0.033 to 2.708 (0.031 to 2.768)
	GTF2E1	rs447978_AG (intron)	0.024 (0.004)	0.409 (0.223)	-1.672 to -0.117 (-2.520 to -0.488)
	ATP7B	rs1801244_CG (missense)	0.078 (0.027)	1.977 (4.626)	-0.073 to 1.384(0.178 to 2.885)
Neutropenia	GTF2E1#	rs447978 (intron)	0.027 (0.039)	0.444 (0.345)	-1.531 to -0.093(-2.01 to -0.053)
	TMEM63A	rs10158985	0.031 (0.09)	2.557 (2.307)	0.035 to 1.52(0.308 to 2.307)
	ERCC4	rs744154 (intron)	0.04 (0.01)	2.176 (3.697)	0.087 to 1.79(-0.158 to 1.83)
Sensory Neuropathy	#	rs1347851_AG	0.016 (0.023)	0.235 (0.225)	-2.625 to -0.27(-2.773 to -0.208)
	ABCG2	rs13120400 (Intron)	0.027 (0.055)	0.271 (0.295)	-2.467 to -0.146(-2.465 to 0.024)
	KLC3#	rs13181_TG	0.073 (0.021)	2.27 (5.893)	-0.076 to 1.715 (0.273 to 3.274)
	XPC	rs2228001_AC (missense)	0.096 (0.045)	0.456 (0.296)	-1.71 to 0.14(-2.406 to -0.03)
Diarrhea	CCDC127	rs9312960_CA (5'UTR)	0.019 (0.09)	2.429 (2.434)	0.149 to 1.626 (-0.138 to 1.918)
	CYP2C8	rs11572080_GA (missense)	0.017 (0.017)	4.396 (4.396)	0.26 to 2.702 (0.26 to 2.702)
	TP53	rs1625895_GA (intron)	0.02 (0.023)	3.004 (3.633)	0.174 to 2.026 (0.179 to 2.401)
Thrombocytopenia	ABCB1	rs2235015_GT	0.074 (0.04)	2.605 (3.467)	-0.093 to 2.004 (0.058 to 2.428)
	#	rs17098912_GA	0.072 (0.027)	2.212 (3.115)	-0.071 to 1.659 (0.131 to 2.142)

Data are presented as: values from additive model, three group analysis (values from dominant model, variant carrier vs non-carrier).

†SNPs not in a gene, these were selected from genome-wide association study.

4.4 Discussion

This comprehensive analysis identified pharmacogenomics variants of clinical significance within key genes (n=29) in the platinating agents pathway and taxane pathway in NSCLC patients treated with a combination of carboplatin and paclitaxel. We found significant associations of drug transporter (ABCC1 and ABCB1) SNPs with PFS. Although we did not observe significant association of ABCB1 rs1045642 with outcome/toxicity, ABCB1 SNPs rs1128503 (occurring in partial linkage disequilibrium with rs1045642) and rs2235015 were associated with PFS and thrombocytopenia, respectively. The functional consequence of the three most commonly studied SNPs (rs1128503, rs1045642 and rs2032582 also occurring in linkage disequilibrium) in ABCB1 is still not completely understood. Associations of these SNPs with mRNA/protein expression have been shown in some studies but not all the studies. rs1045642, a synonymous SNP (3435C>T; Ile1145Ile) has also been indicated to influence protein conformation and substrate specificity ²⁰. Previous studies have identified association of ABCB1 SNPs (especially rs1045642) with survival in osteosarcoma patients treated with cisplatin containing chemotherapy ²¹. In esophageal cancer patients treated with platinating agents, presence of the T allele for rs1045642 is associated with significantly longer survival and reduced risk of recurrence ²². Together, these findings suggest that drug transporter SNPs that might influence outcomes in several malignancies by altering intracellular drug concentrations. NQO1 belongs to the Quinone dehydrogenase family of proteins and has been implicated in metabolizing platinum agents. A missense SNP rs1800566 (NQO1*2: Pro187Ser) is in the active site of the enzyme and has been associated with reduced NQO1 activity. The NQO1 isoform with Ser (NQO1*2) undergoes rapid degradation ^{23,24}. A recent study reported association of the NQO1*2 SNP with shorter OS in NSCLC patients treated with adjuvant radiation therapy with or without platinating agent based chemotherapy ²⁵. Our results are in concordance with these previous observations, with presence of NQO1*2 associated with poor PFS (**Table 4.5**). Toxicity due to platinating agent and taxane-based chemotherapy remains a major challenge

faced by clinicians and patients. There are no well-established biomarkers that can predict such toxicities in individual patients. In our efforts to identify SNPs predictive of major hematological and non-hematological toxicities, we observed that a missense SNP in CYP2C8, a drug-metabolizing enzyme involved in the metabolism of taxanes, was significantly associated with higher incidence of diarrhea (OR: 4.396; $p=0.017$). Previous results in breast cancer patients receiving paclitaxel have shown significant association of CYP2C8*3 (which denotes two highly linked SNPs rs11572080 and rs10509681) with better response and a trend towards greater risk of peripheral neuropathy²⁶. CYP2C8*3 is associated with lower paclitaxel- α hydroxylation activity, and hence carriers of this SNP have lower clearance of paclitaxel, which can contribute to better response but also greater risk of toxicity to normal cells^{27, 28}. In addition to drug-metabolizing enzymes, several SNPs in drug transporters (ABCG2 and ABCB1) were associated with nausea and thrombocytopenia, respectively. These drug transporters have been implicated in efflux of either platinating agents or taxanes. The ABCG2 missense SNP rs2231142 results in a Gln141Lys change; which has been associated with lower expression²⁹ as well as with reduced drug efflux capacity of ABCG2³⁰. Cell lines that are resistant to cisplatin have been shown to express higher levels of the copper transporters ATP7A and ATP7B⁵. ATP7A has been implicated in sequestering both carboplatin and cisplatin in vesicles. Since SNPs in these transporters have not been evaluated for potential clinical impact, we performed an exploratory evaluation of SNPs in ATP7A and ATP7B and found that rs1801244 (valine to leucine change) in ATP7B is associated with nausea. The functional effect of the valine to leucine change resulting from this SNP remains to be determined. Since platinating agents such as cisplatin and carboplatin form platinum–DNA adducts that are repaired by the nucleotide excision repair pathway, interpatient variation in DNA repair mechanism due to presence of SNPs in DNA damage/repair pathway genes may influence treatment outcome. Although none of the SNPs in the DNA repair pathway genes were associated with PFS or OS, significant correlations were observed with toxicities. A 3'-UTR SNP (rs3212986)

and a coding SNP in ERCC1(rs11615 ; Asn118Asn) have been associated with reduced ERCC1 mRNA/protein-expression levels, and lower ERCC1 levels have been associated with better outcomes in NSCLC patients receiving platinating agent based chemotherapy ^{31, 32}. Nevertheless, while multiple studies have evaluated these two ERCC1SNPs, results have been quite variable and remain inconclusive. A recent meta-analysis of 39 previously published studies on these two ERCC1SNPs in lung cancer patients demonstrated that the rs11615 SNP might be a potential biomarker for risk of developing lung cancer as well as a prognostic marker in NSCLC patients treated with platinating agents ³³. In our dataset we did not observe significant association of these SNPs with toxicity of survival. We did observe a marginally increased risk of neutropenia with ERCC4intronic SNP and reduced risk of sensory neuropathy with XPC missense SNP rs2228001 (C>A; Gln939Lys). Though not extensively studied, rs2228001 has been shown to have a trend towards higher risk of ototoxicity in osteosarcoma patients treated with cisplatin ³⁴. Finally, in addition to candidate genes in drug pathways; we tested selected genes that have been identified by genome-wide association study (GWAS). GWAS analysis in colorectal cancer patients treated with 5-fluorouracil with or without oxaliplatin identified SNPs in KCNC1 (rs17718902), GTF2E1 (rs447978), CCDC127(rs9312960) or TMEM63A (rs10158985) to be associated with nausea, vomiting and drug-induced neuropathy ³⁵. In our study, an intronic SNP (rs447978) in the transcription factor GTF2E1 was associated with reduced risk of nausea and neutropenia and an intronic SNP in TMEM63 (rs10158985) was associated with neutropenia. Furthermore, SNPs in KCL3and CCDC127were associated with increased risk of sensory neuropathy and diarrhea, respectively. Although the functional relevance of these genes/SNPs is not known, future studies on functional characterization as well as validation of these SNPs are needed to confirm the results of the GWAS analysis.

4.5 Conclusion

In conclusion, we identified SNPs within key candidate drug pathway genes that are independently predictive of PFS and/or major hematological and non-hematological toxicities in patients with advanced NSCLC treated with platinating agent based chemotherapy. There is a paucity of predictive markers that can be used to guide clinical decisions for such patients. Our results confirm some previously reported associations, but more importantly identify several new candidates that warrant testing in prospective studies, that could contribute towards the development of personalized medicine. Indeed, a recent Phase II trial demonstrated that response rates are higher when chemotherapy is selected based on SNPs in ERCC1 and RR M1 in patients with advanced NSCLC ³⁶, suggesting that incorporation of pharmacogenomics biomarkers into clinical decision making has considerable potential for improving therapeutic outcomes.

4.6 Summary

Lung carcinoma is the most common malignancy and the leading cause of cancer deaths worldwide. The genetic makeup of individual patients in addition to other clinical factors can influence the likelihood of benefit from and tolerability of chemotherapy. We evaluated SNPs in candidate genes of relevance to platinating agents and paclitaxel for association with survival and toxicity in non-small-cell lung cancer patients. After adjusting for covariates, SNPs in drug transporters and NQO1 were associated with progression-free survival and SNPs in drug transporters as well as in the in the DNA repair pathway genes ERCC4 and XPC were associated with toxicity. Comprehensive evaluation of the genetic variants in conjunction with known prognostic factors may help optimize therapeutic decisions to maximize benefit and minimize toxicity in non-small-cell lung cancer patients.

CHAPTER 5 SUMMARY

Platinum drugs are currently used as standard-of-care chemotherapy for various cancers including solid tumors like lung cancer, the most common malignancy in the world, and ovarian cancer, the most lethal gynecological malignancy in the United States. Platinating agents including carboplatin form DNA helix-distorting DNA adducts that lead to strand breakage activating DNA repair mechanisms that ultimately resulting in apoptotic death of cancer cells. Taxanes are another widely used class of chemotherapy drug to treat various cancers including ovarian and lung cancers, primarily in combination with platinating agents. paclitaxel, a taxane, is a mitotic inhibitor that stabilizes microtubules and interferes with cell division leading to induction of apoptosis.

However, despite considerable improvements in therapeutic approaches, the overall response rate of carboplatin/paclitaxel based combination chemotherapy is far from desirable. Drug resistance, interpatient variation in response and toxicity are major causes of concern. Furthermore, a large bulk of the cancer patients who do respond eventually undergo relapse after first line based chemotherapy. Inter-patient variability in treatment response is therefore a commonly observed phenomenon in this drug resistance. However, the complete mechanism governing this spectrum of sensitivity to chemotherapeutic agents still remains largely unclear.

Genetic factors like differential expression and/or activity of genes involved in drug metabolism pathway and the presence of single nucleotide polymorphisms may have an impact on treatment outcome and toxicity in cancer patients being treated with Platinum/Paclitaxel combination. Genetic variation in these key pharmacokinetic and pharmacodynamics pathway genes may therefore contribute to interpatient variation in response and toxicity.

Our approach was to use cell line models derived from epithelial ovarian cancer (EOC) patients to identify genetic polymorphisms and differential gene expression associated with chemo-sensitivity and response to carboplatin and paclitaxel single agent and combination treatments followed by the evaluation of these predictive pharmacogenomics markers in epithelial ovarian cancer and NSCLC patients undergoing Carboplatin/Paclitaxel combination chemotherapy. For the purpose of cell-

based modeling of ovarian cancer, we generated immortalized Lymphoblastoid cell lines (LCLs) from human lymphocytes from Epithelial ovarian cancer (EOC) patients through the infection of Epstein-Barr virus (EBV). Then, we performed a comprehensive profiling in vitro chemosensitivity phenotypes in our panel of ~100 Epstein-Barr virus (EBV) transformed EOC LCLs following treatment with carboplatin, paclitaxel as single agent and in combination. We demonstrated extensive inter-individual variation in drug response as was evident from the drug IC_{50} and area under survival curve (AUC) values. Fold changes between the lowest (most-sensitive) and highest (most-resistant) IC_{50} and AUC values were 8.34 & 3.84, respectively for carboplatin, and 7.72 & 5.65, respectively for paclitaxel. Our drug cytotoxicity values corroborated with the caspase activity assay results demonstrating that drug cytotoxicity indeed reflected alterations in cellular apoptosis and programmed cell death. Furthermore, using Chou-Talalay's Combination index-Isobologram Theorem we also observed, based on Combination index values, wide inter-individual variations in effects of carboplatin and paclitaxel when used in combination treatment. The wide range of dose reduction index values indicated variability in impact of the dose of one drug on another in combination treatment based on the cell line used for the study.

To understand the pharmacogenomics of the inter-individual variation in chemo-sensitivity parameters, we performed a pathway based identification of polymorphisms within the PK/PD pathway of carboplatin and paclitaxel as well as analysis of gene expression of these pathway genes in the EOC LCLs using TaqMan Low Density Arrays (TLDA), followed by genotype-phenotype association analysis.

Our analysis revealed significant association between drug chemosensitivity measures and genetic variations in several key candidate genes involved in carboplatin and paclitaxel pathway.

Genetic variations in drug transporter genes ABCG2 and ABCC2, several mutations in the DNA repair gene ERCC6 and XRCC5 as well as mutations in PMS2 and MSH2 and MSH6 were found associated with carboplatin cytotoxicity whereas. Mutations in microtubule associated protein

MAP4, the transporter genes ABCC1 and SLC22A7, as well as TP53 and EGFR were associated with paclitaxel response. Top among the mutations associated with combination treatment were SNPs within the genes ABCC1, MSH2 and MSH6. Remarkably, we found a number of coding variants associated with drug response. These include K1087N (MAP4; paclitaxel), F150F (TP53; paclitaxel) and S667S (ABCC1; combination), R704R (ERCC6; carboplatin), R338R and P428P (MSH6; carboplatin). Our gene expression analysis revealed changes in expression levels of the following pathway genes were associated with drug response phenotypes: ERCC2 (carboplatin), GSTM1 (carboplatin) and NQO1 (carboplatin); FOXC2 (paclitaxel), FOXL1 (paclitaxel) and BIRC5 (paclitaxel). NER genes have been shown to play significant roles in the identification and repair of platinum-DNA adducts. Earlier studies have shown upregulation of expression of the NER pathway genes ERCC1 and XPD was significantly associated with higher resistance in platinum-based chemotherapy in advanced EOC. Our results thus corroborate with this earlier finding and identifies the NER genes associated with platinum drug resistance.

Subsequently, we performed a comprehensive genome-wide association scans (GWAS) of germline genotype of these patient-derived LCLs to discover predictive pharmacogenomics markers of treatment response within the human genome. At $p < 10^{-4}$, we found genes related to malignant solid tumor and epithelial cancers that were significantly associated with our drug chemosensitivity measures. Notably, the top canonical pathways represented by these genes included Epithelial Adherens Junction Signaling, Sertoli Cell Junction Signaling and Endometrial Cancer Signaling, which seems reasonable as the cells were derived from ovarian cancer patients. We also found mutations in several relevant genes associated with drug response including the tumor suppressor genes CTNNA2, CDH4 and LRP1B; FRAS1, a gene supposedly involved in tumorigenesis and metastasis; and the DNA repair gene BRE.

Subsequently, to understand the impact of genetic variation within the pharmacokinetic/pharmacodynamics pathways of platinating agents and taxanes in NSCLC, we

performed a genotype-phenotype correlation analysis on in advanced NSCLC patients treated primarily with carboplatin/paclitaxel-based combination chemotherapy. Our results showed a number of pathway SNPs associated with treatment outcomes and toxicity including progression-free survival (PFS) and multiple adverse effects, following adjusting for clinical prognostic factors in multivariate models. Our results showed SNPs in key pathway genes were associated with treatment outcome (PFS) including ABCC1, ABCB1 (rs1045642/ Ile1145Ile, rs1128503, rs2235015), NQO1 (NQO1*2: Pro187Ser). Whereas, SNPs in the following genes were also associated with hematological and/or non-hematological toxicities: CYP2C8 (diarrhea); ABCG2 and ABCB1 (thrombocytopenia); (ATP7B rs1801244; nausea); ERCC4 (neutropenia); XPC missense SNP rs2228001 (C>A; Gln939Lys). Among other SNPs associated with toxicities included rs447978 (GTF2E1; nausea and neutropenia, rs10158985 (TMEM63; neutropenia), as well as SNPs in KCL3 (neuropathy) and CCDC127 (diarrhea). The nonsynonymous coding SNP in NQO1, rs1800566 (NQO1*2: Pro187Ser), is in the active site of the NQO1 enzyme and has been associated with reduced NQO1 activity since the NQO1 isoform with Ser (NQO1*2) undergoes rapid degradation.

Most remarkably, the expression of NQO1 was also shown to be associated with our *in vitro* profiling of carboplatin chemosensitivity in EBV-transformed ovarian cancer LCLs. Similarly, SNPs in a number of other pathway genes that were found relevant with regard to the treatment response and/or toxicities resulting from treatment with platinating agents/ taxane-containing chemotherapy regimens in NSCLC patients were also shown associated with *in vitro* chemosensitivity in our EOC LCL panel-based model system of drug resistance. For example, the ABCC1 intronic SNPs rs246240 and rs2238476 were associated with PFS in NSCLC patients after adjusting for covariates, while ABCC1 SNPs were also found significantly correlated with *in vitro* paclitaxel and combination cytotoxicity as well as caspase activity following combination treatment. The ABCG2 missense SNP rs2231142 was associated with nausea, and the intronic SNP

rs13120400 with Sensory neuropathy in NSCLC; ABCG2 SNPs were also associated with *in vitro* carboplatin cytotoxicity. In addition, SNPs in CYP2C8 and TP53 were associated with diarrhea in NSCLC patients as well as response to paclitaxel *in vitro*. Also, ERCC2 SNPs were associated with caspase 3/7 activity following carboplatin treatment *in vitro*, while the ERCC2/KLC3 missense SNP rs13181 that was found significantly associated with sensory neuropathy in NSCLC patients after adjusting for covariates.

Thus, using immortalized LCLs directly derived from patient subjects we could successfully develop *in vitro* drug response models that could be directly correlated with patient clinical responses. Furthermore, we have effectively demonstrated by using candidate-gene based and genome-wide approaches in these immortalized LCLs from patient samples we could identify key pharmacogenomics changes associated with drug response, treatment outcome and toxicity. Finally, we identified genetic variation within the pharmacokinetic/pharmacodynamics pathways of platinating agents and taxanes in real world clinical settings in advanced NSCLC patients undergoing carboplatin-based chemotherapy.

However, although we found a number of genetic variations significantly associated with treatment outcome and toxicities, the functional relevance of most of these SNPs vis-à-vis activity of target genes is still largely unknown. Therefore, for a comprehensive understanding of the role of these variations, we propose future studies focusing on the characterization of the significant SNPs using functional genetic/genomic approaches.

Therefore, results from our study provide an actionable panel of reliable genetic biomarkers of carboplatin and paclitaxel single-agent and combination treatment response as clinically applicable signatures of drug response and toxicity which has the potential to be used in precision medicine approaches following further validation to develop pharmacogenomics-guided treatment approaches to achieve maximum efficacy and minimum toxicity.

REFERENCES

Chapter 1

1. Mescher AL. Junqueira's basic histology: text and atlas. McGraw-hill; 2013.
2. Yancik R. Ovarian cancer. Age contrasts in incidence, histology, disease stage at diagnosis, and mortality. *Cancer*. 1993;71(2 Suppl):517-523.
3. American Cancer Society. American Cancer Society: Cancer Facts and Figures 2012. 2012.
4. Siegel RL, Miller KD, Jemal A. Cancer statistics, 2015. *CA Cancer J Clin*. 2015;65(1):5-29.
5. Chen VW, Ruiz B, Killeen JL, Cote TR, Wu XC, Correa CN. Pathology and classification of ovarian tumors. *Cancer*. 2003;97(10 Suppl):2631-2642.
6. Marsh S. Pharmacogenomics of taxane/platinum therapy in ovarian cancer. *Int J Gynecol Cancer*. 2009;19 Suppl 2:S30-4.
7. Lynch HT, Watson P, Lynch JF, Conway TA, Fili M. Hereditary ovarian cancer. Heterogeneity in age at onset. *Cancer*. 1993;71(2 Suppl):573-581.
8. Pennington KP and Swisher EM. Hereditary ovarian cancer: beyond the usual suspects. *Gynecol Oncol*. 2012;124(2):347-353.
9. Risch HA, McLaughlin JR, Cole DE, et al. Prevalence and penetrance of germline BRCA1 and BRCA2 mutations in a population series of 649 women with ovarian cancer. *Am J Hum Genet*. 2001;68(3):700-710.
10. Pal T, Permuth-Wey J, Betts JA, et al. BRCA1 and BRCA2 mutations account for a large proportion of ovarian carcinoma cases. *Cancer*. 2005;104(12):2807-2816.
11. Hilton JL, Geisler JP, Rathe JA, Hattermann-Zogg MA, DeYoung B, Buller RE. Inactivation of BRCA1 and BRCA2 in ovarian cancer. *J Natl Cancer Inst*. 2002;94(18):1396-1406.
12. Rebbeck TR, Lynch HT, Neuhausen SL, et al. Prophylactic oophorectomy in carriers of BRCA1 or BRCA2 mutations. *N Engl J Med*. 2002;346(21):1616-1622.
13. Beral V, Million Women Study Collaborators, Bull D, Green J, Reeves G. Ovarian cancer and hormone replacement therapy in the Million Women Study. *Lancet*. 2007;369(9574):1703-1710.
14. Zhou B, Sun Q, Cong R, et al. Hormone replacement therapy and ovarian cancer risk: a meta-analysis. *Gynecol Oncol*. 2008;108(3):641-651.
15. Amos CI and Struwing JP. Genetic epidemiology of epithelial ovarian cancer. *Cancer*. 1993;71(2 Suppl):566-572.
16. Adami HO, Hsieh CC, Lambe M, et al. Parity, age at first childbirth, and risk of ovarian cancer. *Lancet*. 1994;344(8932):1250-1254.
17. Risch HA, Marrett LD, Howe GR. Parity, contraception, infertility, and the risk of epithelial ovarian cancer. *Am J Epidemiol*. 1994;140(7):585-597.
18. Calle EE, Rodriguez C, Walker-Thurmond K, Thun MJ. Overweight, obesity, and mortality from cancer in a prospectively studied cohort of U.S. adults. *N Engl J Med*. 2003;348(17):1625-1638.
19. Engeland A, Tretli S, Bjorge T. Height, body mass index, and ovarian cancer: a follow-up of 1.1 million Norwegian women. *J Natl Cancer Inst*. 2003;95(16):1244-1248.
20. Canchola AJ, Chang ET, Bernstein L, et al. Body size and the risk of ovarian cancer by hormone therapy use in the California Teachers Study cohort. *Cancer Causes Control*. 2010;21(12):2241-2248.
21. Schouten LJ, Goldbohm RA, van den Brandt PA. Height, weight, weight change, and ovarian cancer risk in the Netherlands cohort study on diet and cancer. *Am J Epidemiol*. 2003;157(5):424-433.
22. Hankinson SE, Colditz GA, Hunter DJ, Spencer TL, Rosner B, Stampfer MJ. A quantitative assessment of oral contraceptive use and risk of ovarian cancer. *Obstet Gynecol*. 1992;80(4):708-714.
23. Collaborative Group on Epidemiological Studies of Ovarian Cancer, Beral V, Doll R, Hermon C, Peto R, Reeves G. Ovarian cancer and oral contraceptives: collaborative reanalysis of data from 45 epidemiological studies including 23,257 women with ovarian cancer and 87,303 controls. *Lancet*. 2008;371(9609):303-314.
24. Havrilesky LJ, Moorman PG, Lowery WJ, et al. Oral contraceptive pills as primary prevention for ovarian cancer: a systematic review and meta-analysis. *Obstet Gynecol*. 2013;122(1):139-147.
25. Wilailak S, Vipupinyo C, Suraseranivong V, et al. Depot medroxyprogesterone acetate and epithelial ovarian cancer: a multicentre case-control study. *BJOG*. 2012;119(6):672-677.
26. Ness RB, Dodge RC, Edwards RP, Baker JA, Moysich KB. Contraception methods, beyond oral contraceptives and tubal ligation, and risk of ovarian cancer. *Ann Epidemiol*. 2011;21(3):188-196.
27. Cibula D, Widschwendter M, Majek O, Dusek L. Tubal ligation and the risk of ovarian cancer: review and meta-analysis. *Hum Reprod Update*. 2011;17(1):55-67.

28. Sieh W, Salvador S, McGuire V, et al. Tubal ligation and risk of ovarian cancer subtypes: a pooled analysis of case-control studies. *Int J Epidemiol.* 2013;42(2):579-589.
29. Luan NN, Wu QJ, Gong TT, Vogtmann E, Wang YL, Lin B. Breastfeeding and ovarian cancer risk: a meta-analysis of epidemiologic studies. *Am J Clin Nutr.* 2013;98(4):1020-1031.
30. Huncharek M, Geschwind JF, Kupelnick B. Perineal application of cosmetic talc and risk of invasive epithelial ovarian cancer: a meta-analysis of 11,933 subjects from sixteen observational studies. *Anticancer Res.* 2003;23(2C):1955-1960.
31. Terry KL, Karageorgi S, Shvetsov YB, et al. Genital powder use and risk of ovarian cancer: a pooled analysis of 8,525 cases and 9,859 controls. *Cancer Prev Res (Phila).* 2013;6(8):811-821.
32. Heintz AP, Odicino F, Maisonneuve P, et al. Carcinoma of the ovary. *Int J Gynaecol Obstet.* 2003;83 Suppl 1:135-166.
33. Omura GA, Brady MF, Homesley HD, et al. Long-term follow-up and prognostic factor analysis in advanced ovarian carcinoma: the Gynecologic Oncology Group experience. *J Clin Oncol.* 1991;9(7):1138-1150.
34. Hoskins WJ, Bundy BN, Thigpen JT, Omura GA. The influence of cytoreductive surgery on recurrence-free interval and survival in small-volume stage III epithelial ovarian cancer: a Gynecologic Oncology Group study. *Gynecol Oncol.* 1992;47(2):159-166.
35. Thigpen T, Brady MF, Omura GA, et al. Age as a prognostic factor in ovarian carcinoma. The Gynecologic Oncology Group experience. *Cancer.* 1993;71(2 Suppl):606-614.
36. Vencken PM, Kriege M, Hoogwerf D, et al. Chemosensitivity and outcome of BRCA1- and BRCA2-associated ovarian cancer patients after first-line chemotherapy compared with sporadic ovarian cancer patients. *Ann Oncol.* 2011;22(6):1346-1352.
37. Mogensen O. Prognostic value of CA 125 in advanced ovarian cancer. *Gynecol Oncol.* 1992;44(3):207-212.
38. Alberts DS, Liu PY, Hannigan EV, et al. Intraperitoneal cisplatin plus intravenous cyclophosphamide versus intravenous cisplatin plus intravenous cyclophosphamide for stage III ovarian cancer. *N Engl J Med.* 1996;335(26):1950-1955.
39. Markman M, Bundy BN, Alberts DS, et al. Phase III trial of standard-dose intravenous cisplatin plus paclitaxel versus moderately high-dose carboplatin followed by intravenous paclitaxel and intraperitoneal cisplatin in small-volume stage III ovarian carcinoma: an intergroup study of the Gynecologic Oncology Group, Southwestern Oncology Group, and Eastern Cooperative Oncology Group. *J Clin Oncol.* 2001;19(4):1001-1007.
40. Armstrong DK, Bundy B, Wenzel L, et al. Intraperitoneal cisplatin and paclitaxel in ovarian cancer. *N Engl J Med.* 2006;354(1):34-43.
41. Elit L, Oliver TK, Covens A, et al. Intraperitoneal chemotherapy in the first-line treatment of women with stage III epithelial ovarian cancer: a systematic review with metaanalyses. *Cancer.* 2007;109(4):692-702.
42. Katsumata N, Yasuda M, Isonishi S, et al. Long-term results of dose-dense paclitaxel and carboplatin versus conventional paclitaxel and carboplatin for treatment of advanced epithelial ovarian, fallopian tube, or primary peritoneal cancer (JGOG 3016): a randomised, controlled, open-label trial. *Lancet Oncol.* 2013;14(10):1020-1026.
43. du Bois A, Luck HJ, Meier W, et al. A randomized clinical trial of cisplatin/paclitaxel versus carboplatin/paclitaxel as first-line treatment of ovarian cancer. *J Natl Cancer Inst.* 2003;95(17):1320-1329.
44. Neijt JP, Engelholm SA, Tuxen MK, et al. Exploratory phase III study of paclitaxel and cisplatin versus paclitaxel and carboplatin in advanced ovarian cancer. *J Clin Oncol.* 2000;18(17):3084-3092.
45. Ozols RF, Bundy BN, Greer BE, et al. Phase III trial of carboplatin and paclitaxel compared with cisplatin and paclitaxel in patients with optimally resected stage III ovarian cancer: a Gynecologic Oncology Group study. *J Clin Oncol.* 2003;21(17):3194-3200.
46. Vasey PA, Jayson GC, Gordon A, et al. Phase III randomized trial of docetaxel-carboplatin versus paclitaxel-carboplatin as first-line chemotherapy for ovarian carcinoma. *J Natl Cancer Inst.* 2004;96(22):1682-1691.
47. Kristensen GB, Vergote I, Stuart G, et al. First-line treatment of ovarian cancer FIGO stages IIb-IV with paclitaxel/epirubicin/carboplatin versus paclitaxel/carboplatin. *Int J Gynecol Cancer.* 2003;13 Suppl 2:172-177.

48. Bookman MA, Brady MF, McGuire WP, et al. Evaluation of new platinum-based treatment regimens in advanced-stage ovarian cancer: a Phase III Trial of the Gynecologic Cancer Intergroup. *J Clin Oncol.* 2009;27(9):1419-1425.
49. Rustin GJ, van der Burg ME, Griffin CL, et al. Early versus delayed treatment of relapsed ovarian cancer (MRC OV05/EORTC 55955): a randomised trial. *Lancet.* 2010;376(9747):1155-1163.
50. Muggia FM. Overview of carboplatin: replacing, complementing, and extending the therapeutic horizons of cisplatin. *Semin Oncol.* 1989;16(2 Suppl 5):7-13.
51. Monk BJ, Herzog TJ, Kaye SB, et al. Trabectedin plus pegylated liposomal Doxorubicin in recurrent ovarian cancer. *J Clin Oncol.* 2010;28(19):3107-3114.
52. Parmar MK, Ledermann JA, Colombo N, et al. Paclitaxel plus platinum-based chemotherapy versus conventional platinum-based chemotherapy in women with relapsed ovarian cancer: the ICON4/AGO-OVAR-2.2 trial. *Lancet.* 2003;361(9375):2099-2106.
53. Pujade-Lauraine E, Wagner U, Aavall-Lundqvist E, et al. Pegylated liposomal Doxorubicin and Carboplatin compared with Paclitaxel and Carboplatin for patients with platinum-sensitive ovarian cancer in late relapse. *J Clin Oncol.* 2010;28(20):3323-3329.
54. Cantu MG, Buda A, Parma G, et al. Randomized controlled trial of single-agent paclitaxel versus cyclophosphamide, doxorubicin, and cisplatin in patients with recurrent ovarian cancer who responded to first-line platinum-based regimens. *J Clin Oncol.* 2002;20(5):1232-1237.
55. Pfisterer J, Plante M, Vergote I, et al. Gemcitabine plus carboplatin compared with carboplatin in patients with platinum-sensitive recurrent ovarian cancer: an intergroup trial of the AGO-OVAR, the NCIC CTG, and the EORTC GCG. *J Clin Oncol.* 2006;24(29):4699-4707.
56. Wagner U, Marth C, Largillier R, et al. Final overall survival results of phase III GCIG CALYPSO trial of pegylated liposomal doxorubicin and carboplatin vs paclitaxel and carboplatin in platinum-sensitive ovarian cancer patients. *Br J Cancer.* 2012;107(4):588-591.
57. Gordon AN, Tonda M, Sun S, Rackoff W, Doxil Study 30-49 Investigators. Long-term survival advantage for women treated with pegylated liposomal doxorubicin compared with topotecan in a phase 3 randomized study of recurrent and refractory epithelial ovarian cancer. *Gynecol Oncol.* 2004;95(1):1-8.
58. Piccart MJ, Green JA, Lacave AJ, et al. Oxaliplatin or paclitaxel in patients with platinum-pretreated advanced ovarian cancer: A randomized phase II study of the European Organization for Research and Treatment of Cancer Gynecology Group. *J Clin Oncol.* 2000;18(6):1193-1202.
59. Kohn EC, Sarosy G, Bicher A, et al. Dose-intense taxol: high response rate in patients with platinum-resistant recurrent ovarian cancer. *J Natl Cancer Inst.* 1994;86(1):18-24.
60. McGuire WP, Rowinsky EK, Rosenshein NB, et al. Taxol: a unique antineoplastic agent with significant activity in advanced ovarian epithelial neoplasms. *Ann Intern Med.* 1989;111(4):273-279.
61. Einzig AI, Wiernik PH, Sasloff J, Runowicz CD, Goldberg GL. Phase II study and long-term follow-up of patients treated with taxol for advanced ovarian adenocarcinoma. *J Clin Oncol.* 1992;10(11):1748-1753.
62. Thigpen JT, Blessing JA, Ball H, Hummel SJ, Barrett RJ. Phase II trial of paclitaxel in patients with progressive ovarian carcinoma after platinum-based chemotherapy: a Gynecologic Oncology Group study. *J Clin Oncol.* 1994;12(9):1748-1753.
63. Trimble EL, Adams JD, Vena D, et al. Paclitaxel for platinum-refractory ovarian cancer: results from the first 1,000 patients registered to National Cancer Institute Treatment Referral Center 9103. *J Clin Oncol.* 1993;11(12):2405-2410.
64. ten Bokkel Huinink W, Gore M, Carmichael J, et al. Topotecan versus paclitaxel for the treatment of recurrent epithelial ovarian cancer. *J Clin Oncol.* 1997;15(6):2183-2193.
65. Gordon AN, Fleagle JT, Guthrie D, Parkin DE, Gore ME, Lacave AJ. Recurrent epithelial ovarian carcinoma: a randomized phase III study of pegylated liposomal doxorubicin versus topotecan. *J Clin Oncol.* 2001;19(14):3312-3322.
66. du Bois A, Luck HJ, Meier W, et al. Carboplatin/paclitaxel versus cisplatin/paclitaxel as first-line chemotherapy in advanced ovarian cancer: an interim analysis of a randomized phase III trial of the Arbeitsgemeinschaft Gynakologische Onkologie Ovarian Cancer Study Group. *Semin Oncol.* 1997;24(5 Suppl 15):S15-44-S15-52.
67. Recchia F, De Filippis S, Rosselli M, Saggio G, Carta G, Rea S. Primary chemotherapy in stage IV ovarian cancer. A prospective phase II study. *Eur J Gynaecol Oncol.* 2001;22(4):287-291.
68. Schink JC, Weller E, Harris LS, et al. Outpatient taxol and carboplatin chemotherapy for suboptimally debulked epithelial carcinoma of the ovary results in improved quality of life: an Eastern Cooperative Oncology Group Phase II Study (E2E93). *Cancer J.* 2001;7(2):155-164.

69. Ozols RF. Paclitaxel plus carboplatin in the treatment of ovarian cancer. *Semin Oncol.* 1999;26(1 Suppl 2):84-89.
70. Roland PY, Barnes MN, Niwas S, et al. Response to salvage treatment in recurrent ovarian cancer treated initially with paclitaxel and platinum-based combination regimens. *Gynecol Oncol.* 1998;68(2):178-182.
71. Skarlos DV, Aravantinos G, Kosmidis P, et al. Paclitaxel with carboplatin versus paclitaxel with carboplatin alternating with cisplatin as first-line chemotherapy in advanced epithelial ovarian cancer: preliminary results of a Hellenic Cooperative Oncology Group study. *Semin Oncol.* 1997;24(5 Suppl 15):S15-57-S15-61.
72. Pujade-Lauraine E, Guastalla JP, Weber B, et al. Efficacy and safety of the combination paclitaxel/carboplatin in patients with previously treated advanced ovarian carcinoma: a multicenter French Groupe des Investigateurs Nationaux pour l'Etude des Cancers Ovariens phase II study. *Semin Oncol.* 1997;24(5 Suppl 15):S15-30-S15-35.
73. Meerpohl HG, du Bois A, Luck HJ, et al. Paclitaxel combined with carboplatin in the first-line treatment of advanced ovarian cancer: a phase I trial. *Semin Oncol.* 1997;24(1 Suppl 2):S2-17-S2-22.
74. World Health Organization. GLOBOCAN 2012: Estimated cancer incidence, mortality and prevalence worldwide in 2012. Lyon, France: International Agency for Research on Cancer.[Links]. 2014.
75. Reis L, Eisner M, Kosary C, et al. SEER cancer statistics review, 1975–2002. National Cancer Institute, Bethesda, MD. 2005;6.
76. Howlader N, Noone A, Krapcho M, et al. SEER Cancer Statistics Review (CSR) 1975-2010. Bethesda: National Cancer Institute. 2014.
77. Travis WD, Brambilla E, Nicholson AG, et al. The 2015 World Health Organization Classification of Lung Tumors: Impact of Genetic, Clinical and Radiologic Advances Since the 2004 Classification. *J Thorac Oncol.* 2015;10(9):1243-1260.
78. Anonymous Lung. In: Edge S, Byrd DR, Compton CC, Fritz AG, Greene FL and Trotti A, eds. *AJCC Cancer Staging Manual.* New York, NY: Springer; 2010:253-270.
79. Kenfield SA, Wei EK, Stampfer MJ, Rosner BA, Colditz GA. Comparison of aspects of smoking among the four histological types of lung cancer. *Tob Control.* 2008;17(3):198-204.
80. Wingo PA, Ries LA, Giovino GA, et al. Annual report to the nation on the status of cancer, 1973-1996, with a special section on lung cancer and tobacco smoking. *J Natl Cancer Inst.* 1999;91(8):675-690.
81. PDQ Adult Treatment Editorial Board. Non-Small Cell Lung Cancer Treatment (PDQ(R)): Health Professional Version. In: Anonymous PDQ Cancer Information Summaries. Bethesda (MD): 2002:.
82. Sayar A, Turna A, Kilicgun A, Solak O, Urer N, Gurses A. Prognostic significance of surgical-pathologic multiple-station N1 disease in non-small cell carcinoma of the lung. *Eur J Cardiothorac Surg.* 2004;25(3):434-438.
83. Tanaka F, Yanagihara K, Otake Y, et al. Prognostic factors in patients with resected pathologic (p-) T1-2N1M0 non-small cell lung cancer (NSCLC). *Eur J Cardiothorac Surg.* 2001;19(5):555-561.
84. Riquet M, Manac'h D, Le Pimpec-Barthes F, Dujon A, Chehab A. Prognostic significance of surgical-pathologic N1 disease in non-small cell carcinoma of the lung. *Ann Thorac Surg.* 1999;67(6):1572-1576.
85. van Velzen E, Snijder RJ, Brutel de la Riviere A, Elbers HR, van den Bosch JM. Lymph node type as a prognostic factor for survival in T2 N1 M0 non-small cell lung carcinoma. *Ann Thorac Surg.* 1997;63(5):1436-1440.
86. Vansteenkiste JF, De Leyn PR, Deneffe GJ, et al. Survival and prognostic factors in resected N2 non-small cell lung cancer: a study of 140 cases. Leuven Lung Cancer Group. *Ann Thorac Surg.* 1997;63(5):1441-1450.
87. Naruke T, Goya T, Tsuchiya R, Suemasu K. Prognosis and survival in resected lung carcinoma based on the new international staging system. *J Thorac Cardiovasc Surg.* 1988;96(3):440-447.
88. Pao W and Girard N. New driver mutations in non-small-cell lung cancer. *Lancet Oncol.* 2011;12(2):175-180.
89. Lin JJ, Cardarella S, Lydon CA, et al. Five-Year Survival in EGFR-Mutant Metastatic Lung Adenocarcinoma Treated with EGFR-TKIs. *J Thorac Oncol.* 2016;11(4):556-565.
90. Tiseo M, Gelsomino F, Boggiani D, et al. EGFR and EML4-ALK gene mutations in NSCLC: a case report of erlotinib-resistant patient with both concomitant mutations. *Lung Cancer.* 2011;71(2):241-243.
91. Villaruz LC, Socinski MA, Abberbock S, et al. Clinicopathologic features and outcomes of patients with lung adenocarcinomas harboring BRAF mutations in the Lung Cancer Mutation Consortium. *Cancer.* 2015;121(3):448-456.

92. Pignon JP, Tribodet H, Scagliotti GV, et al. Lung adjuvant cisplatin evaluation: a pooled analysis by the LACE Collaborative Group. *J Clin Oncol.* 2008;26(21):3552-3559.
93. Albain KS, Rusch VW, Crowley JJ, et al. Concurrent cisplatin/etoposide plus chest radiotherapy followed by surgery for stages IIIA (N2) and IIIB non-small-cell lung cancer: mature results of Southwest Oncology Group phase II study 8805. *J Clin Oncol.* 1995;13(8):1880-1892.
94. Gilligan D, Nicolson M, Smith I, et al. Preoperative chemotherapy in patients with resectable non-small cell lung cancer: results of the MRC LU22/NVALT 2/EORTC 08012 multicentre randomised trial and update of systematic review. *Lancet.* 2007;369(9577):1929-1937.
95. Suntharalingam M, Paulus R, Edelman MJ, et al. Radiation therapy oncology group protocol 02-29: a phase II trial of neoadjuvant therapy with concurrent chemotherapy and full-dose radiation therapy followed by surgical resection and consolidative therapy for locally advanced non-small cell carcinoma of the lung. *Int J Radiat Oncol Biol Phys.* 2012;84(2):456-463.
96. Winton T, Livingston R, Johnson D, et al. Vinorelbine plus cisplatin vs. observation in resected non-small-cell lung cancer. *N Engl J Med.* 2005;352(25):2589-2597.
97. Arriagada R, Bergman B, Dunant A, et al. Cisplatin-based adjuvant chemotherapy in patients with completely resected non-small-cell lung cancer. *N Engl J Med.* 2004;350(4):351-360.
98. Strauss GM, Herndon JE, 2nd, Maddaus MA, et al. Adjuvant paclitaxel plus carboplatin compared with observation in stage IB non-small-cell lung cancer: CALGB 9633 with the Cancer and Leukemia Group B, Radiation Therapy Oncology Group, and North Central Cancer Treatment Group Study Groups. *J Clin Oncol.* 2008;26(31):5043-5051.
99. Douillard JY, Rosell R, De Lena M, et al. Adjuvant vinorelbine plus cisplatin versus observation in patients with completely resected stage IB-IIIa non-small-cell lung cancer (Adjuvant Navelbine International Trialist Association [ANITA]): a randomised controlled trial. *Lancet Oncol.* 2006;7(9):719-727.
100. Pepe C, Hasan B, Winton TL, et al. Adjuvant vinorelbine and cisplatin in elderly patients: National Cancer Institute of Canada and Intergroup Study JBR.10. *J Clin Oncol.* 2007;25(12):1553-1561.
101. Miller JJ, Jr and Phillips TW. Neodymium:YAG laser and brachytherapy in the management of inoperable bronchogenic carcinoma. *Ann Thorac Surg.* 1990;50(2):190-5; discussion 195-6.
102. Weick JK, Crowley J, Natale RB, et al. A randomized trial of five cisplatin-containing treatments in patients with metastatic non-small-cell lung cancer: a Southwest Oncology Group study. *J Clin Oncol.* 1991;9(7):1157-1162.
103. Ardizzoni A, Boni L, Tiseo M, et al. Cisplatin- versus carboplatin-based chemotherapy in first-line treatment of advanced non-small-cell lung cancer: an individual patient data meta-analysis. *J Natl Cancer Inst.* 2007;99(11):847-857.
104. D'Addario G, Pintilie M, Leighl NB, Feld R, Cerny T, Shepherd FA. Platinum-based versus non-platinum-based chemotherapy in advanced non-small-cell lung cancer: a meta-analysis of the published literature. *J Clin Oncol.* 2005;23(13):2926-2936.
105. Sandler A, Gray R, Perry MC, et al. Paclitaxel-carboplatin alone or with bevacizumab for non-small-cell lung cancer. *N Engl J Med.* 2006;355(24):2542-2550.
106. Reck M, von Pawel J, Zatloukal P, et al. Phase III trial of cisplatin plus gemcitabine with either placebo or bevacizumab as first-line therapy for nonsquamous non-small-cell lung cancer: AVAil. *J Clin Oncol.* 2009;27(8):1227-1234.
107. Lynch TJ, Patel T, Dreisbach L, et al. Cetuximab and first-line taxane/carboplatin chemotherapy in advanced non-small-cell lung cancer: results of the randomized multicenter phase III trial BMS099. *J Clin Oncol.* 2010;28(6):911-917.
108. Pirker R, Pereira JR, Szczesna A, et al. Cetuximab plus chemotherapy in patients with advanced non-small-cell lung cancer (FLEX): an open-label randomised phase III trial. *Lancet.* 2009;373(9674):1525-1531.
109. Paz-Ares L, Mezger J, Ciuleanu TE, et al. Necitumumab plus pemetrexed and cisplatin as first-line therapy in patients with stage IV non-squamous non-small-cell lung cancer (INSPIRE): an open-label, randomised, controlled phase 3 study. *Lancet Oncol.* 2015;16(3):328-337.
110. Thatcher N, Hirsch FR, Luft AV, et al. Necitumumab plus gemcitabine and cisplatin versus gemcitabine and cisplatin alone as first-line therapy in patients with stage IV squamous non-small-cell lung cancer (SQUIRE): an open-label, randomised, controlled phase 3 trial. *Lancet Oncol.* 2015;16(7):763-774.
111. PORT Meta-analysis Trialists Group. Postoperative radiotherapy for non-small cell lung cancer. *Cochrane Database Syst Rev.* 2005;(2)(2):CD002142.

112. Lally BE, Zelterman D, Colasanto JM, Haffty BG, Detterbeck FC, Wilson LD. Postoperative radiotherapy for stage II or III non-small-cell lung cancer using the surveillance, epidemiology, and end results database. *J Clin Oncol.* 2006;24(19):2998-3006.
113. Johnstone DW, Byhardt RW, Ettinger D, Scott CB. Phase III study comparing chemotherapy and radiotherapy with preoperative chemotherapy and surgical resection in patients with non-small-cell lung cancer with spread to mediastinal lymph nodes (N2); final report of RTOG 89-01. Radiation Therapy Oncology Group. *Int J Radiat Oncol Biol Phys.* 2002;54(2):365-369.
114. Taylor NA, Liao ZX, Cox JD, et al. Equivalent outcome of patients with clinical Stage IIIA non-small-cell lung cancer treated with concurrent chemoradiation compared with induction chemotherapy followed by surgical resection. *Int J Radiat Oncol Biol Phys.* 2004;58(1):204-212.
115. van Meerbeeck JP, Kramer GW, Van Schil PE, et al. Randomized controlled trial of resection versus radiotherapy after induction chemotherapy in stage IIIA-N2 non-small-cell lung cancer. *J Natl Cancer Inst.* 2007;99(6):442-450.
116. Auperin A, Le Pechoux C, Pignon JP, et al. Concomitant radio-chemotherapy based on platin compounds in patients with locally advanced non-small cell lung cancer (NSCLC): a meta-analysis of individual data from 1764 patients. *Ann Oncol.* 2006;17(3):473-483.
117. Furuse K, Fukuoka M, Kawahara M, et al. Phase III study of concurrent versus sequential thoracic radiotherapy in combination with mitomycin, vindesine, and cisplatin in unresectable stage III non-small-cell lung cancer. *J Clin Oncol.* 1999;17(9):2692-2699.
118. Zatloukal P, Petruzella L, Zemanova M, et al. Concurrent versus sequential chemoradiotherapy with cisplatin and vinorelbine in locally advanced non-small cell lung cancer: a randomized study. *Lung Cancer.* 2004;46(1):87-98.
119. Hanna N, Neubauer M, Yiannoutsos C, et al. Phase III study of cisplatin, etoposide, and concurrent chest radiation with or without consolidation docetaxel in patients with inoperable stage III non-small-cell lung cancer: the Hoosier Oncology Group and U.S. Oncology. *J Clin Oncol.* 2008;26(35):5755-5760.
120. Lester JF, Macbeth FR, Toy E, Coles B. Palliative radiotherapy regimens for non-small cell lung cancer. *Cochrane Database Syst Rev.* 2006;(4):CD002143.
121. Ung YC, Yu E, Falkson C, et al. The role of high-dose-rate brachytherapy in the palliation of symptoms in patients with non-small-cell lung cancer: a systematic review. *Brachytherapy.* 2006;5(3):189-202.
122. Wisnivesky JP, Yankellevitz D, Henschke CI. Stage of lung cancer in relation to its size: part 2. Evidence. *Chest.* 2005;127(4):1136-1139.
123. Mountain CF. Revisions in the International System for Staging Lung Cancer. *Chest.* 1997;111(6):1710-1717.
124. Chen YM, Perng RP, Lee YC, et al. Paclitaxel plus carboplatin, compared with paclitaxel plus gemcitabine, shows similar efficacy while more cost-effective: a randomized phase II study of combination chemotherapy against inoperable non-small-cell lung cancer previously untreated. *Ann Oncol.* 2002;13(1):108-115.
125. Hainsworth JD and Greco FA. One-hour paclitaxel in the treatment of non-small cell lung cancer. *Semin Oncol.* 1996;23(6 Suppl 16):98-101.
126. Langer CJ, Millenson M, Rosvold E, et al. Paclitaxel (1-hour) and carboplatin (area under the concentration-time curve 7.5) in advanced non-small cell lung cancer: a phase II study of the Fox Chase Cancer Center and its network. *Semin Oncol.* 1997;24(4 Suppl 12):S12-81-S12-88.
127. Johnson DH, Paul DM, Hande KR, DeVore R. Paclitaxel plus carboplatin in the treatment of patients with advanced lung cancer: a Vanderbilt University Cancer Center phase II trial (LUN-46). *Semin Oncol.* 1996;23(6 Suppl 16):42-46.
128. Langer CJ, Leighton JC, Comis RL, et al. Paclitaxel and carboplatin in combination in the treatment of advanced non-small-cell lung cancer: a phase II toxicity, response, and survival analysis. *J Clin Oncol.* 1995;13(8):1860-1870.
129. Glorieux P, Ortmanns P, Marien S, et al. Multi-center study of two dose levels of paclitaxel with carboplatin in locally advanced and metastatic non-small cell lung cancer (NSCLC). *Anticancer Res.* 2001;21(2B):1487-1494.
130. Rosenberg B, VanCamp L, Trosko JE, Mansour VH. Platinum compounds: a new class of potent antitumour agents. *Nature.* 1969;222(5191):385-386.
131. Rabik CA and Dolan ME. Molecular mechanisms of resistance and toxicity associated with platinating agents. *Cancer Treat Rev.* 2007;33(1):9-23.

132. Wheate NJ, Walker S, Craig GE, Oun R. The status of platinum anticancer drugs in the clinic and in clinical trials. *Dalton Trans.* 2010;39(35):8113-8127.
133. Fang G, Haznadar M, Wang W, et al. High-order SNP combinations associated with complex diseases: efficient discovery, statistical power and functional interactions. *PLoS One.* 2012;7(4):e33531.
134. American Psychological Association (APA).

The publication manual of the American Psychological Association Washington, DC: American Psychological Association; 2010.

135. Kartalou M and Essigmann JM. Mechanisms of resistance to cisplatin. *Mutat Res.* 2001;478(1-2):23-43.
136. Guo Y, Smith K, Petris MJ. Cisplatin stabilizes a multimeric complex of the human Ctr1 copper transporter: requirement for the extracellular methionine-rich clusters. *J Biol Chem.* 2004;279(45):46393-46399.
137. Howell SB, Safaei R, Larson CA, Sailor MJ. Copper transporters and the cellular pharmacology of the platinum-containing cancer drugs. *Mol Pharmacol.* 2010;77(6):887-894.
138. Sprowl JA, Ness RA, Sparreboom A. Polymorphic transporters and platinum pharmacodynamics. *Drug Metab Pharmacokinet.* 2013;28(1):19-27.
139. Gupta A, Srivastava S, Prasad R, et al. Smoking intensity, oxidative stress and chemotherapy in nonsmall cell lung cancer: a correlated prognostic study. *Biosci Trends.* 2009;3(5):191-199.
140. Ciarimboli G. Membrane transporters as mediators of Cisplatin effects and side effects. *Scientifica (Cairo).* 2012;2012:473829.
141. Sinani D, Adle DJ, Kim H, Lee J. Distinct mechanisms for Ctr1-mediated copper and cisplatin transport. *J Biol Chem.* 2007;282(37):26775-26785.
142. Holzer AK, Samimi G, Katano K, et al. The copper influx transporter human copper transport protein 1 regulates the uptake of cisplatin in human ovarian carcinoma cells. *Mol Pharmacol.* 2004;66(4):817-823.
143. Song IS, Savaraj N, Siddik ZH, et al. Role of human copper transporter Ctr1 in the transport of platinum-based antitumor agents in cisplatin-sensitive and cisplatin-resistant cells. *Mol Cancer Ther.* 2004;3(12):1543-1549.
144. Kalayda GV, Wagner CH, Jaehde U. Relevance of copper transporter 1 for cisplatin resistance in human ovarian carcinoma cells. *J Inorg Biochem.* 2012;116:1-10.
145. Kilari D, Guancial E, Kim ES. Role of copper transporters in platinum resistance. *World J Clin Oncol.* 2016;7(1):106-113.
146. Zisowsky J, Koegel S, Leyers S, et al. Relevance of drug uptake and efflux for cisplatin sensitivity of tumor cells. *Biochem Pharmacol.* 2007;73(2):298-307.
147. Ishida S, Lee J, Thiele DJ, Herskowitz I. Uptake of the anticancer drug cisplatin mediated by the copper transporter Ctr1 in yeast and mammals. *Proc Natl Acad Sci U S A.* 2002;99(22):14298-14302.
148. Larson CA, Blair BG, Safaei R, Howell SB. The role of the mammalian copper transporter 1 in the cellular accumulation of platinum-based drugs. *Mol Pharmacol.* 2009;75(2):324-330.
149. Jandial DD, Messer K, Farshchi-Heydari S, Pu M, Howell SB. Tumor platinum concentration following intraperitoneal administration of cisplatin versus carboplatin in an ovarian cancer model. *Gynecol Oncol.* 2009;115(3):362-366.
150. Blair BG, Larson CA, Safaei R, Howell SB. Copper transporter 2 regulates the cellular accumulation and cytotoxicity of Cisplatin and Carboplatin. *Clin Cancer Res.* 2009;15(13):4312-4321.
151. Lancaster CS, Sprowl JA, Walker AL, Hu S, Gibson AA, Sparreboom A. Modulation of OATP1B-type transporter function alters cellular uptake and disposition of platinum chemotherapeutics. *Mol Cancer Ther.* 2013;12(8):1537-1544.
152. Mangala LS, Zuzel V, Schmandt R, et al. Therapeutic Targeting of ATP7B in Ovarian Carcinoma. *Clin Cancer Res.* 2009;15(11):3770-3780.
153. Martinez-Balibrea E, Martinez-Cardus A, Musulen E, et al. Increased levels of copper efflux transporter ATP7B are associated with poor outcome in colorectal cancer patients receiving oxaliplatin-based chemotherapy. *Int J Cancer.* 2009;124(12):2905-2910.
154. Samimi G, Katano K, Holzer AK, Safaei R, Howell SB. Modulation of the cellular pharmacology of cisplatin and its analogs by the copper exporters ATP7A and ATP7B. *Mol Pharmacol.* 2004;66(1):25-32.
155. Safaei R. Role of copper transporters in the uptake and efflux of platinum containing drugs. *Cancer Lett.* 2006;234(1):34-39.

156. Leonhardt K, Gebhardt R, Mossner J, Lutsenko S, Huster D. Functional interactions of Cu-ATPase ATP7B with cisplatin and the role of ATP7B in the resistance of cells to the drug. *J Biol Chem.* 2009;284(12):7793-7802.
157. Tadini-Buoninsegni F, Bartolommei G, Moncelli MR, et al. Translocation of platinum anticancer drugs by human copper ATPases ATP7A and ATP7B. *Angew Chem Int Ed Engl.* 2014;53(5):1297-1301.
158. Komatsu M, Sumizawa T, Mutoh M, et al. Copper-transporting P-type adenosine triphosphatase (ATP7B) is associated with cisplatin resistance. *Cancer Res.* 2000;60(5):1312-1316.
159. Calandrini V, Arnesano F, Galliani A, et al. Platination of the copper transporter ATP7A involved in anticancer drug resistance. *Dalton Trans.* 2014;43(31):12085-12094.
160. Kanzaki A, Toi M, Neamati N, et al. Copper-transporting P-type adenosine triphosphatase (ATP7B) is expressed in human breast carcinoma. *Jpn J Cancer Res.* 2002;93(1):70-77.
161. Nakayama K, Kanzaki A, Ogawa K, Miyazaki K, Neamati N, Takebayashi Y. Copper-transporting P-type adenosine triphosphatase (ATP7B) as a cisplatin based chemoresistance marker in ovarian carcinoma: comparative analysis with expression of MDR1, MRP1, MRP2, LRP and BCRP. *Int J Cancer.* 2002;101(5):488-495.
162. Katano K, Kondo A, Safaei R, et al. Acquisition of resistance to cisplatin is accompanied by changes in the cellular pharmacology of copper. *Cancer Res.* 2002;62(22):6559-6565.
163. Rabik CA, Maryon EB, Kasza K, Shafer JT, Bartnik CM, Dolan ME. Role of copper transporters in resistance to platinating agents. *Cancer Chemother Pharmacol.* 2009;64(1):133-142.
164. Francis MJ, Jones EE, Levy ER, Martin RL, Ponnambalam S, Monaco AP. Identification of a di-leucine motif within the C terminus domain of the Menkes disease protein that mediates endocytosis from the plasma membrane. *J Cell Sci.* 1999;112 (Pt 11)(Pt 11):1721-1732.
165. Stephenson SE, Dubach D, Lim CM, Mercer JF, La Fontaine S. A single PDZ domain protein interacts with the Menkes copper ATPase, ATP7A. A new protein implicated in copper homeostasis. *J Biol Chem.* 2005;280(39):33270-33279.
166. Pase L, Voskoboinik I, Greenough M, Camakaris J. Copper stimulates trafficking of a distinct pool of the Menkes copper ATPase (ATP7A) to the plasma membrane and diverts it into a rapid recycling pool. *Biochem J.* 2004;378(Pt 3):1031-1037.
167. Nyasae L, Bustos R, Braiterman L, Eipper B, Hubbard A. Dynamics of endogenous ATP7A (Menkes protein) in intestinal epithelial cells: copper-dependent redistribution between two intracellular sites. *Am J Physiol Gastrointest Liver Physiol.* 2007;292(4):G1181-94.
168. Borst P, Zelcer N, van Helvoort A. ABC transporters in lipid transport. *Biochim Biophys Acta.* 2000;1486(1):128-144.
169. Cole SP, Bhardwaj G, Gerlach JH, et al. Overexpression of a transporter gene in a multidrug-resistant human lung cancer cell line. *Science.* 1992;258(5088):1650-1654.
170. Dean M, Rzhetsky A, Allikmets R. The human ATP-binding cassette (ABC) transporter superfamily. *Genome Res.* 2001;11(7):1156-1166.
171. Moyer AM, Sun Z, Batzler AJ, et al. Glutathione pathway genetic polymorphisms and lung cancer survival after platinum-based chemotherapy. *Cancer Epidemiol Biomarkers Prev.* 2010;19(3):811-821.
172. Cervenak J, Andrikovics H, Ozvegy-Laczka C, et al. The role of the human ABCG2 multidrug transporter and its variants in cancer therapy and toxicology. *Cancer Lett.* 2006;234(1):62-72.
173. Ceckova M, Vackova Z, Radilova H, Libra A, Buncek M, Staud F. Effect of ABCG2 on cytotoxicity of platinum drugs: interference of EGFP. *Toxicol In Vitro.* 2008;22(8):1846-1852.
174. Liedert B, Materna V, Schadendorf D, Thomale J, Lage H. Overexpression of cMOAT (MRP2/ABCC2) is associated with decreased formation of platinum-DNA adducts and decreased G2-arrest in melanoma cells resistant to cisplatin. *J Invest Dermatol.* 2003;121(1):172-176.
175. Materna V, Liedert B, Thomale J, Lage H. Protection of platinum-DNA adduct formation and reversal of cisplatin resistance by anti-MRP2 hammerhead ribozymes in human cancer cells. *Int J Cancer.* 2005;115(3):393-402.
176. Suzuki T, Nishio K, Tanabe S. The MRP family and anticancer drug metabolism. *Curr Drug Metab.* 2001;2(4):367-377.
177. Hayes JD and Pulford DJ. The glutathione S-transferase supergene family: regulation of GST and the contribution of the isoenzymes to cancer chemoprotection and drug resistance. *Crit Rev Biochem Mol Biol.* 1995;30(6):445-600.
178. Sawers L, Ferguson MJ, Ihrig BR, et al. Glutathione S-transferase P1 (GSTP1) directly influences platinum drug chemosensitivity in ovarian tumour cell lines. *Br J Cancer.* 2014;111(6):1150-1158.

179. Yanagie H, Hisa T, Ogata A, et al. Improvement of sensitivity to platinum compound with siRNA knockdown of upregulated genes in platinum complex-resistant ovarian cancer cells in vitro. *Biomed Pharmacother.* 2009;63(8):553-560.
180. Parekh H and Simpkins H. Cross-resistance and collateral sensitivity to natural product drugs in cisplatin-sensitive and -resistant rat lymphoma and human ovarian carcinoma cells. *Cancer Chemother Pharmacol.* 1996;37(5):457-462.
181. Chen J, Solomides C, Simpkins H. Sensitization of mesothelioma cells to platinum-based chemotherapy by GSTpi knockdown. *Biochem Biophys Res Commun.* 2014;447(1):77-82.
182. Kotoh S, Naito S, Yokomizo A, Kohno K, Kuwano M, Kumazawa J. Enhanced expression of gamma-glutamylcysteine synthetase and glutathione S-transferase genes in cisplatin-resistant bladder cancer cells with multidrug resistance phenotype. *J Urol.* 1997;157(3):1054-1058.
183. Tozawa K, Oshima T, Kobayashi T, et al. Oxaliplatin in treatment of the cisplatin-resistant MKN45 cell line of gastric cancer. *Anticancer Res.* 2008;28(4B):2087-2092.
184. Brown DP, Chin-Sinex H, Nie B, Mendonca MS, Wang M. Targeting superoxide dismutase 1 to overcome cisplatin resistance in human ovarian cancer. *Cancer Chemother Pharmacol.* 2009;63(4):723-730.
185. Pasheva EA, Ugrinova I, Spassovska NC, Pashev IG. The binding affinity of HMG1 protein to DNA modified by cis-platin and its analogs correlates with their antitumor activity. *Int J Biochem Cell Biol.* 2002;34(1):87-92.
186. He Y, Yuan J, Qiao Y, et al. The role of carrier ligands of platinum(II) anticancer complexes in the protein recognition of Pt-DNA adducts. *Chem Commun (Camb).* 2015;51(74):14064-14067.
187. Yusein-Myashkova S, Ugrinova I, Pasheva E. Non-histone protein HMGB1 inhibits the repair of damaged DNA by cisplatin in NIH-3T3 murine fibroblasts. *BMB Rep.* 2016;49(2):99-104.
188. He Q, Ohndorf UM, Lippard SJ. Intercalating residues determine the mode of HMG1 domains A and B binding to cisplatin-modified DNA. *Biochemistry.* 2000;39(47):14426-14435.
189. Parsons JL, Dianova II, Allinson SL, Dianov GL. Poly(ADP-ribose) polymerase-1 protects excessive DNA strand breaks from deterioration during repair in human cell extracts. *FEBS J.* 2005;272(8):2012-2021.
190. Guggenheim ER, Xu D, Zhang CX, Chang PV, Lippard SJ. Photoaffinity isolation and identification of proteins in cancer cell extracts that bind to platinum-modified DNA. *Chembiochem.* 2009;10(1):141-157.
191. Guggenheim ER, Ondrus AE, Movassaghi M, Lippard SJ. Poly(ADP-ribose) polymerase-1 activity facilitates the dissociation of nuclear proteins from platinum-modified DNA. *Bioorg Med Chem.* 2008;16(23):10121-10128.
192. Zhang CX, Chang PV, Lippard SJ. Identification of nuclear proteins that interact with platinum-modified DNA by photoaffinity labeling. *J Am Chem Soc.* 2004;126(21):6536-6537.
193. Zhu G, Chang P, Lippard SJ. Recognition of platinum-DNA damage by poly(ADP-ribose) polymerase-1. *Biochemistry.* 2010;49(29):6177-6183.
194. Martin A and Scharff MD. AID and mismatch repair in antibody diversification. *Nat Rev Immunol.* 2002;2(8):605-614.
195. Xiao X, Melton DW, Gourley C. Mismatch repair deficiency in ovarian cancer -- molecular characteristics and clinical implications. *Gynecol Oncol.* 2014;132(2):506-512.
196. Durant ST, Morris MM, Illand M, et al. Dependence on RAD52 and RAD1 for anticancer drug resistance mediated by inactivation of mismatch repair genes. *Curr Biol.* 1999;9(1):51-54.
197. Fink D, Nebel S, Aebi S, et al. The role of DNA mismatch repair in platinum drug resistance. *Cancer Res.* 1996;56(21):4881-4886.
198. Adachi M, Ijichi K, Hasegawa Y, et al. Hypersensitivity to cisplatin after hRev3 mRNA knockdown in head and neck squamous cell carcinoma cells. *Mol Med Rep.* 2008;1(5):695-698.
199. Fink D, Nebel S, Aebi S, Nehme A, Howell S. Loss of DNA mismatch repair due to knockout of MSH2 or PMS2 results in resistance to cisplatin and carboplatin. *Int J Oncol.* 1997;11(3):539-542.
200. Takahashi M, Koi M, Balaguer F, Boland CR, Goel A. MSH3 mediates sensitization of colorectal cancer cells to cisplatin, oxaliplatin, and a poly(ADP-ribose) polymerase inhibitor. *J Biol Chem.* 2011;286(14):12157-12165.
201. Fedier A, Ruefenacht UB, Schwarz VA, Haller U, Fink D. Increased sensitivity of p53-deficient cells to anticancer agents due to loss of Pms2. *Br J Cancer.* 2002;87(9):1027-1033.
202. de Laat WL, Jaspers NG, Hoeijmakers JH. Molecular mechanism of nucleotide excision repair. *Genes Dev.* 1999;13(7):768-785.
203. Chang IY, Kim MH, Kim HB, et al. Small interfering RNA-induced suppression of ERCC1 enhances sensitivity of human cancer cells to cisplatin. *Biochem Biophys Res Commun.* 2005;327(1):225-233.

204. O'Grady S, Finn SP, Cuffe S, Richard DJ, O'Byrne KJ, Barr MP. The role of DNA repair pathways in cisplatin resistant lung cancer. *Cancer Treat Rev.* 2014;40(10):1161-1170.
205. Ge R, Liu L, Dai W, et al. Xeroderma Pigmentosum Group A Promotes Autophagy to Facilitate Cisplatin Resistance in Melanoma Cells through the Activation of PARP1. *J Invest Dermatol.* 2016;136(6):1219-1228.
206. Zhang Q, Wu JX, Wei YP, Hao JQ, Huang SY, Dong L. Effect of XPA expression on the chemotherapy sensitivity of A549/DDP cells. *Xi Bao Yu Fen Zi Mian Yi Xue Za Zhi.* 2012;28(11):1182-1184.
207. Reed E, Yu JJ, Davies A, Gannon J, Armentrout SL. Clear cell tumors have higher mRNA levels of ERCC1 and XPB than other histological types of epithelial ovarian cancer. *Clin Cancer Res.* 2003;9(14):5299-5305.
208. Dabholkar M, Thornton K, Vionnet J, Bostick-Bruton F, Yu JJ, Reed E. Increased mRNA levels of xeroderma pigmentosum complementation group B (XPB) and Cockayne's syndrome complementation group B (CSB) without increased mRNA levels of multidrug-resistance gene (MDR1) or metallothionein-II (MT-II) in platinum-resistant human ovarian cancer tissues. *Biochem Pharmacol.* 2000;60(11):1611-1619.
209. McNeil EM, Astell KR, Ritchie AM, et al. Inhibition of the ERCC1-XPF structure-specific endonuclease to overcome cancer chemoresistance. *DNA Repair (Amst).* 2015;31:19-28.
210. Arora S, Kothandapani A, Tillison K, Kalman-Maltese V, Patrick SM. Downregulation of XPF-ERCC1 enhances cisplatin efficacy in cancer cells. *DNA Repair (Amst).* 2010;9(7):745-753.
211. Mukai M, Kanzaki A, Chen ZS, et al. Enhanced nucleotide excision repair in cisplatin resistant human KB carcinoma cells. *Oncol Rep.* 2002;9(4):839-844.
212. Graf N, Ang WH, Zhu G, Myint M, Lippard SJ. Role of endonucleases XPF and XPG in nucleotide excision repair of platinated DNA and cisplatin/oxaliplatin cytotoxicity. *Chembiochem.* 2011;12(7):1115-1123.
213. Lin K, Ye D, Xie X. Protein expression levels of excision repair cross-complementation group 1 and xeroderma pigmentosum D correlate with response to platinum-based chemotherapy in the patients with advanced epithelial ovarian cancer. *Int J Gynecol Cancer.* 2008;18(5):1007-1012.
214. Arienti C, Tesei A, Verdecchia GM, et al. Peritoneal carcinomatosis from ovarian cancer: chemosensitivity test and tissue markers as predictors of response to chemotherapy. *J Transl Med.* 2011;9:94-5876-9-94.
215. Thacker J and Zdzienicka MZ. The XRCC genes: expanding roles in DNA double-strand break repair. *DNA Repair (Amst).* 2004;3(8-9):1081-1090.
216. Xu W, Wang S, Chen Q, et al. TXNL1-XRCC1 pathway regulates cisplatin-induced cell death and contributes to resistance in human gastric cancer. *Cell Death Dis.* 2014;5:e1055.
217. Abdel-Fatah T, Sultana R, Abbotts R, et al. Clinicopathological and functional significance of XRCC1 expression in ovarian cancer. *Int J Cancer.* 2013;132(12):2778-2786.
218. Lone S, Townson SA, Uljon SN, et al. Human DNA polymerase kappa encircles DNA: implications for mismatch extension and lesion bypass. *Mol Cell.* 2007;25(4):601-614.
219. Doles J, Oliver TG, Cameron ER, et al. Suppression of Rev3, the catalytic subunit of Pol{zeta}, sensitizes drug-resistant lung tumors to chemotherapy. *Proc Natl Acad Sci U S A.* 2010;107(48):20786-20791.
220. Wu F, Lin X, Okuda T, Howell SB. DNA polymerase zeta regulates cisplatin cytotoxicity, mutagenicity, and the rate of development of cisplatin resistance. *Cancer Res.* 2004;64(21):8029-8035.
221. Giovannetti E, Toffalorio F, De Pas T, Peters GJ. Pharmacogenetics of conventional chemotherapy in non-small-cell lung cancer: a changing landscape? *Pharmacogenomics.* 2012;13(9):1073-1086.
222. Ohrvik H and Thiele DJ. The role of Ctr1 and Ctr2 in mammalian copper homeostasis and platinum-based chemotherapy. *J Trace Elem Med Biol.* 2015;31:178-182.
223. Chen HH, Yan JJ, Chen WC, et al. Predictive and prognostic value of human copper transporter 1 (hCtr1) in patients with stage III non-small-cell lung cancer receiving first-line platinum-based doublet chemotherapy. *Lung Cancer.* 2012;75(2):228-234.
224. Warta R, Theile D, Mogler C, et al. Association of drug transporter expression with mortality and progression-free survival in stage IV head and neck squamous cell carcinoma. *PLoS One.* 2014;9(9):e108908.
225. Kim ES, Tang X, Peterson DR, et al. Copper transporter CTR1 expression and tissue platinum concentration in non-small cell lung cancer. *Lung Cancer.* 2014;85(1):88-93.

226. Ogane N, Yasuda M, Kameda Y, et al. Prognostic value of organic anion transporting polypeptide 1B3 and copper transporter 1 expression in endometrial cancer patients treated with paclitaxel and carboplatin. *Biomed Res.* 2013;34(3):143-151.
227. Yang T, Chen M, Chen T, Thakur A. Expression of the copper transporters hCtr1, ATP7A and ATP7B is associated with the response to chemotherapy and survival time in patients with resected non-small cell lung cancer. *Oncol Lett.* 2015;10(4):2584-2590.
228. Xu X, Duan L, Zhou B, Ma R, Zhou H, Liu Z. Genetic polymorphism of copper transporter protein 1 is related to platinum resistance in Chinese non-small cell lung carcinoma patients. *Clin Exp Pharmacol Physiol.* 2012;39(9):786-792.
229. Negoro K, Yamano Y, Nakashima D, et al. Cross-resistance of platinum derivatives in H-1R, a cisplatin-resistant cell line. *Oncol Rep.* 2009;21(2):443-449.
230. Li ZH, Qiu MZ, Zeng ZL, et al. Copper-transporting P-type adenosine triphosphatase (ATP7A) is associated with platinum-resistance in non-small cell lung cancer (NSCLC). *J Transl Med.* 2012;10:21-5876-10-21.
231. Doherty B, Lawlor D, Gillet JP, Gottesman M, O'Leary JJ, Stordal B. Collateral sensitivity to cisplatin in KB-8-5-11 drug-resistant cancer cells. *Anticancer Res.* 2014;34(1):503-507.
232. Fukushima-Uesaka H, Saito Y, Maekawa K, et al. Genetic polymorphisms of copper- and platinum drug-efflux transporters ATP7A and ATP7B in Japanese cancer patients. *Drug Metab Pharmacokinet.* 2009;24(6):565-574.
233. Kim SH, Lee GW, Lee MJ, et al. Clinical significance of ERCC2 haplotype-tagging single nucleotide polymorphisms in patients with unresectable non-small cell lung cancer treated with first-line platinum-based chemotherapy. *Lung Cancer.* 2012;77(3):578-584.
234. Surowiak P, Materna V, Kaplenko I, et al. Augmented expression of metallothionein and glutathione S-transferase pi as unfavourable prognostic factors in cisplatin-treated ovarian cancer patients. *Virchows Arch.* 2005;447(3):626-633.
235. Zhang Q, Li K, Xu JH, et al. Role of ABCG2 expression driven by cisplatin in platinum-containing chemotherapy for gastric cancer. *World J Gastroenterol.* 2013;19(39):6630-6636.
236. Tian C, Ambrosone CB, Darcy KM, et al. Common variants in ABCB1, ABCC2 and ABCG2 genes and clinical outcomes among women with advanced stage ovarian cancer treated with platinum and taxane-based chemotherapy: a Gynecologic Oncology Group study. *Gynecol Oncol.* 2012;124(3):575-581.
237. Muller PJ, Dally H, Klappenecker CN, et al. Polymorphisms in ABCG2, ABCC3 and CNT1 genes and their possible impact on chemotherapy outcome of lung cancer patients. *Int J Cancer.* 2009;124(7):1669-1674.
238. Kim SH, Kim MJ, Cho YJ, et al. Clinical Significance of ABCG2 Haplotype-tagging Single Nucleotide Polymorphisms in Patients With Unresectable Non-Small Cell Lung Cancer Treated With First-line Platinum-based Chemotherapy. *Am J Clin Oncol.* 2015;38(3):294-299.
239. Cuffe S, Azad AK, Qiu X, et al. ABCC2 polymorphisms and survival in the Princess Margaret cohort study and the NCIC clinical trials group BR.24 trial of platinum-treated advanced stage non-small cell lung cancer patients. *Cancer Epidemiol.* 2016;41:50-56.
240. Han B, Gao G, Wu W, et al. Association of ABCC2 polymorphisms with platinum-based chemotherapy response and severe toxicity in non-small cell lung cancer patients. *Lung Cancer.* 2011;72(2):238-243.
241. Sun N, Sun X, Chen B, et al. MRP2 and GSTP1 polymorphisms and chemotherapy response in advanced non-small cell lung cancer. *Cancer Chemother Pharmacol.* 2010;65(3):437-446.
242. Brambila-Tapia AJ. MDR1 (ABCB1) polymorphisms: functional effects and clinical implications. *Rev Invest Clin.* 2013;65(5):445-454.
243. Wei HB, Hu J, Shang LH, et al. A meta-analytic review of ERCC1/MDR1 polymorphism and chemosensitivity to platinum in patients with advanced non-small cell lung cancer. *Chin Med J (Engl).* 2012;125(16):2902-2907.
244. Dogu GG, Kargi A, Turgut S, et al. MDR1 single nucleotide polymorphism C3435T in Turkish patients with non-small-cell lung cancer. *Gene.* 2012;506(2):404-407.
245. Pellegrino P, Cattaneo D, Clementi E. Role of ABCB1 C3435T in platinum-based therapy. *Arch Med Res.* 2012;43(4):329-30; author reply 331.
246. Yan PW, Huang XE, Yan F, Xu L, Jiang Y. Influence of MDR1 gene codon 3435 polymorphisms on outcome of platinum-based chemotherapy for advanced non small cell lung cancer. *Asian Pac J Cancer Prev.* 2011;12(9):2291-2294.

247. Kim HS, Kim MK, Chung HH, et al. Genetic polymorphisms affecting clinical outcomes in epithelial ovarian cancer patients treated with taxanes and platinum compounds: a Korean population-based study. *Gynecol Oncol.* 2009;113(2):264-269.
248. Chen S, Huo X, Lin Y, et al. Association of MDR1 and ERCC1 polymorphisms with response and toxicity to cisplatin-based chemotherapy in non-small-cell lung cancer patients. *Int J Hyg Environ Health.* 2010;213(2):140-145.
249. Lambrechts S, Lambrechts D, Despierre E, et al. Genetic variability in drug transport, metabolism or DNA repair affecting toxicity of chemotherapy in ovarian cancer. *BMC Pharmacol Toxicol.* 2015;16:2-015-0001-5.
250. Ohishi Y, Oda Y, Uchiumi T, et al. ATP-binding cassette superfamily transporter gene expression in human primary ovarian carcinoma. *Clin Cancer Res.* 2002;8(12):3767-3775.
251. Johnson C, Pankratz VS, Velazquez AI, et al. Candidate pathway-based genetic association study of platinum and platinum-taxane related toxicity in a cohort of primary lung cancer patients. *J Neurol Sci.* 2015;349(1-2):124-128.
252. Wang J, Li AJ, Karlan BY. Chemotherapy in epithelial ovarian cancer. *Curr Womens Health Rep.* 2002;2(1):20-26.
253. Hida T, Kuwabara M, Ariyoshi Y, et al. Serum glutathione S-transferase-pi level as a tumor marker for non-small cell lung cancer. Potential predictive value in chemotherapeutic response. *Cancer.* 1994;73(5):1377-1382.
254. Pillot GA, Read WL, Hennenfent KL, et al. A phase II study of irinotecan and carboplatin in advanced non-small cell lung cancer with pharmacogenomic analysis: final report. *J Thorac Oncol.* 2006;1(9):972-978.
255. Joerger M, Burgers SA, Baas P, et al. Germline polymorphisms in patients with advanced nonsmall cell lung cancer receiving first-line platinum-gemcitabine chemotherapy: a prospective clinical study. *Cancer.* 2012;118(9):2466-2475.
256. Ke HG, Li J, Shen Y, et al. Prognostic significance of GSTP1, XRCC1 and XRCC3 polymorphisms in non-small cell lung cancer patients. *Asian Pac J Cancer Prev.* 2012;13(9):4413-4416.
257. Lv H, Han T, Shi X, et al. Genetic polymorphism of GSTP1 and ERCC1 correlated with response to platinum-based chemotherapy in non-small cell lung cancer. *Med Oncol.* 2014;31(8):86-014-0086-5. Epub 2014 Jun 24.
258. Yang Y and Xian L. The association between the GSTP1 A313G and GSTM1 null/present polymorphisms and the treatment response of the platinum-based chemotherapy in non-small cell lung cancer (NSCLC) patients: a meta-analysis. *Tumour Biol.* 2014;35(7):6791-6799.
259. Yin JY, Huang Q, Zhao YC, Zhou HH, Liu ZQ. Meta-analysis on pharmacogenetics of platinum-based chemotherapy in non small cell lung cancer (NSCLC) patients. *PLoS One.* 2012;7(6):e38150.
260. Zhou F, Yu Z, Jiang T, Lv H, Yao R, Liang J. Genetic polymorphisms of GSTP1 and XRCC1: prediction of clinical outcome of platinum-based chemotherapy in advanced non-small cell lung cancer (NSCLC) patients. *Swiss Med Wkly.* 2011;141:w13275.
261. Yang P, Yokomizo A, Tazelaar HD, et al. Genetic determinants of lung cancer short-term survival: the role of glutathione-related genes. *Lung Cancer.* 2002;35(3):221-229.
262. Saip R, Sen F, Vural B, et al. Glutathione S-transferase P1 polymorphisms are associated with time to tumor progression in small cell lung cancer patients. *J BUON.* 2011;16(2):241-246.
263. Booton R, Ward T, Heighway J, et al. Xeroderma pigmentosum group D haplotype predicts for response, survival, and toxicity after platinum-based chemotherapy in advanced nonsmall cell lung cancer. *Cancer.* 2006;106(11):2421-2427.
264. Booton R, Ward T, Heighway J, Ashcroft L, Morris J, Thatcher N. Glutathione-S-transferase P1 isoenzyme polymorphisms, platinum-based chemotherapy, and non-small cell lung cancer. *J Thorac Oncol.* 2006;1(7):679-683.
265. Khrunin AV, Moisseev A, Gorbunova V, Limborska S. Genetic polymorphisms and the efficacy and toxicity of cisplatin-based chemotherapy in ovarian cancer patients. *Pharmacogenomics J.* 2010;10(1):54-61.
266. Khrunin A, Ivanova F, Moisseev A, et al. Pharmacogenomics of cisplatin-based chemotherapy in ovarian cancer patients of different ethnic origins. *Pharmacogenomics.* 2012;13(2):171-178.
267. Kim KH, Do IG, Kim HS, et al. Excision repair cross-complementation group 1 (ERCC1) expression in advanced urothelial carcinoma patients receiving cisplatin-based chemotherapy. *APMIS.* 2010;118(12):941-948.

268. Kolwijck E, Zusterzeel PL, Roelofs HM, Hendriks JC, Peters WH, Massuger LF. GSTP1-1 in ovarian cyst fluid and disease outcome of patients with ovarian cancer. *Cancer Epidemiol Biomarkers Prev.* 2009;18(8):2176-2181.
269. Kraggerud SM, Oldenburg J, Alnaes GI, et al. Functional glutathione S-transferase genotypes among testicular germ cell tumor survivors: associations with primary and post-chemotherapy tumor histology. *Pharmacogenet Genomics.* 2009;19(10):751-759.
270. Xu Z, Chen Y, Gu D, et al. SOD2 rs4880 CT/CC genotype predicts poor survival for Chinese gastric cancer patients received platinum and fluorouracil based adjuvant chemotherapy. *Am J Transl Res.* 2015;7(2):401-410.
271. Wang Z, Chen JQ, Liu JL, Qin XG, Huang Y. Polymorphisms in ERCC1, GSTs, TS and MTHFR predict clinical outcomes of gastric cancer patients treated with platinum/5-Fu-based chemotherapy: a systematic review. *BMC Gastroenterol.* 2012;12:137-230X-12-137.
272. Boku N, Chin K, Hosokawa K, et al. Biological markers as a predictor for response and prognosis of unresectable gastric cancer patients treated with 5-fluorouracil and cis-platinum. *Clin Cancer Res.* 1998;4(6):1469-1474.
273. Li HY, Ge X, Huang GM, et al. GSTP1, ERCC1 and ERCC2 polymorphisms, expression and clinical outcome of oxaliplatin-based adjuvant chemotherapy in colorectal cancer in Chinese population. *Asian Pac J Cancer Prev.* 2012;13(7):3465-3469.
274. Chen YC, Tzeng CH, Chen PM, et al. Influence of GSTP1 I105V polymorphism on cumulative neuropathy and outcome of FOLFOX-4 treatment in Asian patients with colorectal carcinoma. *Cancer Sci.* 2010;101(2):530-535.
275. Yoon HH, Catalano P, Gibson MK, et al. Genetic variation in radiation and platinum pathways predicts severe acute radiation toxicity in patients with esophageal adenocarcinoma treated with cisplatin-based preoperative radiochemotherapy: results from the Eastern Cooperative Oncology Group. *Cancer Chemother Pharmacol.* 2011;68(4):863-870.
276. Lecomte T, Landi B, Beaune P, Laurent-Puig P, Lloria MA. Glutathione S-transferase P1 polymorphism (Ile105Val) predicts cumulative neuropathy in patients receiving oxaliplatin-based chemotherapy. *Clin Cancer Res.* 2006;12(10):3050-3056.
277. Lu C, Spitz MR, Zhao H, et al. Association between glutathione S-transferase pi polymorphisms and survival in patients with advanced nonsmall cell lung carcinoma. *Cancer.* 2006;106(2):441-447.
278. Pereira D, Assis J, Gomes M, Nogueira A, Medeiros R. Improvement of a predictive model in ovarian cancer patients submitted to platinum-based chemotherapy: implications of a GST activity profile. *Eur J Clin Pharmacol.* 2016;72(5):545-553.
279. Medeiros R, Pereira D, Afonso N, et al. Platinum/paclitaxel-based chemotherapy in advanced ovarian carcinoma: glutathione S-transferase genetic polymorphisms as predictive biomarkers of disease outcome. *Int J Clin Oncol.* 2003;8(3):156-161.
280. Li W, Yue W, Zhang L, et al. Polymorphisms in GSTM1, CYP1A1, CYP2E1, and CYP2D6 are associated with susceptibility and chemotherapy response in non-small-cell lung cancer patients. *Lung.* 2012;190(1):91-98.
281. Beeghly A, Katsaros D, Chen H, et al. Glutathione S-transferase polymorphisms and ovarian cancer treatment and survival. *Gynecol Oncol.* 2006;100(2):330-337.
282. Ekhardt C, Rodenhuis S, Smits PH, Beijnen JH, Huitema AD. An overview of the relations between polymorphisms in drug metabolising enzymes and drug transporters and survival after cancer drug treatment. *Cancer Treat Rev.* 2009;35(1):18-31.
283. Peters U, Preisler-Adams S, Hebeisen A, et al. Glutathione S-transferase genetic polymorphisms and individual sensitivity to the ototoxic effect of cisplatin. *Anticancer Drugs.* 2000;11(8):639-643.
284. Brown AL, Lupo PJ, Okcu MF, Lau CC, Rednam S, Scheurer ME. SOD2 genetic variant associated with treatment-related ototoxicity in cisplatin-treated pediatric medulloblastoma. *Cancer Med.* 2015;4(11):1679-1686.
285. Bag A, Pant NK, Jeena LM, Bag N, Jyala NS. GSTT1 null and MPO -463G>a polymorphisms and carboplatin toxicity in an Indian population. *Asian Pac J Cancer Prev.* 2013;14(8):4739-4742.
286. Tian G, Wang M, Xu X. The role of NQO1 polymorphisms in the susceptibility and chemotherapy response of Chinese NSCLC patients. *Cell Biochem Biophys.* 2014;69(3):475-479.
287. Paek J, Lee M, Nam EJ, Kim SW, Kim YT. Clinical impact of high mobility group box 1 protein in epithelial ovarian cancer. *Arch Gynecol Obstet.* 2016;293(3):645-650.

288. Wang Y, Li XP, Yin JY, et al. Association of HMGB1 and HMGB2 genetic polymorphisms with lung cancer chemotherapy response. *Clin Exp Pharmacol Physiol*. 2014;41(6):408-415.
289. Zhao W, Hu L, Xu J, et al. Polymorphisms in the base excision repair pathway modulate prognosis of platinum-based chemotherapy in advanced non-small cell lung cancer. *Cancer Chemother Pharmacol*. 2013;71(5):1287-1295.
290. Velasco A, Corvalan A, Wistuba II, et al. Mismatch repair expression in testicular cancer predicts recurrence and survival. *Int J Cancer*. 2008;122(8):1774-1777.
291. Vaisman A, Varchenko M, Umar A, et al. The role of hMLH1, hMSH3, and hMSH6 defects in cisplatin and oxaliplatin resistance: correlation with replicative bypass of platinum-DNA adducts. *Cancer Res*. 1998;58(16):3579-3585.
292. Ting S, Mairinger FD, Hager T, et al. ERCC1, MLH1, MSH2, MSH6, and betaIII-tubulin: resistance proteins associated with response and outcome to platinum-based chemotherapy in malignant pleural mesothelioma. *Clin Lung Cancer*. 2013;14(5):558-567.e3.
293. Xu XL, Yao YL, Xu WZ, Feng JG, Mao WM. Correlation of MSH3 polymorphisms with response and survival in advanced non-small cell lung cancer patients treated with first-line platinum-based chemotherapy. *Genet Mol Res*. 2015;14(2):3525-3533.
294. Samimi G, Fink D, Varki NM, et al. Analysis of MLH1 and MSH2 expression in ovarian cancer before and after platinum drug-based chemotherapy. *Clin Cancer Res*. 2000;6(4):1415-1421.
295. Scartozzi M, De Nicolis M, Galizia E, et al. Loss of hMLH1 expression correlates with improved survival in stage III-IV ovarian cancer patients. *Eur J Cancer*. 2003;39(8):1144-1149.
296. Rossi D, Rasi S, Di Rocco A, et al. The host genetic background of DNA repair mechanisms is an independent predictor of survival in diffuse large B-cell lymphoma. *Blood*. 2011;117(8):2405-2413.
297. Jia J, Wang Z, Cai J, Zhang Y. PMS2 expression in epithelial ovarian cancer is posttranslationally regulated by Akt and essential for platinum-induced apoptosis. *Tumour Biol*. 2016;37(3):3059-3069.
298. Fu X, Hu J, Han HY, et al. High expression of XPA confers poor prognosis for nasopharyngeal carcinoma patients treated with platinum-based chemoradiotherapy. *Oncotarget*. 2015;6(29):28478-28490.
299. Dabholkar M, Bostick-Bruton F, Weber C, Bohr VA, Egwuagu C, Reed E. ERCC1 and ERCC2 expression in malignant tissues from ovarian cancer patients. *J Natl Cancer Inst*. 1992;84(19):1512-1517.
300. Dabholkar M, Vionnet J, Bostick-Bruton F, Yu JJ, Reed E. Messenger RNA levels of XPAC and ERCC1 in ovarian cancer tissue correlate with response to platinum-based chemotherapy. *J Clin Invest*. 1994;94(2):703-708.
301. Stevens EV, Raffeld M, Espina V, et al. Expression of xeroderma pigmentosum A protein predicts improved outcome in metastatic ovarian carcinoma. *Cancer*. 2005;103(11):2313-2319.
302. Ganzinelli M, Mariani P, Cattaneo D, et al. Expression of DNA repair genes in ovarian cancer samples: biological and clinical considerations. *Eur J Cancer*. 2011;47(7):1086-1094.
303. Cheng H, Qin Q, Sun X, et al. Predictive effect of XPA and XPD polymorphisms on survival of advanced NSCLC patients treated with platinum-based chemotherapy: a three-dimensional (3-D), polyacrylamide gel-based DNA microarray method. *Technol Cancer Res Treat*. 2013;12(5):473-482.
304. Feng J, Sun X, Sun N, et al. XPA A23G polymorphism is associated with the elevated response to platinum-based chemotherapy in advanced non-small cell lung cancer. *Acta Biochim Biophys Sin (Shanghai)*. 2009;41(5):429-435.
305. Butkiewicz D, Drosik A, Suwinski R, et al. Influence of DNA repair gene polymorphisms on prognosis in inoperable non-small cell lung cancer patients treated with radiotherapy and platinum-based chemotherapy. *Int J Cancer*. 2012;131(7):E1100-8.
306. Sullivan I, Salazar J, Majem M, et al. Pharmacogenetics of the DNA repair pathways in advanced non-small cell lung cancer patients treated with platinum-based chemotherapy. *Cancer Lett*. 2014;353(2):160-166.
307. Saldivar JS, Lu KH, Liang D, et al. Moving toward individualized therapy based on NER polymorphisms that predict platinum sensitivity in ovarian cancer patients. *Gynecol Oncol*. 2007;107(1 Suppl 1):S223-9.
308. Powrozek T, Mlak R, Krawczyk P, et al. The relationship between polymorphisms of genes regulating DNA repair or cell division and the toxicity of platinum and vinorelbine chemotherapy in advanced NSCLC patients. *Clin Transl Oncol*. 2016;18(2):125-131.
309. Wyss AB, Weissler MC, Avery CL, et al. Single nucleotide polymorphisms in nucleotide excision repair genes, cancer treatment, and head and neck cancer survival. *Cancer Causes Control*. 2014;25(4):437-450.

310. Kang S, Sun HY, Zhou RM, Wang N, Hu P, Li Y. DNA repair gene associated with clinical outcome of epithelial ovarian cancer treated with platinum-based chemotherapy. *Asian Pac J Cancer Prev.* 2013;14(2):941-946.
311. Zhu XL, Sun XC, Chen BA, et al. XPC Lys939Gln polymorphism is associated with the decreased response to platinum based chemotherapy in advanced non-small-cell lung cancer. *Chin Med J (Engl).* 2010;123(23):3427-3432.
312. Caronia D, Patino-Garcia A, Milne RL, et al. Common variations in ERCC2 are associated with response to cisplatin chemotherapy and clinical outcome in osteosarcoma patients. *Pharmacogenomics J.* 2009;9(5):347-353.
313. Sakano S, Hinoda Y, Sasaki M, et al. Nucleotide excision repair gene polymorphisms may predict acute toxicity in patients treated with chemoradiotherapy for bladder cancer. *Pharmacogenomics.* 2010;11(10):1377-1387.
314. Fleming ND, Agadjanian H, Nassanian H, et al. Xeroderma pigmentosum complementation group C single-nucleotide polymorphisms in the nucleotide excision repair pathway correlate with prolonged progression-free survival in advanced ovarian cancer. *Cancer.* 2012;118(3):689-697.
315. Dong J, Hu Z, Shu Y, et al. Potentially functional polymorphisms in DNA repair genes and non-small-cell lung cancer survival: a pathway-based analysis. *Mol Carcinog.* 2012;51(7):546-552.
316. Li P, Wang YD, Cheng J, Chen JC, Ha MW. Association between polymorphisms of BAG-1 and XPD and chemotherapy sensitivity in advanced non-small-cell lung cancer patients treated with vinorelbine combined cisplatin regimen. *Tumour Biol.* 2015;36(12):9465-9473.
317. Gurubhagavatula S, Liu G, Park S, et al. XPD and XRCC1 genetic polymorphisms are prognostic factors in advanced non-small-cell lung cancer patients treated with platinum chemotherapy. *J Clin Oncol.* 2004;22(13):2594-2601.
318. Qiu M, Yang X, Hu J, et al. Predictive value of XPD polymorphisms on platinum-based chemotherapy in non-small cell lung cancer: a systematic review and meta-analysis. *PLoS One.* 2013;8(8):e72251.
319. Yuan P, Miao XP, Zhang XM, et al. Correlation of genetic polymorphisms in nucleotide excision repair system to sensitivity of advanced non-small cell lung cancer patients to platinum-based chemotherapy. *Ai Zheng.* 2005;24(12):1510-1513.
320. Zhou M, Ding YJ, Feng Y, Zhang QR, Xiang Y, Wan HY. Association of xeroderma pigmentosum group D (Asp312Asn, Lys751Gln) and cytidine deaminase (Lys27Gln, Ala70Thr) polymorphisms with outcome in Chinese non-small cell lung cancer patients treated with cisplatin-gemcitabine. *Genet Mol Res.* 2014;13(2):3310-3318.
321. Geredeli C, Artac M, Yildirim S, et al. Prognostic value of ERCC1, ERCC2, XRCC1, and TP53 single nucleotide polymorphisms in patients with early-stage non-small cell lung cancer. *Tumour Biol.* 2015;36(6):4279-4285.
322. Chen X, Sun H, Ren S, et al. Association of XRCC3 and XPD751 SNP with efficacy of platinum-based chemotherapy in advanced NSCLC patients. *Clin Transl Oncol.* 2012;14(3):207-213.
323. Li XD, Han JC, Zhang YJ, Li HB, Wu XY. Common variations of DNA repair genes are associated with response to platinum-based chemotherapy in NSCLCs. *Asian Pac J Cancer Prev.* 2013;14(1):145-148.
324. Zhang L, Ma W, Li Y, Wu J, Shi GY. Pharmacogenetics of DNA repair gene polymorphisms in non-small-cell lung carcinoma patients on platinum-based chemotherapy. *Genet Mol Res.* 2014;13(1):228-236.
325. Qin Q, Zhang C, Yang X, et al. Polymorphisms in XPD gene could predict clinical outcome of platinum-based chemotherapy for non-small cell lung cancer patients: a meta-analysis of 24 studies. *PLoS One.* 2013;8(11):e79864.
326. Bewick MA, Lafrenie RM, Conlon MS. Nucleotide excision repair polymorphisms and survival outcome for patients with metastatic breast cancer. *J Cancer Res Clin Oncol.* 2011;137(3):543-550.
327. Goekkurt E, Al-Batran SE, Hartmann JT, et al. Pharmacogenetic analyses of a phase III trial in metastatic gastroesophageal adenocarcinoma with fluorouracil and leucovorin plus either oxaliplatin or cisplatin: a study of the arbeitgemeinschaft internistische onkologie. *J Clin Oncol.* 2009;27(17):2863-2873.
328. Park DJ, Stoehlmacher J, Zhang W, Tsao-Wei DD, Groshen S, Lenz HJ. A Xeroderma pigmentosum group D gene polymorphism predicts clinical outcome to platinum-based chemotherapy in patients with advanced colorectal cancer. *Cancer Res.* 2001;61(24):8654-8658.
329. Dong Y, Liu JW, Gao YJ, Zhou T, Chen YM. Relationship between DNA repair gene XPD751 single-nucleotide polymorphisms and prognosis of colorectal cancer. *Genet Mol Res.* 2015;14(2):5390-5398.

330. Yin M, Yan J, Martinez-Balibrea E, et al. ERCC1 and ERCC2 polymorphisms predict clinical outcomes of oxaliplatin-based chemotherapies in gastric and colorectal cancer: a systemic review and meta-analysis. *Clin Cancer Res*. 2011;17(6):1632-1640.
331. Sakano S, Wada T, Matsumoto H, et al. Single nucleotide polymorphisms in DNA repair genes might be prognostic factors in muscle-invasive bladder cancer patients treated with chemoradiotherapy. *Br J Cancer*. 2006;95(5):561-570.
332. Ren S, Zhou S, Wu F, et al. Association between polymorphisms of DNA repair genes and survival of advanced NSCLC patients treated with platinum-based chemotherapy. *Lung Cancer*. 2012;75(1):102-109.
333. Biason P, Hattinger CM, Innocenti F, et al. Nucleotide excision repair gene variants and association with survival in osteosarcoma patients treated with neoadjuvant chemotherapy. *Pharmacogenomics J*. 2012;12(6):476-483.
334. Chen YM, Wu XL, Zhang LW, Xu X, Liu JW. Relationship between single nucleotide polymorphism in repair gene XPD751 and prognosis in colorectal carcinoma patients. *Zhonghua Zhong Liu Za Zhi*. 2012;34(7):501-505.
335. Le Morvan V, Smith D, Laurand A, et al. Determination of ERCC2 Lys751Gln and GSTP1 Ile105Val gene polymorphisms in colorectal cancer patients: relationships with treatment outcome. *Pharmacogenomics*. 2007;8(12):1693-1703.
336. Pare L, Marcuello E, Altes A, et al. Pharmacogenetic prediction of clinical outcome in advanced colorectal cancer patients receiving oxaliplatin/5-fluorouracil as first-line chemotherapy. *Br J Cancer*. 2008;99(7):1050-1055.
337. Lai JI, Tzeng CH, Chen PM, et al. Very low prevalence of XPD K751Q polymorphism and its association with XPD expression and outcomes of FOLFOX-4 treatment in Asian patients with colorectal carcinoma. *Cancer Sci*. 2009;100(7):1261-1266.
338. Ding C, Zhang H, Chen K, Zhao C, Gao J. Genetic variability of DNA repair mechanisms influences treatment outcome of gastric cancer. *Oncol Lett*. 2015;10(4):1997-2002.
339. Zhou J, Liu ZY, Li CB, et al. Genetic polymorphisms of DNA repair pathways influence the response to chemotherapy and overall survival of gastric cancer. *Tumour Biol*. 2015;36(4):3017-3023.
340. Wu W, Zhang W, Qiao R, et al. Association of XPD polymorphisms with severe toxicity in non-small cell lung cancer patients in a Chinese population. *Clin Cancer Res*. 2009;15(11):3889-3895.
341. Wu W, Li H, Wang H, et al. Effect of polymorphisms in XPD on clinical outcomes of platinum-based chemotherapy for Chinese non-small cell lung cancer patients. *PLoS One*. 2012;7(3):e33200.
342. Zhou F, Zhu M, Wang M, et al. Genetic variants of DNA repair genes predict the survival of patients with esophageal squamous cell cancer receiving platinum-based adjuvant chemotherapy. *J Transl Med*. 2016;14(1):154-016-0903-z.
343. Seiwert TY, Wang X, Heitmann J, et al. DNA repair biomarkers XPF and phospho-MAPKAP kinase 2 correlate with clinical outcome in advanced head and neck cancer. *PLoS One*. 2014;9(7):e102112.
344. Ceccaldi R, O'Connor KW, Mouw KW, et al. A unique subset of epithelial ovarian cancers with platinum sensitivity and PARP inhibitor resistance. *Cancer Res*. 2015;75(4):628-634.
345. Liu D, Wu J, Shi GY, Zhou HF, Yu Y. Role of XRCC1 and ERCC5 polymorphisms on clinical outcomes in advanced non-small cell lung cancer. *Genet Mol Res*. 2014;13(2):3100-3107.
346. Jin ZY, Zhao XT, Zhang LN, Wang Y, Yue WT, Xu SF. Effects of polymorphisms in the XRCC1, XRCC3, and XPG genes on clinical outcomes of platinum-based chemotherapy for treatment of non-small cell lung cancer. *Genet Mol Res*. 2014;13(3):7617-7625.
347. He C, Duan Z, Li P, Xu Q, Yuan Y. Role of ERCC5 promoter polymorphisms in response to platinum-based chemotherapy in patients with advanced non-small-cell lung cancer. *Anticancer Drugs*. 2013;24(3):300-305.
348. Yuli Y, Zhe S, Xia W, et al. XPG is a novel biomarker of clinical outcome in advanced non-small-cell lung cancer. *Pak J Med Sci*. 2013;29(3):762-767.
349. Zhang L, Gao G, Li X, et al. Association between single nucleotide polymorphisms (SNPs) and toxicity of advanced non-small-cell lung cancer patients treated with chemotherapy. *PLoS One*. 2012;7(10):e48350.
350. Matakidou A, el Galta R, Webb EL, et al. Genetic variation in the DNA repair genes is predictive of outcome in lung cancer. *Hum Mol Genet*. 2007;16(19):2333-2340.
351. Caiola E, Porcu L, Fruscio R, et al. DNA-damage response gene polymorphisms and therapeutic outcomes in ovarian cancer. *Pharmacogenomics J*. 2013;13(2):159-172.
352. Somers J, Wilson LA, Kilday JP, et al. A common polymorphism in the 5' UTR of ERCC5 creates an upstream ORF that confers resistance to platinum-based chemotherapy. *Genes Dev*. 2015;29(18):1891-1896.

353. Hu W, Pan J, Zhao P, Yang G, Yang S. Genetic polymorphisms in XPG could predict clinical outcome of platinum-based chemotherapy for advanced non-small cell lung cancer. *Tumour Biol.* 2014;35(6):5561-5567.
354. Zhang T, Sun J, Lv M, et al. XPG is predictive gene of clinical outcome in advanced non-small-cell lung cancer with platinum drug therapy. *Asian Pac J Cancer Prev.* 2013;14(2):701-705.
355. Amable L. Cisplatin resistance and opportunities for precision medicine. *Pharmacol Res.* 2016;106:27-36.
356. Li W, Sun Q, Lu M. Correlation between expression of ERCC1 and the treatment of cisplatin-based chemotherapy in local advanced nasopharyngeal carcinoma. *Lin Chung Er Bi Yan Hou Tou Jing Wai Ke Za Zhi.* 2015;29(2):144-146.
357. Song P, Yin Q, Lu M, Fu BO, Wang B, Zhao Q. Prognostic value of excision repair cross-complementation group 1 expression in gastric cancer: A meta-analysis. *Exp Ther Med.* 2015;9(4):1393-1400.
358. Miura JT, Xiu J, Thomas J, et al. Tumor profiling of gastric and esophageal carcinoma reveal different treatment options. *Cancer Biol Ther.* 2015;16(5):764-769.
359. Sakano S, Ogawa S, Yamamoto Y, et al. ERCC1 and XRCC1 expression predicts survival in bladder cancer patients receiving combined trimodality therapy. *Mol Clin Oncol.* 2013;1(3):403-410.
360. Metzger R, Bollschweiler E, Holscher AH, Warnecke-Eberz U. ERCC1: impact in multimodality treatment of upper gastrointestinal cancer. *Future Oncol.* 2010;6(11):1735-1749.
361. Bussey KJ and Demeure MJ. Toward a pathway-centered approach for the treatment of adrenocortical carcinoma. *Curr Opin Oncol.* 2011;23(1):34-44.
362. Vergote I, Calvert H, Kania M, Kaiser C, Zimmermann AH, Sehouli J. A randomised, double-blind, phase II study of two doses of pemetrexed in the treatment of platinum-resistant, epithelial ovarian or primary peritoneal cancer. *Eur J Cancer.* 2009;45(8):1415-1423.
363. Felip E and Rosell R. Testing for excision repair cross-complementing 1 in patients with non-small-cell lung cancer for chemotherapy response. *Expert Rev Mol Diagn.* 2007;7(3):261-268.
364. Altaha R, Liang X, Yu JJ, Reed E. Excision repair cross complementing-group 1: gene expression and platinum resistance. *Int J Mol Med.* 2004;14(6):959-970.
365. Rosell R and Felip E. Predicting response to paclitaxel/carboplatin-based therapy in non-small cell lung cancer. *Semin Oncol.* 2001;28(4 Suppl 14):37-44.
366. Steffensen KD, Waldstrom M, Jeppesen U, Brandslund I, Jakobsen A. Prediction of response to chemotherapy by ERCC1 immunohistochemistry and ERCC1 polymorphism in ovarian cancer. *Int J Gynecol Cancer.* 2008;18(4):702-710.
367. Ameri A, Mortazavi N, Khoshbakht Ahmadi H, Novin K. ERCC1 Expression Can Predict Response to Platinum-Based Induction Chemotherapy in Head and Neck Cancer Cases. *Asian Pac J Cancer Prev.* 2016;17 Spec No.:87-91.
368. Bogush TA, Popova AS, Dudko EA, et al. ERCC1 as a Marker of Ovarian Cancer Resistance to Platinum Drugs. *Antibiot Khimioter.* 2015;60(3-4):42-50.
369. Li FY, Ren XB, Xie XY, Zhang J. Meta-analysis of excision repair cross-complementation group 1 (ERCC1) association with response to platinum- based chemotherapy in ovarian cancer. *Asian Pac J Cancer Prev.* 2013;14(12):7203-7206.
370. Muallem MZ, Braicu I, Nassir M, Richter R, Sehouli J, Arsenic R. ERCC1 expression as a predictor of resistance to platinum-based chemotherapy in primary ovarian cancer. *Anticancer Res.* 2014;34(1):393-399.
371. Allingham-Hawkins D, Lea A, Levine S. ERCC1 Expression Analysis to Guide Therapy in Non-Small Cell Lung Cancer. *PLoS Curr.* 2010;2:RRN1202.
372. Scheil-Bertram S, Tylus-Schaaf P, du Bois A, et al. Excision repair cross-complementation group 1 protein overexpression as a predictor of poor survival for high-grade serous ovarian adenocarcinoma. *Gynecol Oncol.* 2010;119(2):325-331.
373. Yu J, Dabholkar M, Bennett W, et al. Platinum-sensitive and platinum-resistant ovarian cancer tissues show differences in the relationships between mRNA levels of p53, ERCC1 and XPA. *Int J Oncol.* 1996;8(2):313-317.
374. Milovic-Kovacevic M, Srdic-Rajic T, Radulovic S, Bjelogric S, Gavrilovic D. Expression of ERCC1 protein in biopsy specimen predicts survival in advanced ovarian cancer patients treated with platinum-based chemotherapy. *J BUON.* 2011;16(4):708-714.
375. Weberpals J, Garbuio K, O'Brien A, et al. The DNA repair proteins BRCA1 and ERCC1 as predictive markers in sporadic ovarian cancer. *Int J Cancer.* 2009;124(4):806-815.

376. Xie C, Yin RT, Li YL, Kang DY, Xu L, Yang KX. The protein expression of ERCC1 and survivin in epithelial ovarian carcinoma and their clinical significance. *Sichuan Da Xue Xue Bao Yi Xue Ban*. 2011;42(1):86-89.
377. Steffensen KD, Smoter M, Waldstrom M, et al. Resistance to first line platinum paclitaxel chemotherapy in serous epithelial ovarian cancer: the prediction value of ERCC1 and Tau expression. *Int J Oncol*. 2014;44(5):1736-1744.
378. Steffensen KD, Waldstrom M, Jakobsen A. The relationship of platinum resistance and ERCC1 protein expression in epithelial ovarian cancer. *Int J Gynecol Cancer*. 2009;19(5):820-825.
379. Park JS, Jeon EK, Chun SH, et al. ERCC1 (excision repair cross-complementation group 1) expression as a predictor for response of neoadjuvant chemotherapy for FIGO stage 2B uterine cervix cancer. *Gynecol Oncol*. 2011;120(2):275-279.
380. Yao A, Wang Y, Peng X, et al. Predictive value of excision repair cross-complementation group 1 expression for platinum-based chemotherapy and survival in gastric cancer: a meta-analysis. *J Cancer Res Clin Oncol*. 2014;140(12):2107-2117.
381. Wei KK, Jiang L, Wei YY, et al. The prognostic value of ERCC1 expression in gastric cancer patients treated with platinum-based chemotherapy: a meta-analysis. *Tumour Biol*. 2014;35(9):8721-8731.
382. Qi YJ, Cui S, Yang YZ, et al. Excision repair cross-complementation group 1 codon 118 polymorphism, micro ribonucleic acid and protein expression, clinical outcome of the advanced gastric cancer response to first-line FOLFOX-4 in Qinghai-Tibetan plateau population. *J Cancer Res Ther*. 2013;9(3):410-415.
383. Liu YP, Ling Y, Qi QF, et al. The effects of ERCC1 expression levels on the chemosensitivity of gastric cancer cells to platinum agents and survival in gastric cancer patients treated with oxaliplatin-based adjuvant chemotherapy. *Oncol Lett*. 2013;5(3):935-942.
384. Ozkan M, Akbudak IH, Deniz K, et al. Prognostic value of excision repair cross-complementing gene 1 expression for cisplatin-based chemotherapy in advanced gastric cancer. *Asian Pac J Cancer Prev*. 2010;11(1):181-185.
385. Bamias A, Karina M, Papakostas P, et al. A randomized phase III study of adjuvant platinum/docetaxel chemotherapy with or without radiation therapy in patients with gastric cancer. *Cancer Chemother Pharmacol*. 2010;65(6):1009-1021.
386. Lee HW, Choi YW, Han JH, et al. Expression of excision repair cross-complementation group 1 protein predicts poor outcome in advanced non-small cell lung cancer patients treated with platinum-based doublet chemotherapy. *Lung Cancer*. 2009;65(3):377-382.
387. Azuma K, Sasada T, Kawahara A, et al. Expression of ERCC1 and class III beta-tubulin in non-small cell lung cancer patients treated with carboplatin and paclitaxel. *Lung Cancer*. 2009;64(3):326-333.
388. Wei J, Zou Z, Qian X, et al. ERCC1 mRNA levels and survival of advanced gastric cancer patients treated with a modified FOLFOX regimen. *Br J Cancer*. 2008;98(8):1398-1402.
389. Bonanno L. Predictive models for customizing chemotherapy in advanced non-small cell lung cancer (NSCLC). *Transl Lung Cancer Res*. 2013;2(3):160-171.
390. Qiao H, Huang X, Guo H, Liu Y, Yue C. ERCC1, RRM1 and TUBB3 mRNA expression on the tumor response and overall survival of non-small cell lung cancer treated with platinum-based chemotherapy. *Pak J Med Sci*. 2014;30(6):1403-1408.
391. Huang ZL, Cao X, Luo RZ, Chen YF, Zhu LC, Wen Z. Analysis of ERCC1, BRCA1, RRM1 and TUBB3 as predictors of prognosis in patients with non-small cell lung cancer who received cisplatin-based adjuvant chemotherapy: A prospective study. *Oncol Lett*. 2016;11(1):299-305.
392. Deng Q, Yang H, Lin Y, et al. Prognostic value of ERCC1 mRNA expression in non-small cell lung cancer, breast cancer, and gastric cancer in patients from Southern China. *Int J Clin Exp Pathol*. 2014;7(12):8312-8321.
393. Karim NA, Bui H, Pathrose P, et al. The use of pharmacogenomics for selection of therapy in non-small-cell lung cancer. *Clin Med Insights Oncol*. 2014;8:139-144.
394. Zhang H, Li J, Zhang Y, et al. ERCC1 mRNA expression is associated with the clinical outcome of non-small cell lung cancer treated with platinum-based chemotherapy. *Genet Mol Res*. 2014;13(4):10215-10222.
395. Xian-Jun F, Xiu-Guang Q, Li Z, et al. ERCC1 and BRCA1 mRNA expression predicts the clinical outcome of non-small cell lung cancer receiving platinum-based chemotherapy. *Pak J Med Sci*. 2014;30(3):488-492.
396. Vassalou H, Stathopoulos E, Fiolitaki G, et al. Excision-repair-cross-complement-1 protein as a prognostic factor in patients with advanced non-small cell lung cancer treated with platinum-based first-line chemotherapy. *Lung Cancer*. 2013;82(2):324-329.

397. Tiseo M, Bordi P, Bortesi B, et al. ERCC1/BRCA1 expression and gene polymorphisms as prognostic and predictive factors in advanced NSCLC treated with or without cisplatin. *Br J Cancer*. 2013;108(8):1695-1703.
398. Sun S, Shi W, Wu Z, Zhang G, Yang BO, Jiao S. Prognostic significance of the mRNA expression of ERCC1, RRM1, TUBB3 and TYMS genes in patients with non-small cell lung cancer. *Exp Ther Med*. 2015;10(3):937-941.
399. Jiang H, Wang H, Wang S, et al. Expression of ERCC1, TYMS, RRM1, TUBB3, non-muscle myosin II, myoglobin and MyoD1 in lung adenocarcinoma pleural effusions predicts survival in patients receiving platinum-based chemotherapy. *Mol Med Rep*. 2015;11(5):3523-3532.
400. Wang TB, Zhang NL, Wang SH, Li HY, Chen SW, Zheng YG. Expression of ERCC1 and BRCA1 predict the clinical outcome of non-small cell lung cancer in patients receiving platinum-based chemotherapy. *Genet Mol Res*. 2014;13(2):3704-3710.
401. Qin X, Yao W, Li W, et al. ERCC1 and BRCA1 mRNA expressions are associated with clinical outcome of non-small cell lung cancer treated with platinum-based chemotherapy. *Tumour Biol*. 2014;35(5):4697-4704.
402. Das M, Riess JW, Frankel P, et al. ERCC1 expression in circulating tumor cells (CTCs) using a novel detection platform correlates with progression-free survival (PFS) in patients with metastatic non-small-cell lung cancer (NSCLC) receiving platinum chemotherapy. *Lung Cancer*. 2012;77(2):421-426.
403. Leng XF, Chen MW, Xian L, Dai L, Ma GY, Li MH. Combined analysis of mRNA expression of ERCC1, BAG-1, BRCA1, RRM1 and TUBB3 to predict prognosis in patients with non-small cell lung cancer who received adjuvant chemotherapy. *J Exp Clin Cancer Res*. 2012;31:25-9966-31-25.
404. Papadaki C, Sfakianaki M, Ioannidis G, et al. ERCC1 and BRCA1 mRNA expression levels in the primary tumor could predict the effectiveness of the second-line cisplatin-based chemotherapy in pretreated patients with metastatic non-small cell lung cancer. *J Thorac Oncol*. 2012;7(4):663-671.
405. Jiang J, Liang X, Zhou X, Huang R, Chu Z, Zhan Q. ERCC1 expression as a prognostic and predictive factor in patients with non-small cell lung cancer: a meta-analysis. *Mol Biol Rep*. 2012;39(6):6933-6942.
406. Roth JA and Carlson JJ. Prognostic role of ERCC1 in advanced non-small-cell lung cancer: a systematic review and meta-analysis. *Clin Lung Cancer*. 2011;12(6):393-401.
407. Joerger M, deJong D, Burylo A, et al. Tubulin, BRCA1, ERCC1, Abraxas, RAP80 mRNA expression, p53/p21 immunohistochemistry and clinical outcome in patients with advanced non small-cell lung cancer receiving first-line platinum-gemcitabine chemotherapy. *Lung Cancer*. 2011;74(2):310-317.
408. Zhang S, Li Q, Zhang Q, et al. Expression of ERCC1 and class III α -tubulin in resected non-small cell lung cancer and its correlation with platinum-based adjuvant chemotherapy. *Int J Biol Markers*. 2010;25(3):141-149.
409. Chen S, Zhang J, Wang R, Luo X, Chen H. The platinum-based treatments for advanced non-small cell lung cancer, is low/negative ERCC1 expression better than high/positive ERCC1 expression? A meta-analysis. *Lung Cancer*. 2010;70(1):63-70.
410. Ren S, Zhou S, Zhang L, et al. High-level mRNA of excision repair cross-complementation group 1 gene is associated with poor outcome of platinum-based doublet chemotherapy of advanced nonsmall cell lung cancer patients. *Cancer Invest*. 2010;28(10):1078-1083.
411. Su C, Zhou S, Zhang L, et al. ERCC1, RRM1 and BRCA1 mRNA expression levels and clinical outcome of advanced non-small cell lung cancer. *Med Oncol*. 2011;28(4):1411-1417.
412. Wang X, Zhao J, Yang L, et al. Positive expression of ERCC1 predicts a poorer platinum-based treatment outcome in Chinese patients with advanced non-small-cell lung cancer. *Med Oncol*. 2010;27(2):484-490.
413. Schettino C, Bareschino MA, Maione P, Rossi A, Ciardiello F, Gridelli C. The potential role of pharmacogenomic and genomic in the adjuvant treatment of early stage non small cell lung cancer. *Curr Genomics*. 2008;9(4):252-262.
414. Ota S, Ishii G, Goto K, et al. Immunohistochemical expression of BCRP and ERCC1 in biopsy specimen predicts survival in advanced non-small-cell lung cancer treated with cisplatin-based chemotherapy. *Lung Cancer*. 2009;64(1):98-104.
415. Hwang IG, Jang JS, Do JH, et al. Different relation between ERCC1 overexpression and treatment outcomes of two platinum agents in advanced biliary tract adenocarcinoma patients. *Cancer Chemother Pharmacol*. 2011;68(4):935-944.

416. Ceppi P, Longo M, Volante M, et al. Excision repair cross complementing-1 and topoisomerase IIalpha gene expression in small-cell lung cancer patients treated with platinum and etoposide: a retrospective study. *J Thorac Oncol.* 2008;3(6):583-589.
417. Suk R, Gurubhagavatula S, Park S, et al. Polymorphisms in ERCC1 and grade 3 or 4 toxicity in non-small cell lung cancer patients. *Clin Cancer Res.* 2005;11(4):1534-1538.
418. Zhao H, Zhang H, Du Y, Gu X. Prognostic significance of BRCA1, ERCC1, RRM1, and RRM2 in patients with advanced non-small cell lung cancer receiving chemotherapy. *Tumour Biol.* 2014;35(12):12679-12688.
419. Azuma K, Komohara Y, Sasada T, et al. Excision repair cross-complementation group 1 predicts progression-free and overall survival in non-small cell lung cancer patients treated with platinum-based chemotherapy. *Cancer Sci.* 2007;98(9):1336-1343.
420. Li Z, Qing Y, Guan W, et al. Predictive value of APE1, BRCA1, ERCC1 and TUBB3 expression in patients with advanced non-small cell lung cancer (NSCLC) receiving first-line platinum-paclitaxel chemotherapy. *Cancer Chemother Pharmacol.* 2014;74(4):777-786.
421. Olaussen KA, Dunant A, Fouret P, et al. DNA repair by ERCC1 in non-small-cell lung cancer and cisplatin-based adjuvant chemotherapy. *N Engl J Med.* 2006;355(10):983-991.
422. Zhang L, Liu T, Zhang JQ. Relationship between the protein expression of ERCC1, BRCA, beta-tubulin and K-ras and the efficacy and prognosis in advanced non-small cell lung cancer. *Zhonghua Zhong Liu Za Zhi.* 2011;33(3):212-216.
423. Chen W and Bepler G. A method for biomarker directed survival prediction in advanced non-small-cell lung cancer patients treated with Carboplatin-based therapy. *J Pers Med.* 2013;3(3):10.3390/jpm3030251.
424. Ikeda S, Takabe K, Suzuki K. Expression of ERCC1 and class IIIbeta tubulin for predicting effect of carboplatin/paclitaxel in patients with advanced inoperable non-small cell lung cancer. *Pathol Int.* 2009;59(12):863-867.
425. Vilmar A and Sorensen JB. Excision repair cross-complementation group 1 (ERCC1) in platinum-based treatment of non-small cell lung cancer with special emphasis on carboplatin: a review of current literature. *Lung Cancer.* 2009;64(2):131-139.
426. Yang Y and Xian L. The association between the ERCC1/2 polymorphisms and the clinical outcomes of the platinum-based chemotherapy in non-small cell lung cancer (NSCLC): a systematic review and meta-analysis. *Tumour Biol.* 2014;35(4):2905-2921.
427. Xie KJ, He HE, Sun AJ, Liu XB, Sun LP, Dong XJ. Expression of ERCC1, MSH2 and PARP1 in non-small cell lung cancer and prognostic value in patients treated with platinum-based chemotherapy. *Asian Pac J Cancer Prev.* 2014;15(6):2591-2596.
428. Cubukcu E, Fatih Olmez O, Saraydaroglu O, et al. Immunohistochemical expression of excision repair cross-complementing 1 (ERCC1) in non-small-cell lung cancer: implications for patient outcome. *Clin Transl Oncol.* 2011;13(11):826-830.
429. Bisof V, Zajc Petranovic M, Rakusic Z, Samardzic KR, Juretic A. The prognostic and predictive value of excision repair cross-complementation group 1 (ERCC1) protein in 1288 patients with head and neck squamous cell carcinoma treated with platinum-based therapy: a meta-analysis. *Eur Arch Otorhinolaryngol.* 2015.
430. Jun HJ, Ahn MJ, Kim HS, et al. ERCC1 expression as a predictive marker of squamous cell carcinoma of the head and neck treated with cisplatin-based concurrent chemoradiation. *Br J Cancer.* 2008;99(1):167-172.
431. Bellmunt J, Paz-Ares L, Cuello M, et al. Gene expression of ERCC1 as a novel prognostic marker in advanced bladder cancer patients receiving cisplatin-based chemotherapy. *Ann Oncol.* 2007;18(3):522-528.
432. Li S, Wu J, Chen Y, et al. ERCC1 expression levels predict the outcome of platinum-based chemotherapies in advanced bladder cancer: a meta-analysis. *Anticancer Drugs.* 2014;25(1):106-114.
433. Zucali PA, Giovannetti E, Destro A, et al. Thymidylate synthase and excision repair cross-complementing group-1 as predictors of responsiveness in mesothelioma patients treated with pemetrexed/carboplatin. *Clin Cancer Res.* 2011;17(8):2581-2590.
434. Lee SH, Noh KB, Lee JS, et al. Thymidylate synthase and ERCC1 as predictive markers in patients with pulmonary adenocarcinoma treated with pemetrexed and cisplatin. *Lung Cancer.* 2013;81(1):102-108.
435. Sun JM, Ahn MJ, Park MJ, et al. Expression of excision repair cross-complementation group 1 as predictive marker for nasopharyngeal cancer treated with concurrent chemoradiotherapy. *Int J Radiat Oncol Biol Phys.* 2011;80(3):655-660.

436. Mancuso A, Sacchetta S, Saletti PC, et al. Clinical and molecular determinants of survival in pancreatic cancer patients treated with second-line chemotherapy: results of an Italian/Swiss multicenter survey. *Anticancer Res.* 2010;30(10):4289-4295.
437. Fareed KR, Al-Attar A, Soomro IN, et al. Tumour regression and ERCC1 nuclear protein expression predict clinical outcome in patients with gastro-oesophageal cancer treated with neoadjuvant chemotherapy. *Br J Cancer.* 2010;102(11):1600-1607.
438. Joshi MB, Shiota Y, Danenberg KD, et al. High gene expression of TS1, GSTP1, and ERCC1 are risk factors for survival in patients treated with trimodality therapy for esophageal cancer. *Clin Cancer Res.* 2005;11(6):2215-2221.
439. Ronchi CL, Sbiera S, Kraus L, et al. Expression of excision repair cross complementing group 1 and prognosis in adrenocortical carcinoma patients treated with platinum-based chemotherapy. *Endocr Relat Cancer.* 2009;16(3):907-918.
440. Zheng DL, Tang GD, Chen YN, Zhang T, Qin MB. Genetic variability of ERCC1 and ERCC2 genes involved in the nucleotide excision repair pathway influences the treatment outcome of gastric cancer. *Genet Mol Res.* 2016;15(2):10.4238/gmr.15027384.
441. Mo J, Luo M, Cui J, Zhou S. Prognostic value of ERCC1 and ERCC2 gene polymorphisms in patients with gastric cancer receiving platinum-based chemotherapy. *Int J Clin Exp Pathol.* 2015;8(11):15065-15071.
442. Bai Y, Wang L, Li G, Fang X, Li Y, Yang S. Genetic variability of ERCC1 genes in NER pathway influences the treatment outcome of gastric cancer. *Int J Clin Exp Pathol.* 2015;8(10):13367-13373.
443. Kalikaki A, Voutsina A, Koutsopoulos A, et al. ERCC1 SNPs as Potential Predictive Biomarkers in Non-Small Cell Lung Cancer Patients Treated With Platinum-Based Chemotherapy. *Cancer Invest.* 2015;33(4):107-113.
444. Takenaka T, Yano T, Kiyohara C, et al. Effects of excision repair cross-complementation group 1 (ERCC1) single nucleotide polymorphisms on the prognosis of non-small cell lung cancer patients. *Lung Cancer.* 2010;67(1):101-107.
445. Zhou W, Gurubhagavatula S, Liu G, et al. Excision repair cross-complementation group 1 polymorphism predicts overall survival in advanced non-small cell lung cancer patients treated with platinum-based chemotherapy. *Clin Cancer Res.* 2004;10(15):4939-4943.
446. Chen WH, Xin PL, Pan QX, et al. ERCC1 single nucleotide polymorphism C8092A, but not its expression is associated with survival of esophageal squamous cell carcinoma patients from Fujian province, China. *PLoS One.* 2014;9(9):e106600.
447. Li Y, Hu P, Cao Y, Wang GY, Wang N, Zhou RM. Predicting the outcome of platinum-based chemotherapies in epithelial ovarian cancer using the 8092C/A polymorphism of ERCC1: a meta-analysis. *Biomarkers.* 2014;19(2):128-134.
448. Moxley KM, Benbrook DM, Queimado L, et al. The role of single nucleotide polymorphisms of the ERCC1 and MMS19 genes in predicting platinum-sensitivity, progression-free and overall survival in advanced epithelial ovarian cancer. *Gynecol Oncol.* 2013;130(2):377-382.
449. Krivak TC, Darcy KM, Tian C, et al. Relationship between ERCC1 polymorphisms, disease progression, and survival in the Gynecologic Oncology Group Phase III Trial of intraperitoneal versus intravenous cisplatin and paclitaxel for stage III epithelial ovarian cancer. *J Clin Oncol.* 2008;26(21):3598-3606.
450. Erculj N, Kovac V, Hmeljak J, Dolzan V. The influence of platinum pathway polymorphisms on the outcome in patients with malignant mesothelioma. *Ann Oncol.* 2012;23(4):961-967.
451. Krawczyk P, Wojas-Krawczyk K, Mlak R, Kucharczyk T, Biernacka B, Milanowski J. Predictive value of ERCC1 single-nucleotide polymorphism in patients receiving platinum-based chemotherapy for locally-advanced and advanced non-small cell lung cancer--a pilot study. *Folia Histochem Cytobiol.* 2012;50(1):80-86.
452. Mazzoni F, Cecere FL, Meoni G, et al. Phase II trial of customized first line chemotherapy according to ERCC1 and RRM1 SNPs in patients with advanced non-small-cell lung cancer. *Lung Cancer.* 2013;82(2):288-293.
453. Smith S, Su D, Rigault de la Longrais IA, et al. ERCC1 genotype and phenotype in epithelial ovarian cancer identify patients likely to benefit from paclitaxel treatment in addition to platinum-based therapy. *J Clin Oncol.* 2007;25(33):5172-5179.
454. Su D, Ma S, Liu P, et al. Genetic polymorphisms and treatment response in advanced non-small cell lung cancer. *Lung Cancer.* 2007;56(2):281-288.

455. Yan L, Shu-Ying Y, Shan K, et al. Association between polymorphisms of ERCC1 and survival in epithelial ovarian cancer patients with chemotherapy. *Pharmacogenomics*. 2012;13(4):419-427.
456. Krivak TC, Darcy KM, Tian C, et al. Single nucleotide polymorphisms in ERCC1 are associated with disease progression, and survival in patients with advanced stage ovarian and primary peritoneal carcinoma; a Gynecologic Oncology Group study. *Gynecol Oncol*. 2011;122(1):121-126.
457. Hao T, Feng W, Zhang J, Sun YJ, Wang G. Association of four ERCC1 and ERCC2 SNPs with survival of bone tumour patients. *Asian Pac J Cancer Prev*. 2012;13(8):3821-3824.
458. Chai H, Pan J, Zhang X, et al. ERCC1 C118T associates with response to FOLFOX4 chemotherapy in colorectal cancer patients in Han Chinese. *Int J Clin Exp Med*. 2012;5(2):186-194.
459. Reed E. ERCC1 and clinical resistance to platinum-based therapy. *Clin Cancer Res*. 2005;11(17):6100-6102.
460. Li F, Xie X, Ren X, Zhang J. A meta-analysis identifies ERCC1 gene polymorphism as a predictor of better patient response to treatment with radiochemotherapy. *Cancer Chemother Pharmacol*. 2016;77(6):1183-1191.
461. Zhao X, Zhang Z, Yuan Y, Yuan X. Polymorphisms in ERCC1 gene could predict clinical outcome of platinum-based chemotherapy for non-small cell lung cancer patients. *Tumour Biol*. 2014;35(8):8335-8341.
462. Xu TP, Shen H, Liu LX, Shu YQ. Association of ERCC1-C118T and -C8092A polymorphisms with lung cancer risk and survival of advanced-stage non-small cell lung cancer patients receiving platinum-based chemotherapy: a pooled analysis based on 39 reports. *Gene*. 2013;526(2):265-274.
463. Vaezi A, Feldman CH, Niedernhofer LJ. ERCC1 and XRCC1 as biomarkers for lung and head and neck cancer. *Pharmacogenomics Pers Med*. 2011;4:47-63.
464. Zhou C, Ren S, Zhou S, et al. Predictive effects of ERCC1 and XRCC3 SNP on efficacy of platinum-based chemotherapy in advanced NSCLC patients. *Jpn J Clin Oncol*. 2010;40(10):954-960.
465. Li F, Sun X, Sun N, et al. Association between polymorphisms of ERCC1 and XPD and clinical response to platinum-based chemotherapy in advanced non-small cell lung cancer. *Am J Clin Oncol*. 2010;33(5):489-494.
466. Wei SZ, Zhan P, Shi MQ, et al. Predictive value of ERCC1 and XPD polymorphism in patients with advanced non-small cell lung cancer receiving platinum-based chemotherapy: a systematic review and meta-analysis. *Med Oncol*. 2011;28(1):315-321.
467. Okuda K, Sasaki H, Hikosaka Y, et al. Excision repair cross complementation group 1 polymorphisms predict overall survival after platinum-based chemotherapy for completely resected non-small-cell lung cancer. *J Surg Res*. 2011;168(2):206-212.
468. Kalikaki A, Kanaki M, Vassalou H, et al. DNA repair gene polymorphisms predict favorable clinical outcome in advanced non-small-cell lung cancer. *Clin Lung Cancer*. 2009;10(2):118-123.
469. Huang SJ, Wang YF, Jin ZY, Sun JY, Guo ZL. Role of ERCC1 variants in response to chemotherapy and clinical outcome of advanced non-small cell lung cancer. *Tumour Biol*. 2014;35(5):4023-4029.
470. Isla D, Sarries C, Rosell R, et al. Single nucleotide polymorphisms and outcome in docetaxel-cisplatin-treated advanced non-small-cell lung cancer. *Ann Oncol*. 2004;15(8):1194-1203.
471. Kalikaki A, Koutsopoulos A, Hatzidaki D, et al. Clinical outcome of patients with non-small cell lung cancer receiving front-line chemotherapy according to EGFR and K-RAS mutation status. *Lung Cancer*. 2010;69(1):110-115.
472. Palomba G, Atzori F, Budroni M, et al. ERCC1 polymorphisms as prognostic markers in T4 breast cancer patients treated with platinum-based chemotherapy. *J Transl Med*. 2014;12:272-014-0272-4.
473. Nikitas N, Karadimou A, Tsitoura E, et al. Association of ERCC1 SNPs with outcome in platinum-treated patients with advanced urothelial cancer: a Hellenic Cooperative Oncology Group study. *Pharmacogenomics*. 2012;13(14):1595-1607.
474. Yu D, Zhang X, Liu J, et al. Characterization of functional excision repair cross-complementation group 1 variants and their association with lung cancer risk and prognosis. *Clin Cancer Res*. 2008;14(9):2878-2886.
475. Gu AQ, Wang WM, Chen WY, Shi CL, Lu JH, Han JQ. XRCC1 genetic polymorphisms and sensitivity to platinum-based drugs in non-small cell lung cancer: an update meta-analysis based on 4708 subjects. *Int J Clin Exp Med*. 2015;8(1):145-154.
476. Li L, Wan C, Wen FQ. Polymorphisms in the XRCC1 gene are associated with treatment response to platinum chemotherapy in advanced non-small cell lung cancer patients based on meta-analysis. *Genet Mol Res*. 2014;13(2):3772-3786.

477. Peng Y, Li Z, Zhang S, et al. Association of DNA base excision repair genes (OGG1, APE1 and XRCC1) polymorphisms with outcome to platinum-based chemotherapy in advanced nonsmall-cell lung cancer patients. *Int J Cancer*. 2014;135(11):2687-2696.
478. Wu J, Liu J, Zhou Y, et al. Predictive value of XRCC1 gene polymorphisms on platinum-based chemotherapy in advanced non-small cell lung cancer patients: a systematic review and meta-analysis. *Clin Cancer Res*. 2012;18(14):3972-3981.
479. Liao WY, Shih JY, Chang GC, et al. Genetic polymorphism of XRCC1 Arg399Gln is associated with survival in non-small-cell lung cancer patients treated with gemcitabine/platinum. *J Thorac Oncol*. 2012;7(6):973-981.
480. Cui Z, Yin Z, Li X, Wu W, Guan P, Zhou B. Association between polymorphisms in XRCC1 gene and clinical outcomes of patients with lung cancer: a meta-analysis. *BMC Cancer*. 2012;12:71-2407-12-71.
481. Osawa T, Davies D, Hartley JA. Mechanism of cell death resulting from DNA interstrand cross-linking in mammalian cells. *Cell Death Dis*. 2011;2:e187.
482. Sun X, Li F, Sun N, et al. Polymorphisms in XRCC1 and XPG and response to platinum-based chemotherapy in advanced non-small cell lung cancer patients. *Lung Cancer*. 2009;65(2):230-236.
483. Giachino DF, Ghio P, Regazzoni S, et al. Prospective assessment of XPD Lys751Gln and XRCC1 Arg399Gln single nucleotide polymorphisms in lung cancer. *Clin Cancer Res*. 2007;13(10):2876-2881.
484. Li D, Zhou Q, Liu Y, Yang Y, Li Q. DNA repair gene polymorphism associated with sensitivity of lung cancer to therapy. *Med Oncol*. 2012;29(3):1622-1628.
485. Xu J, Ma J, Zong HT, Wang SY, Zhou JW. Pharmacogenetic role of XRCC1 polymorphisms on the clinical outcome of gastric cancer patients with platinum-based chemotherapy: a systematic review and meta-analysis. *Genet Mol Res*. 2014;13(1):1438-1446.
486. Cao Z, Song J, Wang J, Guo X, Yu S, Dong W. Association between polymorphisms in XRCC1 gene and treatment outcomes of patients with advanced gastric cancer: a systematic review and meta-analysis. *PLoS One*. 2014;9(1):e85357.
487. Wu H, Xu C, Chen G, Wang J. X-ray repair cross-complementing 1 polymorphism and prognosis of platinum-based chemotherapy in gastric and colorectal cancer: a meta-analysis. *J Gastroenterol Hepatol*. 2014;29(5):926-933.
488. Liu B, Wei J, Zou Z, et al. Polymorphism of XRCC1 predicts overall survival of gastric cancer patients receiving oxaliplatin-based chemotherapy in Chinese population. *Eur J Hum Genet*. 2007;15(10):1049-1053.
489. Stoeckelmacher J, Ghaderi V, Iobal S, et al. A polymorphism of the XRCC1 gene predicts for response to platinum based treatment in advanced colorectal cancer. *Anticancer Res*. 2001;21(4B):3075-3079.
490. Li K and Li W. Association between polymorphisms of XRCC1 and ADPRT genes and ovarian cancer survival with platinum-based chemotherapy in Chinese population. *Mol Cell Biochem*. 2013;372(1-2):27-33.
491. Cheng CX, Xue M, Li K, Li WS. Predictive value of XRCC1 and XRCC3 gene polymorphisms for risk of ovarian cancer death after chemotherapy. *Asian Pac J Cancer Prev*. 2012;13(6):2541-2545.
492. Shuai HL, Luo X, Yan RL, Li J, Chen DL. XRCC1 polymorphisms are associated with cervical cancer risk and response to chemotherapy: a systematic review and meta-analysis. *Asian Pac J Cancer Prev*. 2012;13(12):6423-6427.
493. Cheng XD, Lu WG, Ye F, Wan XY, Xie X. The association of XRCC1 gene single nucleotide polymorphisms with response to neoadjuvant chemotherapy in locally advanced cervical carcinoma. *J Exp Clin Cancer Res*. 2009;28:91-9966-28-91.
494. Chung HH, Kim MK, Kim JW, et al. XRCC1 R399Q polymorphism is associated with response to platinum-based neoadjuvant chemotherapy in bulky cervical cancer. *Gynecol Oncol*. 2006;103(3):1031-1037.
495. Kim K, Kang SB, Chung HH, Kim JW, Park NH, Song YS. XRCC1 Arginine194Tryptophan and GGH-401Cytosine/Thymine polymorphisms are associated with response to platinum-based neoadjuvant chemotherapy in cervical cancer. *Gynecol Oncol*. 2008;111(3):509-515.
496. Chen J, Wang Z, Zou T, et al. Pharmacogenomics of platinum-based chemotherapy response in NSCLC: a genotyping study and a pooled analysis. *Oncotarget*. 2016.
497. Yuan Z, Li J, Hu R, Jiao Y, Han Y, Weng Q. Predictive assessment in pharmacogenetics of XRCC1 gene on clinical outcomes of advanced lung cancer patients treated with platinum-based chemotherapy. *Sci Rep*. 2015;5:16482.

498. Fan X and Xiu Q. Effect of X-ray repair cross complementing group 1 polymorphisms on the efficacy of platinum-based chemotherapy in patients with nonsmall cell lung cancer. *J Cancer Res Ther.* 2015;11(3):571-574.
499. Giovannetti E, Pacetti P, Reni M, et al. Association between DNA-repair polymorphisms and survival in pancreatic cancer patients treated with combination chemotherapy. *Pharmacogenomics.* 2011;12(12):1641-1652.
500. Wang R, Wang W, Zhang JW. Correlation between XRCC2 and XRCC5 single nucleotide polymorphisms and drug-sensitivity of human lung cancer cells. *Zhonghua Yi Xue Za Zhi.* 2008;88(43):3059-3062.
501. Chen J, Wu L, Wang Y, et al. Effect of transporter and DNA repair gene polymorphisms to lung cancer chemotherapy toxicity. *Tumour Biol.* 2016;37(2):2275-2284.
502. Shen XY, Lu FZ, Wu Y, Zhao LT, Lin ZF. XRCC3 Thr241Met polymorphism and clinical outcomes of NSCLC patients receiving platinum-based chemotherapy: a systematic review and meta-analysis. *PLoS One.* 2013;8(8):e69553.
503. Xu C, Wang X, Zhang Y, Li L. Effect of the XRCC1 and XRCC3 genetic polymorphisms on the efficacy of platinum-based chemotherapy in patients with advanced non-small cell lung cancer. *Zhongguo Fei Ai Za Zhi.* 2011;14(12):912-917.
504. Ye J, Chu T, Li R, et al. Pol zeta polymorphisms are associated with platinum based chemotherapy response and side effects among non-small cell lung cancer patients. *Neoplasma.* 2015;62(5):833-839.
505. Iwatsuki M, Mimori K, Yokobori T, et al. A platinum agent resistance gene, POLB, is a prognostic indicator in colorectal cancer. *J Surg Oncol.* 2009;100(3):261-266.
506. Zhang X, Chen Q, Chen J, et al. Association of polymorphisms in translesion synthesis genes with prognosis of advanced non-small-cell lung cancer patients treated with platinum-based chemotherapy. *J Surg Oncol.* 2016;113(1):17-23.
507. Hertz DL. Germline pharmacogenetics of paclitaxel for cancer treatment. *Pharmacogenomics.* 2013;14(9):1065-1084.
508. Travis LB, Fossa SD, Sessa HD, et al. Chemotherapy-induced peripheral neurotoxicity and ototoxicity: new paradigms for translational genomics. *J Natl Cancer Inst.* 2014;106(5):10.1093/jnci/dju044.
509. Roberts D, Schick J, Conway S, et al. Identification of genes associated with platinum drug sensitivity and resistance in human ovarian cancer cells. *Br J Cancer.* 2005;92(6):1149-1158.
510. Konstantinopoulos PA, Fountzilas E, Pillay K, et al. Carboplatin-induced gene expression changes in vitro are prognostic of survival in epithelial ovarian cancer. *BMC Med Genomics.* 2008;1:59-8794-1-59.
511. Ferriss JS, Kim Y, Duska L, et al. Multi-gene expression predictors of single drug responses to adjuvant chemotherapy in ovarian carcinoma: predicting platinum resistance. *PLoS One.* 2012;7(2):e30550.
512. Peters GJ, Van Moorsel CJ, Lakerveld B, et al. Effects of gemcitabine on cis-platinum-DNA adduct formation and repair in a panel of gemcitabine and cisplatin-sensitive or -resistant human ovarian cancer cell lines. *Int J Oncol.* 2006;28(1):237-244.
513. Nevedomskaya E, Perryman R, Solanki S, Syed N, Mayboroda OA, Keun HC. A Systems Oncology Approach Identifies NT5E as a Key Metabolic Regulator in Tumor Cells and Modulator of Platinum Sensitivity. *J Proteome Res.* 2016;15(1):280-290.
514. Cao S, Wang S, Ma H, et al. Genome-wide association study of myelosuppression in non-small-cell lung cancer patients with platinum-based chemotherapy. *Pharmacogenomics J.* 2016;16(1):41-46.
515. Cao S, Wang C, Ma H, et al. Genome-wide Association Study on Platinum-induced Hepatotoxicity in Non-Small Cell Lung Cancer Patients. *Sci Rep.* 2015;5:11556.
516. LaCroix B, Gamazon ER, Lenkala D, et al. Integrative analyses of genetic variation, epigenetic regulation, and the transcriptome to elucidate the biology of platinum sensitivity. *BMC Genomics.* 2014;15:292-2164-15-292.
517. Hu L, Wu C, Zhao X, et al. Genome-wide association study of prognosis in advanced non-small cell lung cancer patients receiving platinum-based chemotherapy. *Clin Cancer Res.* 2012;18(19):5507-5514.
518. Ziliak D, O'Donnell PH, Im HK, et al. Germline polymorphisms discovered via a cell-based, genome-wide approach predict platinum response in head and neck cancers. *Transl Res.* 2011;157(5):265-272.
519. Wu X, Ye Y, Rosell R, et al. Genome-wide association study of survival in non-small cell lung cancer patients receiving platinum-based chemotherapy. *J Natl Cancer Inst.* 2011;103(10):817-825.
520. Li D, Wei L, Xu B, et al. Association of GWAS-identified lung cancer susceptibility loci with survival length in patients with small-cell lung cancer treated with platinum-based chemotherapy. *PLoS One.* 2014;9(11):e113574.

521. Wu C, Xu B, Yuan P, et al. Genome-wide interrogation identifies YAP1 variants associated with survival of small-cell lung cancer patients. *Cancer Res.* 2010;70(23):9721-9729.
522. Tan XL, Moyer AM, Fridley BL, et al. Genetic variation predicting cisplatin cytotoxicity associated with overall survival in lung cancer patients receiving platinum-based chemotherapy. *Clin Cancer Res.* 2011;17(17):5801-5811.
523. French JD, Johnatty SE, Lu Y, et al. Germline polymorphisms in an enhancer of PSIP1 are associated with progression-free survival in epithelial ovarian cancer. *Oncotarget.* 2016;7(6):6353-6368.
524. Xu H, Robinson GW, Huang J, et al. Common variants in ACYP2 influence susceptibility to cisplatin-induced hearing loss. *Nat Genet.* 2015;47(3):263-266.
525. International HapMap Consortium. The International HapMap Project. *Nature.* 2003;426(6968):789-796.
526. O'Donnell PH, Gamazon E, Zhang W, et al. Population differences in platinum toxicity as a means to identify novel genetic susceptibility variants. *Pharmacogenet Genomics.* 2010;20(5):327-337.
527. Huang RS, Johnatty SE, Gamazon ER, et al. Platinum sensitivity-related germline polymorphism discovered via a cell-based approach and analysis of its association with outcome in ovarian cancer patients. *Clin Cancer Res.* 2011;17(16):5490-5500.
528. Wheeler HE, Gamazon ER, Stark AL, et al. Genome-wide meta-analysis identifies variants associated with platinating agent susceptibility across populations. *Pharmacogenomics J.* 2013;13(1):35-43.
529. Wani MC, Taylor HL, Wall ME, Coggon P, McPhail AT. Plant antitumor agents. VI. The isolation and structure of taxol, a novel antileukemic and antitumor agent from *Taxus brevifolia*. *J Am Chem Soc.* 1971;93(9):2325-2327.
530. Schiff PB, Fant J, Horwitz SB. Promotion of microtubule assembly in vitro by taxol. *Nature.* 1979;277(5698):665-667.
531. de Weger VA, Beijnen JH, Schellens JH. Cellular and clinical pharmacology of the taxanes docetaxel and paclitaxel--a review. *Anticancer Drugs.* 2014;25(5):488-494.
532. Jordan MA and Wilson L. Microtubules as a target for anticancer drugs. *Nat Rev Cancer.* 2004;4(4):253-265.
533. Kumar N. Taxol-induced polymerization of purified tubulin. Mechanism of action. *J Biol Chem.* 1981;256(20):10435-10441.
534. Hamel E, del Campo AA, Lowe MC, Lin CM. Interactions of taxol, microtubule-associated proteins, and guanine nucleotides in tubulin polymerization. *J Biol Chem.* 1981;256(22):11887-11894.
535. De Brabander M, Geuens G, Nuydens R, Willebrords R, De Mey J. Taxol induces the assembly of free microtubules in living cells and blocks the organizing capacity of the centrosomes and kinetochores. *Proc Natl Acad Sci U S A.* 1981;78(9):5608-5612.
536. Nogales E, Wolf SG, Khan IA, Luduena RF, Downing KH. Structure of tubulin at 6.5 Å and location of the taxol-binding site. *Nature.* 1995;375(6530):424-427.
537. Schiff PB and Horwitz SB. Taxol stabilizes microtubules in mouse fibroblast cells. *Proc Natl Acad Sci U S A.* 1980;77(3):1561-1565.
538. Dostal V and Libusova L. Microtubule drugs: action, selectivity, and resistance across the kingdoms of life. *Protoplasma.* 2014;251(5):991-1005.
539. Sherman-Baust CA, Becker KG, Wood Iii WH, Zhang Y, Morin PJ. Gene expression and pathway analysis of ovarian cancer cells selected for resistance to cisplatin, paclitaxel, or doxorubicin. *J Ovarian Res.* 2011;4(1):21-2215-4-21.
540. Weiss RB, Donehower RC, Wiernik PH, et al. Hypersensitivity reactions from taxol. *J Clin Oncol.* 1990;8(7):1263-1268.
541. McGuire WP, Hoskins WJ, Brady MF, et al. Cyclophosphamide and cisplatin compared with paclitaxel and cisplatin in patients with stage III and stage IV ovarian cancer. *N Engl J Med.* 1996;334(1):1-6.
542. McGuire WP, Hoskins WJ, Brady MF, et al. Cyclophosphamide and cisplatin versus paclitaxel and cisplatin: a phase III randomized trial in patients with suboptimal stage III/IV ovarian cancer (from the Gynecologic Oncology Group). *Semin Oncol.* 1996;23(5 Suppl 12):40-47.
543. Piccart MJ, Bertelsen K, James K, et al. Randomized intergroup trial of cisplatin-paclitaxel versus cisplatin-cyclophosphamide in women with advanced epithelial ovarian cancer: three-year results. *J Natl Cancer Inst.* 2000;92(9):699-708.
544. Joshi M, Liu X, Belani CP. Taxanes, past, present, and future impact on non-small cell lung cancer. *Anticancer Drugs.* 2014;25(5):571-583.

545. Kampan NC, Madondo MT, McNally OM, Quinn M, Plebanski M. Paclitaxel and Its Evolving Role in the Management of Ovarian Cancer. *Biomed Res Int*. 2015;2015:413076.
546. Nogales E. Structural insights into microtubule function. *Annu Rev Biochem*. 2000;69:277-302.
547. Rahman A, Korzekwa KR, Grogan J, Gonzalez FJ, Harris JW. Selective biotransformation of taxol to 6 alpha-hydroxytaxol by human cytochrome P450 2C8. *Cancer Res*. 1994;54(21):5543-5546.
548. Czejka M, Schueller J, Schnait H, Springer B, Eder I. Clinical pharmacokinetics and metabolism of paclitaxel after a three-hour infusion: comparison of two preparations. *Journal of Oncology Pharmacy Practice*. 2003;9(4):129-139.
549. Cresteil T, Monsarrat B, Dubois J, Sonnier M, Alvinerie P, Gueritte F. Regioselective metabolism of taxoids by human CYP3A4 and 2C8: structure-activity relationship. *Drug Metab Dispos*. 2002;30(4):438-445.
550. Lagas JS, Vlaming ML, van Tellingen O, et al. Multidrug resistance protein 2 is an important determinant of paclitaxel pharmacokinetics. *Clin Cancer Res*. 2006;12(20 Pt 1):6125-6132.
551. Sparreboom A, van Tellingen O, Nooijen WJ, Beijnen JH. Tissue distribution, metabolism and excretion of paclitaxel in mice. *Anticancer Drugs*. 1996;7(1):78-86.
552. Sparreboom A, Huizing MT, Boesen JJ, Nooijen WJ, van Tellingen O, Beijnen JH. Isolation, purification, and biological activity of mono- and dihydroxylated paclitaxel metabolites from human feces. *Cancer Chemother Pharmacol*. 1995;36(4):299-304.
553. Malingre MM, Terwogt JM, Beijnen JH, et al. Phase I and pharmacokinetic study of oral paclitaxel. *J Clin Oncol*. 2000;18(12):2468-2475.
554. Kuh HJ, Jang SH, Wientjes MG, Weaver JR, Au JL. Determinants of paclitaxel penetration and accumulation in human solid tumor. *J Pharmacol Exp Ther*. 1999;290(2):871-880.
555. Smith NF, Marsh S, Scott-Horton TJ, et al. Variants in the SLCO1B3 gene: interethnic distribution and association with paclitaxel pharmacokinetics. *Clin Pharmacol Ther*. 2007;81(1):76-82.
556. Smith NF, Acharya MR, Desai N, Figg WD, Sparreboom A. Identification of OATP1B3 as a high-affinity hepatocellular transporter of paclitaxel. *Cancer Biol Ther*. 2005;4(8):815-818.
557. Kobayashi Y, Ohshiro N, Sakai R, Ohbayashi M, Kohyama N, Yamamoto T. Transport mechanism and substrate specificity of human organic anion transporter 2 (hOat2 [SLC22A7]). *J Pharm Pharmacol*. 2005;57(5):573-578.
558. Schellens JH, Malingre MM, Kruijtz CM, et al. Modulation of oral bioavailability of anticancer drugs: from mouse to man. *Eur J Pharm Sci*. 2000;12(2):103-110.
559. Szakacs G, Paterson JK, Ludwig JA, Booth-Genthe C, Gottesman MM. Targeting multidrug resistance in cancer. *Nat Rev Drug Discov*. 2006;5(3):219-234.
560. Fellner S, Bauer B, Miller DS, et al. Transport of paclitaxel (Taxol) across the blood-brain barrier in vitro and in vivo. *J Clin Invest*. 2002;110(9):1309-1318.
561. NicAmhlaoibh R, Heenan M, Cleary I, et al. Altered expression of mRNAs for apoptosis-modulating proteins in a low level multidrug resistant variant of a human lung carcinoma cell line that also expresses *mdr1* mRNA. *Int J Cancer*. 1999;82(3):368-376.
562. Yeh JJ, Hsu WH, Wang JJ, Ho ST, Kao A. Predicting chemotherapy response to paclitaxel-based therapy in advanced non-small-cell lung cancer with P-glycoprotein expression. *Respiration*. 2003;70(1):32-35.
563. Penson RT, Oliva E, Skates SJ, et al. Expression of multidrug resistance-1 protein inversely correlates with paclitaxel response and survival in ovarian cancer patients: a study in serial samples. *Gynecol Oncol*. 2004;93(1):98-106.
564. Alberts D, Taetle R, Frutiger E, et al. Multidrug resistance expression as an independent risk factor for poor survival in optimal stage III ovarian cancer. In: Anonymous PROCEEDINGS OF THE ANNUAL MEETING-AMERICAN SOCIETY OF CLINICAL ONCOLOGY. AMERICAN SOCIETY OF CLINICAL ONCOLOGY; 1998:1356-1356.
565. Yague E, Higgins CF, Raguz S. Complete reversal of multidrug resistance by stable expression of small interfering RNAs targeting MDR1. *Gene Ther*. 2004;11(14):1170-1174.
566. Borst P, Evers R, Kool M, Wijnholds J. A family of drug transporters: the multidrug resistance-associated proteins. *J Natl Cancer Inst*. 2000;92(16):1295-1302.
567. Allen JD, Brinkhuis RF, van Deemter L, Wijnholds J, Schinkel AH. Extensive contribution of the multidrug transporters P-glycoprotein and Mrp1 to basal drug resistance. *Cancer Res*. 2000;60(20):5761-5766.

568. Brooks TA, Minderman H, O'Loughlin KL, et al. Taxane-based reversal agents modulate drug resistance mediated by P-glycoprotein, multidrug resistance protein, and breast cancer resistance protein. *Mol Cancer Ther.* 2003;2(11):1195-1205.
569. Huisman MT, Chhatta AA, van Tellingen O, Beijnen JH, Schinkel AH. MRP2 (ABCC2) transports taxanes and confers paclitaxel resistance and both processes are stimulated by probenecid. *Int J Cancer.* 2005;116(5):824-829.
570. Green H, Soderkvist P, Rosenberg P, et al. Pharmacogenetic studies of Paclitaxel in the treatment of ovarian cancer. *Basic Clin Pharmacol Toxicol.* 2009;104(2):130-137.
571. Sissung TM, Mross K, Steinberg SM, et al. Association of ABCB1 genotypes with paclitaxel-mediated peripheral neuropathy and neutropenia. *Eur J Cancer.* 2006;42(17):2893-2896.
572. Leskela S, Jara C, Leandro-Garcia LJ, et al. Polymorphisms in cytochromes P450 2C8 and 3A5 are associated with paclitaxel neurotoxicity. *Pharmacogenomics J.* 2011;11(2):121-129.
573. DeLoia JA, Zamboni WC, Jones JM, Strychor S, Kelley JL, Gallion HH. Expression and activity of taxane-metabolizing enzymes in ovarian tumors. *Gynecol Oncol.* 2008;108(2):355-360.
574. Dai D, Zeldin DC, Blaisdell JA, et al. Polymorphisms in human CYP2C8 decrease metabolism of the anticancer drug paclitaxel and arachidonic acid. *Pharmacogenetics.* 2001;11(7):597-607.
575. Harris JW, Katki A, Anderson LW, Chmurny GN, Paukstelis JV, Collins JM. Isolation, structural determination, and biological activity of 6 alpha-hydroxytaxol, the principal human metabolite of taxol. *J Med Chem.* 1994;37(5):706-709.
576. Monsarrat B, Alvinerie P, Wright M, et al. Hepatic metabolism and biliary excretion of Taxol in rats and humans. *J Natl Cancer Inst Monogr.* 1993;(15)(15):39-46.
577. Ghose R, Mallick P, Taneja G, Chu C, Moorthy B. In Vitro Approaches to Study Regulation of Hepatic Cytochrome P450 (CYP) 3A Expression by Paclitaxel and Rifampicin. *Methods Mol Biol.* 2016;1395:55-68.
578. Orr GA, Verdier-Pinard P, McDaid H, Horwitz SB. Mechanisms of Taxol resistance related to microtubules. *Oncogene.* 2003;22(47):7280-7295.
579. Wosnitzer MS, Domingo-Domenech J, Castillo-Martin M, et al. Predictive value of microtubule associated proteins tau and stathmin in patients with nonmuscle invasive bladder cancer receiving adjuvant intravesical taxane therapy. *J Urol.* 2011;186(5):2094-2100.
580. Sun R, Liu Z, Wang L, et al. Overexpression of stathmin is resistant to paclitaxel treatment in patients with non-small cell lung cancer. *Tumour Biol.* 2015;36(9):7195-7204.
581. Yu J, Gao J, Lu Z, et al. Combination of microtubule associated protein-tau and beta-tubulin III predicts chemosensitivity of paclitaxel in patients with advanced gastric cancer. *Eur J Cancer.* 2014;50(13):2328-2335.
582. Xie S, Ogden A, Aneja R, Zhou J. Microtubule-Binding Proteins as Promising Biomarkers of Paclitaxel Sensitivity in Cancer Chemotherapy. *Med Res Rev.* 2016;36(2):300-312.
583. Sun X, Shi X, Liu M, et al. Mdp3 is a novel microtubule-binding protein that regulates microtubule assembly and stability. *Cell Cycle.* 2011;10(22):3929-3937.
584. Tala, Xie S, Sun X, et al. Microtubule-associated protein Mdp3 promotes breast cancer growth and metastasis. *Theranostics.* 2014;4(10):1052-1061.
585. Tala, Sun X, Chen J, et al. Microtubule stabilization by Mdp3 is partially attributed to its modulation of HDAC6 in addition to its association with tubulin and microtubules. *PLoS One.* 2014;9(3):e90932.
586. Drewes G, Ebnet A, Preuss U, Mandelkow EM, Mandelkow E. MARK, a novel family of protein kinases that phosphorylate microtubule-associated proteins and trigger microtubule disruption. *Cell.* 1997;89(2):297-308.
587. Bauer JA, Chakravarthy AB, Rosenbluth JM, et al. Identification of markers of taxane sensitivity using proteomic and genomic analyses of breast tumors from patients receiving neoadjuvant paclitaxel and radiation. *Clin Cancer Res.* 2010;16(2):681-690.
588. Wagner P, Wang B, Clark E, Lee H, Rouzier R, Pusztai L. Microtubule Associated Protein (MAP)-Tau: a novel mediator of paclitaxel sensitivity in vitro and in vivo. *Cell Cycle.* 2005;4(9):1149-1152.
589. Rouzier R, Rajan R, Wagner P, et al. Microtubule-associated protein tau: a marker of paclitaxel sensitivity in breast cancer. *Proc Natl Acad Sci U S A.* 2005;102(23):8315-8320.
590. Pusztai L. Markers predicting clinical benefit in breast cancer from microtubule-targeting agents. *Ann Oncol.* 2007;18 Suppl 12:xiii15-20.

591. Andre F, Hatzis C, Anderson K, et al. Microtubule-associated protein-tau is a bifunctional predictor of endocrine sensitivity and chemotherapy resistance in estrogen receptor-positive breast cancer. *Clin Cancer Res.* 2007;13(7):2061-2067.
592. Bonneau C, Gurard-Levin ZA, Andre F, Pusztai L, Rouzier R. Predictive and Prognostic Value of the TauProtein in Breast Cancer. *Anticancer Res.* 2015;35(10):5179-5184.
593. Hait WN and Yang JM. The individualization of cancer therapy: the unexpected role of p53. *Trans Am Clin Climatol Assoc.* 2006;117:85-101; discussion 101.
594. Zhang CC, Yang JM, White E, Murphy M, Levine A, Hait WN. The role of MAP4 expression in the sensitivity to paclitaxel and resistance to vinca alkaloids in p53 mutant cells. *Oncogene.* 1998;16(12):1617-1624.
595. Marklund U, Larsson N, Gradin HM, Brattsand G, Gullberg M. Oncoprotein 18 is a phosphorylation-responsive regulator of microtubule dynamics. *EMBO J.* 1996;15(19):5290-5298.
596. Amayed P, Pantaloni D, Carlier MF. The effect of stathmin phosphorylation on microtubule assembly depends on tubulin critical concentration. *J Biol Chem.* 2002;277(25):22718-22724.
597. Alli E, Yang JM, Ford JM, Hait WN. Reversal of stathmin-mediated resistance to paclitaxel and vinblastine in human breast carcinoma cells. *Mol Pharmacol.* 2007;71(5):1233-1240.
598. Mistry SJ, Bank A, Atweh GF. Synergistic antiangiogenic effects of stathmin inhibition and taxol exposure. *Mol Cancer Res.* 2007;5(8):773-782.
599. Alli E, Bash-Babula J, Yang JM, Hait WN. Effect of stathmin on the sensitivity to antimicrotubule drugs in human breast cancer. *Cancer Res.* 2002;62(23):6864-6869.
600. Iancu C, Mistry SJ, Arkin S, Wallenstein S, Atweh GF. Effects of stathmin inhibition on the mitotic spindle. *J Cell Sci.* 2001;114(Pt 5):909-916.
601. Bieche I, Lachkar S, Becette V, et al. Overexpression of the stathmin gene in a subset of human breast cancer. *Br J Cancer.* 1998;78(6):701-709.
602. Melhem R, Hailat N, Kuick R, Hanash SM. Quantitative analysis of Op18 phosphorylation in childhood acute leukemia. *Leukemia.* 1997;11(10):1690-1695.
603. Su D, Smith SM, Preti M, et al. Stathmin and tubulin expression and survival of ovarian cancer patients receiving platinum treatment with and without paclitaxel. *Cancer.* 2009;115(11):2453-2463.
604. Murphy M, Hinman A, Levine AJ. Wild-type p53 negatively regulates the expression of a microtubule-associated protein. *Genes Dev.* 1996;10(23):2971-2980.
605. Lavarino C, Pilotti S, Oggionni M, et al. P53 Gene Status and Response to Platinum/paclitaxel-Based Chemotherapy in Advanced Ovarian Carcinoma. *J Clin Oncol.* 2000;18(23):3936-3945.
606. Luo Y, Li D, Ran J, et al. End-binding protein 1 stimulates paclitaxel sensitivity in breast cancer by promoting its actions toward microtubule assembly and stability. *Protein Cell.* 2014;5(6):469-479.
607. Thomas GE, Sreeja JS, Gireesh KK, Gupta H, Manna TK. +TIP EB1 downregulates paclitaxel-induced proliferation inhibition and apoptosis in breast cancer cells through inhibition of paclitaxel binding on microtubules. *Int J Oncol.* 2015;46(1):133-146.
608. Ganguly A, Yang H, Pedroza M, Bhattacharya R, Cabral F. Mitotic centromere-associated kinesin (MCAK) mediates paclitaxel resistance. *J Biol Chem.* 2011;286(42):36378-36384.
609. Ganguly A, Yang H, Cabral F. Overexpression of mitotic centromere-associated Kinesin stimulates microtubule detachment and confers resistance to paclitaxel. *Mol Cancer Ther.* 2011;10(6):929-937.
610. Wang H, Liu B, Zhang C, et al. Parkin regulates paclitaxel sensitivity in breast cancer via a microtubule-dependent mechanism. *J Pathol.* 2009;218(1):76-85.
611. Fetz V, Bier C, Habtemichael N, et al. Inducible NO synthase confers chemoresistance in head and neck cancer by modulating survivin. *Int J Cancer.* 2009;124(9):2033-2041.
612. Lu J, Tan M, Huang WC, et al. Mitotic deregulation by survivin in ErbB2-overexpressing breast cancer cells contributes to Taxol resistance. *Clin Cancer Res.* 2009;15(4):1326-1334.
613. Sudo T, Nitta M, Saya H, Ueno NT. Dependence of paclitaxel sensitivity on a functional spindle assembly checkpoint. *Cancer Res.* 2004;64(7):2502-2508.
614. Javeed A, Ashraf M, Riaz A, Ghafoor A, Afzal S, Mukhtar MM. Paclitaxel and immune system. *European Journal of Pharmaceutical Sciences.* 2009;38(4):283-290.
615. Ferlini C, Cicchillitti L, Raspaglio G, et al. Paclitaxel directly binds to Bcl-2 and functionally mimics activity of Nur77. *Cancer Res.* 2009;69(17):6906-6914.
616. Agarwal R and Kaye SB. Ovarian cancer: strategies for overcoming resistance to chemotherapy. *Nat Rev Cancer.* 2003;3(7):502-516.

617. Komlodi-Pasztor E, Sackett D, Wilkerson J, Fojo T. Mitosis is not a key target of microtubule agents in patient tumors. *Nat Rev Clin Oncol*. 2011;8(4):244-250.
618. Mitchison TJ. The proliferation rate paradox in antimitotic chemotherapy. *Mol Biol Cell*. 2012;23(1):1-6.
619. Oshiro C, Marsh S, McLeod H, Carrillo MW, Klein T, Altman R. Taxane pathway. *Pharmacogenet Genomics*. 2009;19(12):979-983.
620. Park HS, Lim SM, Shin HJ, et al. Pharmacogenetic analysis of advanced non-small-cell lung cancer patients treated with first-line paclitaxel and carboplatin chemotherapy. *Pharmacogenet Genomics*. 2016;26(3):116-125.
621. Mechetner E, Kyshtoobayeva A, Zonis S, et al. Levels of multidrug resistance (MDR1) P-glycoprotein expression by human breast cancer correlate with in vitro resistance to taxol and doxorubicin. *Clin Cancer Res*. 1998;4(2):389-398.
622. Johnatty SE, Beesley J, Gao B, et al. ABCB1 (MDR1) polymorphisms and ovarian cancer progression and survival: a comprehensive analysis from the Ovarian Cancer Association Consortium and The Cancer Genome Atlas. *Gynecol Oncol*. 2013;131(1):8-14.
623. Yakirevich E, Sabo E, Naroditsky I, Sova Y, Lavie O, Resnick MB. Multidrug resistance-related phenotype and apoptosis-related protein expression in ovarian serous carcinomas. *Gynecol Oncol*. 2006;100(1):152-159.
624. Ludwig AH and Kupryjanczyk J. Does MDR-1 G2677T/A polymorphism really associate with ovarian cancer response to paclitaxel chemotherapy? *Clin Cancer Res*. 2006;12(20 Pt 1):6204.
625. Leschziner GD, Andrew T, Pirmohamed M, Johnson MR. ABCB1 genotype and PGP expression, function and therapeutic drug response: a critical review and recommendations for future research. *Pharmacogenomics J*. 2007;7(3):154-179.
626. Hamidovic A, Hahn K, Kolesar J. Clinical significance of ABCB1 genotyping in oncology. *J Oncol Pharm Pract*. 2010;16(1):39-44.
627. Subhani S, Jamil K, Nirni SS. Association of MDR1 gene (C3435T) polymorphism and gene expression profiling in lung cancer patients treated with platinum-based chemotherapy. *Mol Diagn Ther*. 2015;19(5):289-297.
628. Johnatty SE, Beesley J, Paul J, et al. ABCB1 (MDR 1) polymorphisms and progression-free survival among women with ovarian cancer following paclitaxel/carboplatin chemotherapy. *Clin Cancer Res*. 2008;14(17):5594-5601.
629. Gao B, Russell A, Beesley J, et al. Paclitaxel sensitivity in relation to ABCB1 expression, efflux and single nucleotide polymorphisms in ovarian cancer. *Sci Rep*. 2014;4:4669.
630. Bergmann TK, Brasch-Andersen C, Green H, et al. Impact of ABCB1 variants on neutrophil depression: a pharmacogenomic study of paclitaxel in 92 women with ovarian cancer. *Basic Clin Pharmacol Toxicol*. 2012;110(2):199-204.
631. Meng X, Wang G, Liu P, et al. ATP-binding cassette B1 gene polymorphisms, mRNA expression and chemosensitivity to paclitaxel in non-small cell lung cancer cells. *Respirology*. 2011;16(8):1228-1234.
632. Johnatty SE, Tyrer JP, Kar S, et al. Genome-wide Analysis Identifies Novel Loci Associated with Ovarian Cancer Outcomes: Findings from the Ovarian Cancer Association Consortium. *Clin Cancer Res*. 2015;21(23):5264-5276.
633. Vella N, Aiello M, Russo AE, et al. 'Genetic profiling' and ovarian cancer therapy (review). *Mol Med Rep*. 2011;4(5):771-777.
634. Grimm C, Polterauer S, Zeillinger R, et al. Two multidrug-resistance (ABCB1) gene polymorphisms as prognostic parameters in women with ovarian cancer. *Anticancer Res*. 2010;30(9):3487-3491.
635. Nakajima M, Fujiki Y, Kyo S, et al. Pharmacokinetics of paclitaxel in ovarian cancer patients and genetic polymorphisms of CYP2C8, CYP3A4, and MDR1. *J Clin Pharmacol*. 2005;45(6):674-682.
636. Chang H, Rha SY, Jeung HC, et al. Association of the ABCB1 gene polymorphisms 2677G>T/A and 3435C>T with clinical outcomes of paclitaxel monotherapy in metastatic breast cancer patients. *Ann Oncol*. 2009;20(2):272-277.
637. Tecza K, Pamula-Pilat J, Kolosza Z, Radlak N, Grzybowska E. Genetic polymorphisms and gene-dosage effect in ovarian cancer risk and response to paclitaxel/cisplatin chemotherapy. *J Exp Clin Cancer Res*. 2015;34:2-015-0124-y.
638. Green H, Soderkvist P, Rosenberg P, Horvath G, Peterson C. ABCB1 G1199A polymorphism and ovarian cancer response to paclitaxel. *J Pharm Sci*. 2008;97(6):2045-2048.

639. Hasmats J, Kupersmidt I, Rodriguez-Antona C, et al. Identification of candidate SNPs for drug induced toxicity from differentially expressed genes in associated tissues. *Gene*. 2012;506(1):62-68.
640. Niemi M, Leathart JB, Neuvonen M, Backman JT, Daly AK, Neuvonen PJ. Polymorphism in CYP2C8 is associated with reduced plasma concentrations of repaglinide. *Clin Pharmacol Ther*. 2003;74(4):380-387.
641. Rodriguez-Antona C, Niemi M, Backman JT, et al. Characterization of novel CYP2C8 haplotypes and their contribution to paclitaxel and repaglinide metabolism. *Pharmacogenomics J*. 2008;8(4):268-277.
642. Bergmann TK, Brasch-Andersen C, Green H, et al. Impact of CYP2C8*3 on paclitaxel clearance: a population pharmacokinetic and pharmacogenomic study in 93 patients with ovarian cancer. *Pharmacogenomics J*. 2011;11(2):113-120.
643. Green H, Khan MS, Jakobsen-Falk I, Avall-Lundqvist E, Peterson C. Impact of CYP3A5*3 and CYP2C8-HapC on paclitaxel/carboplatin-induced myelosuppression in patients with ovarian cancer. *J Pharm Sci*. 2011;100(10):4205-4209.
644. Hu L, Lv QL, Guo Y, et al. Genetic variation of CYP3A5 influences paclitaxel/carboplatin-induced toxicity in Chinese epithelial ovarian cancer patients. *J Clin Pharmacol*. 2016;56(3):349-354.
645. Assis J, Pereira D, Gomes M, et al. Influence of CYP3A4 genotypes in the outcome of serous ovarian cancer patients treated with first-line chemotherapy: implication of a CYP3A4 activity profile. *Int J Clin Exp Med*. 2013;6(7):552-561.
646. Nakajima Y, Yoshitani T, Fukushima-Uesaka H, et al. Impact of the haplotype CYP3A4*16B harboring the Thr185Ser substitution on paclitaxel metabolism in Japanese patients with cancer. *Clin Pharmacol Ther*. 2006;80(2):179-191.
647. Zhang HL, Ruan L, Zheng LM, Whyte D, Tzeng CM, Zhou XW. Association between class III beta-tubulin expression and response to paclitaxel/vinorelbine-based chemotherapy for non-small cell lung cancer: a meta-analysis. *Lung Cancer*. 2012;77(1):9-15.
648. Kelley MJ, Li S, Harpole DH. Genetic analysis of the beta-tubulin gene, TUBB, in non-small-cell lung cancer. *J Natl Cancer Inst*. 2001;93(24):1886-1888.
649. Leandro-Garcia LJ, Leskela S, Jara C, et al. Regulatory polymorphisms in beta-tubulin IIa are associated with paclitaxel-induced peripheral neuropathy. *Clin Cancer Res*. 2012;18(16):4441-4448.
650. Powrozek T, Mlak R, Krawczyk P, et al. Retrospective analysis of second-line chemotherapy outcomes with paclitaxel or docetaxel in correlation with STMN1 polymorphism in advanced non-small cell lung cancer patients. *Clin Transl Oncol*. 2016;18(1):33-39.
651. Santibanez M, Gallardo D, Morales F, et al. The MAD1 1673 G --> A polymorphism alters the function of the mitotic spindle assembly checkpoint and is associated with a worse response to induction chemotherapy and sensitivity to treatment in patients with advanced epithelial ovarian cancer. *Pharmacogenet Genomics*. 2013;23(4):190-199.
652. Pellicciotta I, Yang CP, Venditti CA, Goldberg GL, Shahabi S. Response to microtubule-interacting agents in primary epithelial ovarian cancer cells. *Cancer Cell Int*. 2013;13(1):33-2867-13-33.
653. Malisic E, Jankovic R, Jakovljevic K, Radulovic S. TP53 gene status and human papilloma virus infection in response to platinum plus taxane-based chemotherapy of epithelial ovarian carcinomas. *J BUON*. 2011;16(4):701-707.
654. Gadducci A, Di Cristofano C, Zavaglia M, et al. P53 gene status in patients with advanced serous epithelial ovarian cancer in relation to response to paclitaxel- plus platinum-based chemotherapy and long-term clinical outcome. *Anticancer Res*. 2006;26(1B):687-693.
655. Santos AM, Sousa H, Portela C, et al. TP53 and P21 polymorphisms: response to cisplatin/paclitaxel-based chemotherapy in ovarian cancer. *Biochem Biophys Res Commun*. 2006;340(1):256-262.
656. Vikhanskaya F, Siddique MM, Kei Lee M, Brogini M, Sabapathy K. Evaluation of the combined effect of p53 codon 72 polymorphism and hotspot mutations in response to anticancer drugs. *Clin Cancer Res*. 2005;11(12):4348-4356.
657. Lee SY, Kang HG, Yoo SS, et al. Polymorphisms in DNA repair and apoptosis-related genes and clinical outcomes of patients with non-small cell lung cancer treated with first-line paclitaxel-cisplatin chemotherapy. *Lung Cancer*. 2013;82(2):330-339.
658. Lee SY, Jung DK, Choi JE, et al. PD-L1 polymorphism can predict clinical outcomes of non-small cell lung cancer patients treated with first-line paclitaxel-cisplatin chemotherapy. *Sci Rep*. 2016;6:25952.
659. Do SK, Lee SY, Choi JE, Hong MJ, Lee JH, Park JY. 155P: PD-L1 polymorphism can predict clinical outcomes of non-small cell lung cancer patients treated with first-line paclitaxel-cisplatin chemotherapy. *J Thorac Oncol*. 2016;11(4 Suppl):S125-0864(16)30265-9. Epub 2016 Apr 15.

660. Liu D, Xu W, Zhang ZW, et al. RB1 polymorphism contributes to the efficacy of platinum-taxanes in advanced squamous cell lung cancer. *Asian Pac J Cancer Prev*. 2015;16(2):775-781.
661. Niu N, Schaid DJ, Abo RP, et al. Genetic association with overall survival of taxane-treated lung cancer patients - a genome-wide association study in human lymphoblastoid cell lines followed by a clinical association study. *BMC Cancer*. 2012;12:422-2407-12-422.
662. Komatsu M, Wheeler HE, Chung S, et al. Pharmacoeethnicity in Paclitaxel-Induced Sensory Peripheral Neuropathy. *Clin Cancer Res*. 2015;21(19):4337-4346.
663. Sato Y, Yamamoto N, Kunitoh H, et al. Genome-wide association study on overall survival of advanced non-small cell lung cancer patients treated with carboplatin and paclitaxel. *J Thorac Oncol*. 2011;6(1):132-138.
664. Hong MJ, Lee SY, Choi JE, et al. A genetic variation in microRNA target site of ETS2 is associated with clinical outcomes of paclitaxel-cisplatin chemotherapy in non-small cell lung cancer. *Oncotarget*. 2016;7(13):15948-15958.
665. DiNucci EM. Energy healing: a complementary treatment for orthopaedic and other conditions. *Orthop Nurs*. 2005;24(4):259-269.
666. Lucas A and Cole TJ. Breast milk and neonatal necrotising enterocolitis. *Lancet*. 1990;336(8730):1519-1523.
667. Stessman HA, Baughn LB, Sarver A, et al. Profiling bortezomib resistance identifies secondary therapies in a mouse myeloma model. *Mol Cancer Ther*. 2013.
668. Varma RR, Hector SM, Clark K, Greco WR, Hawthorn L, Pendyala L. Gene expression profiling of a clonal isolate of oxaliplatin-resistant ovarian carcinoma cell line A2780/C10. *Oncol Rep*. 2005;14(4):925-932.
669. Arioka H, Nishio K, Ishida T, et al. Enhancement of cisplatin sensitivity in high mobility group 2 cDNA-transfected human lung cancer cells. *Jpn J Cancer Res*. 1999;90(1):108-115.

Chapter 2

1. Varma RR, Hector SM, Clark K, Greco WR, Hawthorn L, Pendyala L. Gene expression profiling of a clonal isolate of oxaliplatin-resistant ovarian carcinoma cell line A2780/C10. *Oncol Rep.* 2005;14(4):925-932.
2. So PS, Jiang Y, Qin Y. Touch therapies for pain relief in adults. *Cochrane Database Syst Rev.* 2008;(4):CD006535. doi(4):CD006535.
3. Siegel RL, Miller KD, Jemal A. Cancer statistics, 2015. *CA Cancer J Clin.* 2015;65(1):5-29.
4. Chen VW, Ruiz B, Killeen JL, Cote TR, Wu XC, Correa CN. Pathology and classification of ovarian tumors. *Cancer.* 2003;97(10 Suppl):2631-2642.
5. Molldrem JJ. Preparing basic and translational grant proposals: thoughts from the trenches. *Hematology Am Soc Hematol Educ Program.* 2010;2010:181-184.
6. Wheate NJ, Walker S, Craig GE, Oun R. The status of platinum anticancer drugs in the clinic and in clinical trials. *Dalton Trans.* 2010;39(35):8113-8127.
7. Rabik CA and Dolan ME. Molecular mechanisms of resistance and toxicity associated with platinating agents. *Cancer Treat Rev.* 2007;33(1):9-23.
8. Zorbas H and Keppler BK. Cisplatin damage: are DNA repair proteins saviors or traitors to the cell? *Chembiochem.* 2005;6(7):1157-1166.
9. Sprowl JA, Ness RA, Sparreboom A. Polymorphic transporters and platinum pharmacodynamics. *Drug Metab Pharmacokinet.* 2013;28(1):19-27.
10. Marsh S, McLeod H, Dolan E, et al. Platinum pathway. *Pharmacogenet Genomics.* 2009;19(7):563-564.
11. Jordan MA and Wilson L. Microtubules as a target for anticancer drugs. *Nat Rev Cancer.* 2004;4(4):253-265.
12. Huizing MT, Misser VH, Pieters RC, et al. Taxanes: a new class of antitumor agents. *Cancer Invest.* 1995;13(4):381-404.
13. McGrogan BT, Gilmartin B, Carney DN, McCann A. Taxanes, microtubules and chemoresistant breast cancer. *Biochim Biophys Acta.* 2008;1785(2):96-132.
14. Dai D, Zeldin DC, Blaisdell JA, et al. Polymorphisms in human CYP2C8 decrease metabolism of the anticancer drug paclitaxel and arachidonic acid. *Pharmacogenetics.* 2001;11(7):597-607.
15. Cresteil T, Monsarrat B, Dubois J, Sonnier M, Alvinerie P, Gueritte F. Regioselective metabolism of taxoids by human CYP3A4 and 2C8: structure-activity relationship. *Drug Metab Dispos.* 2002;30(4):438-445.
16. Marsh S. Pharmacogenomics of taxane/platinum therapy in ovarian cancer. *Int J Gynecol Cancer.* 2009;19 Suppl 2:S30-4.
17. Rustin GJ, van der Burg ME, Griffin CL, et al. Early versus delayed treatment of relapsed ovarian cancer (MRC OV05/EORTC 55955): a randomised trial. *Lancet.* 2010;376(9747):1155-1163.
18. Kramer MS, Platt RW, Wen SW, et al. A new and improved population-based Canadian reference for birth weight for gestational age. *Pediatrics.* 2001;108(2):E35.
19. Miller G. Immortalization of human lymphocytes by Epstein-Barr virus. *Yale J Biol Med.* 1982;55(3-4):305-310.
20. Huang RS, Johnatty SE, Gamazon ER, et al. Platinum sensitivity-related germline polymorphism discovered via a cell-based approach and analysis of its association with outcome in ovarian cancer patients. *Clin Cancer Res.* 2011;17(16):5490-5500.
21. International HapMap Consortium. The International HapMap Project. *Nature.* 2003;426(6968):789-796.
22. Njiaju UO, Gamazon ER, Gorsic LK, et al. Whole-genome studies identify solute carrier transporters in cellular susceptibility to paclitaxel. *Pharmacogenet Genomics.* 2012;22(7):498-507.
23. 1000 Genomes Project Consortium, Auton A, Brooks LD, et al. A global reference for human genetic variation. *Nature.* 2015;526(7571):68-74.
24. Wu X, Ye Y, Rosell R, et al. Genome-wide association study of survival in non-small cell lung cancer patients receiving platinum-based chemotherapy. *J Natl Cancer Inst.* 2011;103(10):817-825.
25. Niu N, Schaid DJ, Abo RP, et al. Genetic association with overall survival of taxane-treated lung cancer patients - a genome-wide association study in human lymphoblastoid cell lines followed by a clinical association study. *BMC Cancer.* 2012;12:422-2407-12-422.
26. Li L, Fridley BL, Kalari K, et al. Discovery of genetic biomarkers contributing to variation in drug response of cytidine analogues using human lymphoblastoid cell lines. *BMC Genomics.* 2014;15:93-2164-15-93.

27. Fridley BL, Ghosh TM, Wang A, et al. Genome-Wide Study of Response to Platinum, Taxane, and Combination Therapy in Ovarian Cancer: In vitro Phenotypes, Inherited Variation, and Disease Recurrence. *Front Genet.* 2016;7:37.
28. Pritchard JK, Stephens M, Rosenberg NA, Donnelly P. Association mapping in structured populations. *Am J Hum Genet.* 2000;67(1):170-181.
29. Howie B, Fuchsberger C, Stephens M, Marchini J, Abecasis GR. Fast and accurate genotype imputation in genome-wide association studies through pre-phasing. *Nat Genet.* 2012;44(8):955-959.
30. Hartigan JA and Wong MA. Algorithm AS 136: A k-means clustering algorithm. *Journal of the Royal Statistical Society. Series C (Applied Statistics).* 1979;28(1):100-108.
31. Chou TC. Drug combination studies and their synergy quantification using the Chou-Talalay method. *Cancer Res.* 2010;70(2):440-446.
32. Kartalou M and Essigmann JM. Mechanisms of resistance to cisplatin. *Mutat Res.* 2001;478(1-2):23-43.
33. Ganzzinelli M, Mariani P, Cattaneo D, et al. Expression of DNA repair genes in ovarian cancer samples: biological and clinical considerations. *Eur J Cancer.* 2011;47(7):1086-1094.
34. Kim SH, Lee GW, Lee MJ, et al. Clinical significance of ERCC2 haplotype-tagging single nucleotide polymorphisms in patients with unresectable non-small cell lung cancer treated with first-line platinum-based chemotherapy. *Lung Cancer.* 2012;77(3):578-584.
35. Surowiak P, Materna V, Kaplenko I, et al. Augmented expression of metallothionein and glutathione S-transferase pi as unfavourable prognostic factors in cisplatin-treated ovarian cancer patients. *Virchows Arch.* 2005;447(3):626-633.
36. Zhang Q, Li K, Xu JH, et al. Role of ABCG2 expression driven by cisplatin in platinum-containing chemotherapy for gastric cancer. *World J Gastroenterol.* 2013;19(39):6630-6636.
37. Warta R, Theile D, Mogler C, et al. Association of drug transporter expression with mortality and progression-free survival in stage IV head and neck squamous cell carcinoma. *PLoS One.* 2014;9(9):e108908.
38. Tian C, Ambrosone CB, Darcy KM, et al. Common variants in ABCB1, ABCC2 and ABCG2 genes and clinical outcomes among women with advanced stage ovarian cancer treated with platinum and taxane-based chemotherapy: a Gynecologic Oncology Group study. *Gynecol Oncol.* 2012;124(3):575-581.
39. Thacker J and Zdzienicka MZ. The XRCC genes: expanding roles in DNA double-strand break repair. *DNA Repair (Amst).* 2004;3(8-9):1081-1090.
40. Chen J, Wu L, Wang Y, et al. Effect of transporter and DNA repair gene polymorphisms to lung cancer chemotherapy toxicity. *Tumour Biol.* 2016;37(2):2275-2284.
41. Martin A and Scharff MD. AID and mismatch repair in antibody diversification. *Nat Rev Immunol.* 2002;2(8):605-614.
42. Hait WN and Yang JM. The individualization of cancer therapy: the unexpected role of p53. *Trans Am Clin Climatol Assoc.* 2006;117:85-101; discussion 101.
43. Zhang CC, Yang JM, White E, Murphy M, Levine A, Hait WN. The role of MAP4 expression in the sensitivity to paclitaxel and resistance to vinca alkaloids in p53 mutant cells. *Oncogene.* 1998;16(12):1617-1624.
44. Pereira D, Assis J, Gomes M, Nogueira A, Medeiros R. Improvement of a predictive model in ovarian cancer patients submitted to platinum-based chemotherapy: implications of a GST activity profile. *Eur J Clin Pharmacol.* 2016;72(5):545-553.
45. Moyer AM, Sun Z, Batzler AJ, et al. Glutathione pathway genetic polymorphisms and lung cancer survival after platinum-based chemotherapy. *Cancer Epidemiol Biomarkers Prev.* 2010;19(3):811-821.
46. Lin K, Ye D, Xie X. Protein expression levels of excision repair cross-complementation group 1 and xeroderma pigmentosum D correlate with response to platinum-based chemotherapy in the patients with advanced epithelial ovarian cancer. *Int J Gynecol Cancer.* 2008;18(5):1007-1012.
47. Arienti C, Tesei A, Verdecchia GM, et al. Peritoneal carcinomatosis from ovarian cancer: chemosensitivity test and tissue markers as predictors of response to chemotherapy. *J Transl Med.* 2011;9:94-5876-9-94.
48. de Laat WL, Jaspers NG, Hoeijmakers JH. Molecular mechanism of nucleotide excision repair. *Genes Dev.* 1999;13(7):768-785.
49. Lee SY, Kang HG, Yoo SS, et al. Polymorphisms in DNA repair and apoptosis-related genes and clinical outcomes of patients with non-small cell lung cancer treated with first-line paclitaxel-cisplatin chemotherapy. *Lung Cancer.* 2013;82(2):330-339.

Chapter 3

1. Siegel RL, Miller KD, Jemal A. Cancer statistics, 2015. *CA Cancer J Clin.* 2015;65(1):5-29.
2. Marsh S. Pharmacogenomics of taxane/platinum therapy in ovarian cancer. *Int J Gynecol Cancer.* 2009;19 Suppl 2:S30-4.
3. Dekou V, Whincup P, Papacosta O, et al. The effect of the C677T and A1298C polymorphisms in the methylenetetrahydrofolate reductase gene on homocysteine levels in elderly men and women from the British regional heart study. *Atherosclerosis.* 2001;154(3):659-666.
4. Jordan MA and Wilson L. Microtubules as a target for anticancer drugs. *Nat Rev Cancer.* 2004;4(4):253-265.
5. Huizing MT, Misser VH, Pieters RC, et al. Taxanes: a new class of antitumor agents. *Cancer Invest.* 1995;13(4):381-404.
6. Hewett M, Oliver DE, Rubin DL, et al. PharmGKB: the Pharmacogenetics Knowledge Base. *Nucleic Acids Res.* 2002;30(1):163-165.
7. Rodriguez-Antona C. Pharmacogenomics of paclitaxel. *Pharmacogenomics.* 2010;11(5):621-623.
8. Peethambaram P, Fridley BL, Vierkant RA, et al. Polymorphisms in ABCB1 and ERCC2 associated with ovarian cancer outcome. *Int J Mol Epidemiol Genet.* 2011;2(2):185-195.
9. Johnatty SE, Beesley J, Gao B, et al. ABCB1 (MDR1) polymorphisms and ovarian cancer progression and survival: a comprehensive analysis from the Ovarian Cancer Association Consortium and The Cancer Genome Atlas. *Gynecol Oncol.* 2013;131(1):8-14.
10. White KL, Vierkant RA, Fogarty ZC, et al. Analysis of over 10,000 Cases finds no association between previously reported candidate polymorphisms and ovarian cancer outcome. *Cancer Epidemiol Biomarkers Prev.* 2013;22(5):987-992.
11. Li F, Fridley BL, Matimba A, et al. Ecto-5'-nucleotidase and thiopurine cellular circulation: association with cytotoxicity. *Drug Metab Dispos.* 2010;38(12):2329-2338.
12. Li L, Fridley BL, Kalari K, et al. Gemcitabine and cytosine arabinoside pharmacogenomics: genome-wide association and drug response biomarkers. *PLoS One.* 2009;4(11):e7765.
13. Li L, Fridley B, Kalari K, et al. Gemcitabine and cytosine arabinoside cytotoxicity: association with lymphoblastoid cell expression. *Cancer Res.* 2008;68(17):7050-7058.
14. Niu N, Qin Y, Fridley BL, et al. Radiation pharmacogenomics: a genome-wide association approach to identify radiation response biomarkers using human lymphoblastoid cell lines. *Genome Res.* 2010;20(11):1482-1492.
15. Huang RS, Johnatty SE, Gamazon ER, et al. Platinum sensitivity-related germline polymorphism discovered via a cell-based approach and analysis of its association with outcome in ovarian cancer patients. *Clin Cancer Res.* 2011;17(16):5490-5500.
16. Wu TY, Fridley BL, Jenkins GD, Batzler A, Wang L, Weinshilboum RM. Mycophenolic acid response biomarkers: a cell line model system-based genome-wide screen. *Int Immunopharmacol.* 2011;11(8):1057-1064.
17. Gamazon ER, Huang RS, Cox NJ, Dolan ME. Chemotherapeutic drug susceptibility associated SNPs are enriched in expression quantitative trait loci. *Proc Natl Acad Sci U S A.* 2010;107(20):9287-9292.
18. Abo R, Hebring S, Ji Y, et al. Merging pharmacometabolomics with pharmacogenomics using '1000 Genomes' single-nucleotide polymorphism imputation: selective serotonin reuptake inhibitor response pharmacogenomics. *Pharmacogenet Genomics.* 2012;22(4):247-253.
19. Fridley BL, Jenkins GD, Tsai YY, et al. Gene set analysis of survival following ovarian cancer implicates macrolide binding and intracellular signaling genes. *Cancer Epidemiol Biomarkers Prev.* 2012;21(3):529-536.
20. White KL, Schildkraut JM, Palmieri RT, et al. Ovarian cancer risk associated with inherited inflammation-related variants. *Cancer Res.* 2012;72(5):1064-1069.
21. Pharoah PD, Tsai YY, Ramus SJ, et al. GWAS meta-analysis and replication identifies three new susceptibility loci for ovarian cancer. *Nat Genet.* 2013;45(4):362-70, 370e1-2.
22. Shen H, Fridley BL, Song H, et al. Epigenetic analysis leads to identification of HNF1B as a subtype-specific susceptibility gene for ovarian cancer. *Nat Commun.* 2013;4:1628.
23. Pritchard JK, Stephens M, Donnelly P. Inference of population structure using multilocus genotype data. *Genetics.* 2000;155(2):945-959.
24. 1000 Genomes Project Consortium, Auton A, Brooks LD, et al. A global reference for human genetic variation. *Nature.* 2015;526(7571):68-74.

25. Howie B, Fuchsberger C, Stephens M, Marchini J, Abecasis GR. Fast and accurate genotype imputation in genome-wide association studies through pre-phasing. *Nat Genet.* 2012;44(8):955-959.
26. Kutner MH, Nachtsheim CJ, Neter J, Li W. *Applied linear statistical models.* McGraw-Hill Irwin New York; 2005.
27. Willer CJ, Li Y, Abecasis GR. METAL: fast and efficient meta-analysis of genomewide association scans. *Bioinformatics.* 2010;26(17):2190-2191.
28. Fridley BL, Jenkins GD, Batzler A, et al. Multivariate models to detect genomic signatures for a class of drugs: application to thiopurines pharmacogenomics. *Pharmacogenomics J.* 2012;12(2):105-110.
29. Bush WS, Dudek SM, Ritchie MD. Biofilter: a knowledge-integration system for the multi-locus analysis of genome-wide association studies. *Pac Symp Biocomput.* 2009:368-379.
30. Niu N, Schaid DJ, Abo RP, et al. Genetic association with overall survival of taxane-treated lung cancer patients - a genome-wide association study in human lymphoblastoid cell lines followed by a clinical association study. *BMC Cancer.* 2012;12:422-2407-12-422.
31. Fanjul-Fernandez M, Quesada V, Cabanillas R, et al. Cell-cell adhesion genes CTNNA2 and CTNNA3 are tumour suppressors frequently mutated in laryngeal carcinomas. *Nat Commun.* 2013;4:2531.
32. Haryono SJ, Datasena IG, Santosa WB, Mulyarahardja R, Sari K. A pilot genome-wide association study of breast cancer susceptibility loci in Indonesia. *Asian Pac J Cancer Prev.* 2015;16(6):2231-2235.
33. McGranahan N, Favero F, de Bruin EC, Birkbak NJ, Szallasi Z, Swanton C. Clonal status of actionable driver events and the timing of mutational processes in cancer evolution. *Sci Transl Med.* 2015;7(283):283ra54.
34. Mexal S, Berger R, Pearce L, et al. Regulation of a novel alphaN-catenin splice variant in schizophrenic smokers. *Am J Med Genet B Neuropsychiatr Genet.* 2008;147B(6):759-768.
35. Song C and Zhang H. TARV: tree-based analysis of rare variants identifying risk modifying variants in CTNNA2 and CNTNAP2 for alcohol addiction. *Genet Epidemiol.* 2014;38(6):552-559.
36. Du C, Huang T, Sun D, et al. CDH4 as a novel putative tumor suppressor gene epigenetically silenced by promoter hypermethylation in nasopharyngeal carcinoma. *Cancer Lett.* 2011;309(1):54-61.
37. Miotto E, Sabbioni S, Veronese A, et al. Frequent aberrant methylation of the CDH4 gene promoter in human colorectal and gastric cancer. *Cancer Res.* 2004;64(22):8156-8159.
38. Chen T, Long B, Ren G, et al. Protocadherin20 Acts as a Tumor Suppressor Gene: Epigenetic Inactivation in Nasopharyngeal Carcinoma. *J Cell Biochem.* 2015;116(8):1766-1775.
39. Lv J, Zhu P, Yang Z, et al. PCDH20 functions as a tumour-suppressor gene through antagonizing the Wnt/beta-catenin signalling pathway in hepatocellular carcinoma. *J Viral Hepat.* 2015;22(2):201-211.
40. Zhan Q, Huang RF, Liang XH, et al. FRAS1 knockdown reduces A549 cells migration and invasion through downregulation of FAK signaling. *Int J Clin Exp Med.* 2014;7(7):1692-1697.
41. Li L, Yoo H, Becker FF, Ali-Osman F, Chan JY. Identification of a brain- and reproductive-organs-specific gene responsive to DNA damage and retinoic acid. *Biochem Biophys Res Commun.* 1995;206(2):764-774.
42. Noordermeer SM, Wennemers M, Bergevoet SM, et al. Expression of the BRCA1 complex member BRE predicts disease free survival in breast cancer. *Breast Cancer Res Treat.* 2012;135(1):125-133.
43. Chui YL, Ma CH, Li W, et al. Anti-apoptotic protein BRE/BRCC45 attenuates apoptosis through maintaining the expression of caspase inhibitor XIAP in mouse Lewis lung carcinoma D122 cells. *Apoptosis.* 2014;19(5):829-840.
44. Langbein S, Szakacs O, Wilhelm M, et al. Alteration of the LRP1B gene region is associated with high grade of urothelial cancer. *Lab Invest.* 2002;82(5):639-643.
45. Sonoda I, Imoto I, Inoue J, et al. Frequent silencing of low density lipoprotein receptor-related protein 1B (LRP1B) expression by genetic and epigenetic mechanisms in esophageal squamous cell carcinoma. *Cancer Res.* 2004;64(11):3741-3747.
46. Nakagawa T, Pimkhaokham A, Suzuki E, Omura K, Inazawa J, Imoto I. Genetic or epigenetic silencing of low density lipoprotein receptor-related protein 1B expression in oral squamous cell carcinoma. *Cancer Sci.* 2006;97(10):1070-1074.
47. Prazeres H, Torres J, Rodrigues F, et al. Chromosomal, epigenetic and microRNA-mediated inactivation of LRP1B, a modulator of the extracellular environment of thyroid cancer cells. *Oncogene.* 2011;30(11):1302-1317.
48. Lu G, Zhang G, Zhang C, Chen C, Liu R. A study of 131iodine-labeling of histamine-indomethacin: its in vivo therapeutic effect and anti-tumor mechanisms in Lewis-bearing lung cancer. *Radiat Oncol.* 2013;8:74-717X-8-74.

49. Poduslo SE, Huang R, Spiro A,3rd. A genome screen of successful aging without cognitive decline identifies LRP1B by haplotype analysis. *Am J Med Genet B Neuropsychiatr Genet.* 2010;153B(1):114-119.
50. Gamazon ER, Lamba JK, Pounds S, et al. Comprehensive genetic analysis of cytarabine sensitivity in a cell-based model identifies polymorphisms associated with outcome in AML patients. *Blood.* 2013;121(21):4366-4376.

Chapter 4

1. World Health Organization. GLOBOCAN 2012: Estimated cancer incidence, mortality and prevalence worldwide in 2012. Lyon, France: International Agency for Research on Cancer.[Links]. 2014.
2. Howlader N, Noone A, Krapcho M, et al. SEER Cancer Statistics Review (CSR) 1975-2010. Bethesda: National Cancer Institute. 2014.
3. Groome PA, Bolejack V, Crowley JJ, et al. The IASLC Lung Cancer Staging Project: validation of the proposals for revision of the T, N, and M descriptors and consequent stage groupings in the forthcoming (seventh) edition of the TNM classification of malignant tumours. *J Thorac Oncol*. 2007;2(8):694-705.
4. Delbaldo C, Michiels S, Syz N, Soria JC, Le Chevalier T, Pignon JP. Benefits of adding a drug to a single-agent or a 2-agent chemotherapy regimen in advanced non-small-cell lung cancer: a meta-analysis. *JAMA*. 2004;292(4):470-484.
5. Rabik CA and Dolan ME. Molecular mechanisms of resistance and toxicity associated with platinating agents. *Cancer Treat Rev*. 2007;33(1):9-23.
6. Sprowl JA, Ness RA, Sparreboom A. Polymorphic transporters and platinum pharmacodynamics. *Drug Metab Pharmacokinet*. 2013;28(1):19-27.
7. Lu C, Spitz MR, Zhao H, et al. Association between glutathione S-transferase pi polymorphisms and survival in patients with advanced nonsmall cell lung carcinoma. *Cancer*. 2006;106(2):441-447.
8. Okcu MF, Selvan M, Wang LE, et al. Glutathione S-transferase polymorphisms and survival in primary malignant glioma. *Clin Cancer Res*. 2004;10(8):2618-2625.
9. Peters U, Preisler-Adams S, Hebeisen A, et al. Glutathione S-transferase genetic polymorphisms and individual sensitivity to the ototoxic effect of cisplatin. *Anticancer Drugs*. 2000;11(8):639-643.
10. Meijer C, Timmer A, De Vries EG, et al. Role of metallothionein in cisplatin sensitivity of germ-cell tumours. *Int J Cancer*. 2000;85(6):777-781.
11. van der Straaten T, Kweekel D, Tiller M, Bogaartz J, Guchelaar HJ. Multiplex pyrosequencing of two polymorphisms in DNA repair gene XRCC1. *J Mol Diagn*. 2006;8(4):444-448.
12. Sakano S, Wada T, Matsumoto H, et al. Single nucleotide polymorphisms in DNA repair genes might be prognostic factors in muscle-invasive bladder cancer patients treated with chemoradiotherapy. *Br J Cancer*. 2006;95(5):561-570.
13. Wheeler HE, Gamazon ER, Stark AL, et al. Genome-wide meta-analysis identifies variants associated with platinating agent susceptibility across populations. *Pharmacogenomics J*. 2013;13(1):35-43.
14. Marsh S, McLeod H, Dolan E, et al. Platinum pathway. *Pharmacogenet Genomics*. 2009;19(7):563-564.
15. Marsh S, Paul J, King CR, Gifford G, McLeod HL, Brown R. Pharmacogenetic assessment of toxicity and outcome after platinum plus taxane chemotherapy in ovarian cancer: the Scottish Randomised Trial in Ovarian Cancer. *J Clin Oncol*. 2007;25(29):4528-4535.
16. Green H, Soderkvist P, Rosenberg P, Horvath G, Peterson C. mdr-1 single nucleotide polymorphisms in ovarian cancer tissue: G2677T/A correlates with response to paclitaxel chemotherapy. *Clin Cancer Res*. 2006;12(3 Pt 1):854-859.
17. Kim HS, Kim MK, Chung HH, et al. Genetic polymorphisms affecting clinical outcomes in epithelial ovarian cancer patients treated with taxanes and platinum compounds: a Korean population-based study. *Gynecol Oncol*. 2009;113(2):264-269.
18. Oshiro C, Marsh S, McLeod H, Carrillo MW, Klein T, Altman R. Taxane pathway. *Pharmacogenet Genomics*. 2009;19(12):979-983.
19. Hewett M, Oliver DE, Rubin DL, et al. PharmGKB: the Pharmacogenetics Knowledge Base. *Nucleic Acids Res*. 2002;30(1):163-165.
20. Kimchi-Sarfaty C, Oh JM, Kim IW, et al. A "silent" polymorphism in the MDR1 gene changes substrate specificity. *Science*. 2007;315(5811):525-528.
21. Caronia D, Patino-Garcia A, Perez-Martinez A, et al. Effect of ABCB1 and ABCC3 polymorphisms on osteosarcoma survival after chemotherapy: a pharmacogenetic study. *PLoS One*. 2011;6(10):e26091.
22. Wu X, Gu J, Wu TT, et al. Genetic variations in radiation and chemotherapy drug action pathways predict clinical outcomes in esophageal cancer. *J Clin Oncol*. 2006;24(23):3789-3798.
23. Siegel D, Yan C, Ross D. NAD(P)H:quinone oxidoreductase 1 (NQO1) in the sensitivity and resistance to antitumor quinones. *Biochem Pharmacol*. 2012;83(8):1033-1040.
24. Traver RD, Siegel D, Beall HD, et al. Characterization of a polymorphism in NAD(P)H: quinone oxidoreductase (DT-diaphorase). *Br J Cancer*. 1997;75(1):69-75.

25. Kolesar JM, Dahlberg SE, Marsh S, et al. The NQO1*2/*2 polymorphism is associated with poor overall survival in patients following resection of stages II and IIIa non-small cell lung cancer. *Oncol Rep.* 2011;25(6):1765-1772.
26. Hertz DL, Motsinger-Reif AA, Drobish A, et al. CYP2C8*3 predicts benefit/risk profile in breast cancer patients receiving neoadjuvant paclitaxel. *Breast Cancer Res Treat.* 2012;134(1):401-410.
27. Bahadur N, Leathart JB, Mutch E, et al. CYP2C8 polymorphisms in Caucasians and their relationship with paclitaxel 6 α -hydroxylase activity in human liver microsomes. *Biochem Pharmacol.* 2002;64(11):1579-1589.
28. Soyama A, Saito Y, Hanioka N, et al. Non-synonymous single nucleotide alterations found in the CYP2C8 gene result in reduced in vitro paclitaxel metabolism. *Biol Pharm Bull.* 2001;24(12):1427-1430.
29. Imai Y, Nakane M, Kage K, et al. C421A polymorphism in the human breast cancer resistance protein gene is associated with low expression of Q141K protein and low-level drug resistance. *Mol Cancer Ther.* 2002;1(8):611-616.
30. Morisaki K, Robey RW, Ozvegy-Laczka C, et al. Single nucleotide polymorphisms modify the transporter activity of ABCG2. *Cancer Chemother Pharmacol.* 2005;56(2):161-172.
31. Karachaliou N, Papadaki C, Lagoudaki E, et al. Predictive value of BRCA1, ERCC1, ATP7B, PKM2, TOPOI, TOPOmicron-IIA, TOPOIIB and C-MYC genes in patients with small cell lung cancer (SCLC) who received first line therapy with cisplatin and etoposide. *PLoS One.* 2013;8(9):e74611.
32. Vassalou H, Stathopoulos E, Fiolitaki G, et al. Excision-repair-cross-complement-1 protein as a prognostic factor in patients with advanced non-small cell lung cancer treated with platinum-based first-line chemotherapy. *Lung Cancer.* 2013;82(2):324-329.
33. Xu TP, Shen H, Liu LX, Shu YQ. Association of ERCC1-C118T and -C8092A polymorphisms with lung cancer risk and survival of advanced-stage non-small cell lung cancer patients receiving platinum-based chemotherapy: a pooled analysis based on 39 reports. *Gene.* 2013;526(2):265-274.
34. Caronia D, Patino-Garcia A, Milne RL, et al. Common variations in ERCC2 are associated with response to cisplatin chemotherapy and clinical outcome in osteosarcoma patients. *Pharmacogenomics J.* 2009;9(5):347-353.
35. Fernandez-Rozadilla C, Cazier JB, Moreno V, et al. Pharmacogenomics in colorectal cancer: a genome-wide association study to predict toxicity after 5-fluorouracil or FOLFOX administration. *Pharmacogenomics J.* 2013;13(3):209-217.
36. Mazzoni F, Cecere FL, Meoni G, et al. Phase II trial of customized first line chemotherapy according to ERCC1 and RRM1 SNPs in patients with advanced non-small-cell lung cancer. *Lung Cancer.* 2013;82(2):288-293.